Numbers Sequences and Series

Lecture Notes

Dr. Silvio Fanzon

6 Dec 2024

Academic Year 2024/25

Department of Mathematics

University of Hull

Table of contents

We	lcon	ne g	;
	Revi)
	Digit	tal Notes	;
	Read	lings	;
1	Intro	oduction	7
2	Prel	iminaries 15	;
	2.1	Sets	5
	2.2	Logic)
	2.3	Operations on sets	7
		2.3.1 Infinite union and intersection)
		2.3.2 Complement	L
		2.3.3 Power set	2
		2.3.4 Product of sets	3
	2.4	Equivalence relation	3
	2.5	Order relation	ŝ
	2.6	Intervals)
	2.7	Functions	L
	2.8	Absolute value or Modulus	L
	2.9	Triangle inequality	ŝ
	2.10	Proofs in Mathematics	1
	2.11	Induction	L
		*	
3	Real	l Numbers 49)
	3.1	Fields)
	3.2	Ordered fields	3
	3.3	Cut Property)
	3.4	Supremum and infimum	Ś
	•	3.4.1 Upper bound, supremum, maximum	7
		3.4.2 Lower bound, infimum, minimum)
	3.5	Completeness	ł
	3.6	Equivalence of Completeness and Cut Property	ŝ
	J	Axioms of Real Numbers	2
	3.7		
	3.7 3.8	Special subsets of R	5
	3.7 3.8	Special subsets of \mathbb{R}	5
	3.7 3.8	Special subsets of \mathbb{R} 83.8.1Natural numbers83.8.2Principle of induction8	5

		3.8.4 Rational numbers					
4	Prop	erties of R 96					
	4.1	Archimedean Property					
	4.2	Nested Interval Property					
	4.3	Revisiting Sup and inf					
	4.4	Existence of k -th Roots					
	4.5	Density of Q in R \ldots					
	4.6	Cardinality					
_	Com	n Jaw Nium hawa					
5	Con	The field C					
	5.1	$\begin{array}{c} \text{Ine field } \mathbb{C} \\ \text{Division in } \mathbb{C} \\ \end{array}$					
		5.1.1 Division in \mathbb{C}					
		$5.1.2$ U is not ordered \ldots 137					
		5.1.3 Completeness of \mathbb{C}					
	5.2	Complex conjugates					
	5.3	The complex plane					
		5.3.1 Distance on $\mathbb C$					
		5.3.2 Properties of modulus					
	5.4	Polar coordinates					
	5.5	Exponential form					
	5.6	Fundamental Theorem of Algebra					
	5.7	Solving polynomial equations					
		5.7.1 Quadratic polynomials					
		5.7.2 Higher order polynomials					
	5.8	Roots of unity					
	5.9	Roots in \mathbb{C}					
~	S						
0	Sequ	Ences in R 173					
	6.1	Definition of sequence					
	6.2	Convergent sequences					
	6.3	Divergent sequences					
	6.4	Uniqueness of limit 182					
	6.5	Bounded sequences					
	6.6	Algebra of limits					
	6.7	Fractional powers					
	6.8	Limit Tests					
		6.8.1 Squeeze Theorem					
		6.8.2 Geometric sequences					
		6.8.3 Ratio Test					
	6.9	Monotone sequences					
		6.9.1 Example: Euler's Number					
	6.10	Special limits					
7	Sequences in C 228						
	7.1	Algebra of limits in $\mathbb C$					

	7.2	Conve	rgence to zero	231			
	7.3	Geome	etric sequence Test and Ratio Test in $\mathbb C$	234			
	7.4	Conve	rgence of real and imaginary part	237			
8	Seri	es		241			
	8.1	Conve	rgent series	242			
	8.2	Geome	etric series	248			
	8.3	Algebi	a of Limits for Series	254			
	8.4	Non-n	egative series	258			
		8.4.1	Cauchy Condensation Test	259			
		8.4.2	Comparison Test	267			
		8.4.3	Limit Comparison Test	270			
		8.4.4	Ratio Test for positive series	276			
	8.5	Genera	al series	280			
		8.5.1	Absolute Convergence Test	280			
		8.5.2	Ratio Test for general series	284			
		8.5.3	Exponential function and Euler's Number	286			
		8.5.4	Conditional convergence	287			
		8.5.5	Dirichlet and Alternating Series Tests	289			
		8.5.6	Abel's Test	294			
Li	License 295						
	Reus	е		295			
	Cita	tion		295			
References							

Welcome

These are the Lecture Notes of **Numbers Sequences and Series 400297** for 2024/25 at the University of Hull. I will use this material during lectures. If you have any question or find any typo, please email me at

S.Fanzon@hull.ac.uk

Up to date information about the course, Tutorials and Homework will be published on the University of Hull **Canvas Website**

canvas.hull.ac.uk/courses/73579

Revision Guide

A Revision Guide to prepare for the Exam is available at

silviofanzon.com/2024-NSS-Revision

Digital Notes

Digital version of these notes available at

silviofanzon.com/2024-NSS-Notes

Readings

We will study the set of real numbers \mathbb{R} , and then sequences and series in \mathbb{R} . I will follow mainly the textbook by Bartle and Sherbert [2]. Another good reading is the book by Abbott [1]. I also point out the classic book by Rudin [3], although this is a more difficult read.

You are not expected to purchase any of the above books. These lecture notes will cover 100% of the topics you are expected to known in order to excel in the Homework and Final Exam.

1 Introduction

The first aim of this lecture notes is to rigorously introduce the set of **Real Numbers**, which is denoted by \mathbb{R} . But what do we mean by real numbers? To start our discussion, introduce the set of natural numbers (or non-negative integers)

$$\mathbb{N} = \{0, 1, 2, 3, 4, 5, \dots\}$$

On this set we have a notion of **sum** of two numbers, denoted as usual by

n + m

for $n, m \in \mathbb{N}$. Here the symbol \in denotes that m and n belong to \mathbb{N} . For example 3 + 7 results in 10.

Question 1.1

Can the sum be inverted? That is, given any $n, m \in \mathbb{N}$, can you always find $x \in \mathbb{N}$ such that

$$n + x = m? \tag{1.1}$$

Of course to invert (1.1) we can just perform a **subtraction**, implying that

$$x = m - n$$

But there is a catch. In general x does not need to be in \mathbb{N} . For example, take n = 10 and m = 1. Then x = -9, which does not belong to \mathbb{N} . Therefore the answer to Question 1.1 is **NO**.

To make sure that we can always invert the sum, we need to **extend** the set \mathbb{N} . This is done simply by introducing the set of **integers**

$$\mathbb{Z} := \{-n, n : n \in \mathbb{N}\},\$$

that is, the set

$$\mathbb{Z} := \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}.$$

The sum can be extended to \mathbb{Z} , by defining

$$(-n) + (-m) := -(m+n)$$
 (1.2)

for all $m, n \in \mathbb{N}$. Now every element of \mathbb{Z} possesses an **inverse**, that is, for each $n \in \mathbb{Z}$, there exists $m \in \mathbb{Z}$, such that

$$n+m=0.$$

Can we characterize m explicitly? Yes; Seeing the definition at (1.2), we simply have

m = -n.

On the set \mathbb{Z} we can also define the operation of **multiplication** in the usual way. For $n, m \in \mathbb{Z}$, we denote the multiplication of *n* times *m* by

nm or $n \cdot m$.

For example $7 \cdot 2 = 14$ and $1 \cdot (-1) = -1$.

Question 1.2

Can the multiplication in \mathbb{Z} be inverted? That is, given any $n, m \in \mathbb{Z}$, can you always find $x \in \mathbb{Z}$ such that

$$nx = m? \tag{1.3}$$

To invert (1.3) if $n \neq 0$, we can just perform a **division**, to obtain

$$x = \frac{m}{n}$$

But again there is a catch. Indeed taking n = 2 and m = 1 yields x = 1/2, which does not belong to \mathbb{Z} . The answer to Question 1.2 is therefore **NO**.

Thus, in order to invert the multiplication, we need to **extend** the set of integers \mathbb{Z} . This extension is called the set of **rational numbers**, defined by

$$\mathbb{Q} := \left\{ \frac{m}{n} : m, n \in \mathbb{Z}, n \neq 0 \right\}$$

We then extend the operations of sum and multiplication to Q by defining

$$\frac{m}{n} + \frac{p}{q} := \frac{mq + np}{nq}$$
$$\frac{m}{n} \cdot \frac{p}{q} := \frac{mp}{nq}$$

and

Now the multiplication is invertible in \mathbb{Q} . Specifically, each non-zero element has an inverse: the inverse of m/n is given by n/m.

To summarize, we have extended \mathbb{N} to \mathbb{Z} , and \mathbb{Z} to \mathbb{Q} . By construction we have

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q}$$
.

Moreover sum and product are invertible in Q. Now we are happy right? So and so.

Question 1.3

Can we represent the set Q with a drawing?

It is clear how to draw \mathbb{Z} , as seen below.

-4 -3 -2 -1 0 1 2 3 4

Figure 1.1: Representation of integers \mathbb{Z}

However \mathbb{Q} is much **larger** than the set \mathbb{Z} represented by the ticks in Figure 1.1. What do we mean by **larger**? For example, consider $0 \in \mathbb{Q}$.

Question 1.4

What is the number $x \in \mathbb{Q}$ which is closest to 0?

There is no right answer to the above question, since whichever rational number m/n you consider, you can always squeeze the rational number m/(2n) in between:

$$0<\frac{m}{2n}<\frac{m}{n}\,.$$

For example think about the case of the numbers

$$\frac{1}{n}$$
 for $n \in \mathbb{N}$, $n \neq 0$.

Such numbers get arbitrarily close to 0, as depicted below.



Figure 1.2: Fractions $\frac{1}{n}$ can get arbitrarily close to 0

Maybe if we do the same reasoning with other progressively smaller rational numbers, we manage to fill up the interval [0, 1]. In other words, we might conjecture the following.

Conjecture 1.5

Maybe \mathbbm{Q} can be represented by a continuous line.

Do you think the above conjecture is true? Conjecture 1.5 is **false**, as shown by the Theorem below.

Theorem 1.6

The number $\sqrt{2}$ does not belong to \mathbb{Q} .

Theorem 1.6 is the reason why $\sqrt{2}$ is called an **irrational number**. For reference, a few digits of $\sqrt{2}$ are given by

 $\sqrt{2} = 1.414213562373095048...$

and the situation is as in the picture below.



Figure 1.3: Representing $\sqrt{2}$ on the numbers line.

We can therefore see that Conjecture 1.5 is **false**, and Q is not a line: indeed Q has a **gap** at $\sqrt{2}$. Let us see why Theorem 1.6 is true.

Proof: Proof of Theorem 1.6

We prove that

 $\sqrt{2} \notin \mathbb{Q}$

by contradiction.

Wait, what does this mean? Proving the claim by contradiction means assuming that the claim is **false**. This means we **assume** that

$$\sqrt{2} \in \mathbb{Q} \,. \tag{1.4}$$

From this assumption we then start deducing other statements, hoping to encounter a statement which is **FALSE**. But if (1.4) leads to a false statement, then it must be that (1.4) was **FALSE** to begin with. Thus the contrary of (1.4) must hold, meaning that

 $\sqrt{2} \notin \mathbb{Q}$

as we wanted to show. This would conclude the proof. Now we need to actually show that assuming (1.4) will lead to a contradiction. Since this is our first proof, let us take it slowly, step-by-step.

1. Assuming (1.4) just means that there exists $q \in \mathbb{Q}$ such that

$$q = \sqrt{2} \,. \tag{1.5}$$

2. Since $q \in \mathbb{Q}$, by definition we have

$$q = \frac{m}{n}$$

for some $m, n \in \mathbb{N}$ with $n \neq 0$.

3. Recalling (1.5), we then have

$$\frac{m}{n} = \sqrt{2}$$
.

4. We can square the above equation to get

$$\frac{m^2}{n^2} = 2. (1.6)$$

5. Withouth loss of generality, we can assume that *m* and *n* have no common factors.

Wait. What does Step 5 mean? You will encounter the sentence *withouth loss of generality* many times in mathematics. It is often abbreviated in **WLOG**. WLOG means that we can make some extra assumption which does not affect the validity of the proof in general.

For example in our case we can assume that m and n have no common factor. This is because if m and n had common factors, then it would mean

$$m = a\tilde{m}, \quad n = a\tilde{n}$$

for some factor $a \in \mathbb{N}$ with $a \neq 0$. Then

$$\frac{m}{n} = \frac{a\tilde{m}}{a\tilde{n}} = \frac{\tilde{m}}{\tilde{n}}$$

Therefore by (1.6)

$$\frac{\tilde{m}^2}{\tilde{n}^2} = 2$$

and this time \tilde{m} and \tilde{n} have no common factors by construction. The proof can now proceed in the same way it would have proceeded from Step 4, but in addition we have the hypothesis that \tilde{m} and \tilde{n} have no common factors.

6. Equation (1.6) implies

$$m^2 = 2n^2$$
. (1.7)

Therefore the integer m^2 is an even number.

Why is m^2 even? As you already know, **even** numbers are

0, 2, 4, 6, 8, 10, 12, ...

All these numbers have in common that they can be divided by 2, and so they can be written as

2p

for some $p \in \mathbb{N}$. For example 52 is even, because

$$52 = 2 \cdot 26$$

Instead, odd numbers are

These can be all written as

2p + 1

for some $p \in \mathbb{N}$. For example 53 is odd, because

$$52 = 2 \cdot 26 + 1$$

7. Since m^2 is an even number, it follows that also *m* is an even number. Then there exists $p \in \mathbb{N}$ such that

$$m = 2p. \tag{1.8}$$

Why is *m* even if m^2 is even? Let us see what happens if we take the square of an even number m = 2p

$$m^2 = (2p)^2 = 4p^2 = 2(2p^2) = 2q$$

Thus $m^2 = 2q$ for some $q \in \mathbb{N}$, and so m^2 is an even number. If instead m is odd, then m = 2p + 1 and

$$m^{2} = (2p+1)^{2} = 4p^{2} + 4p + 1 = 2(2p^{2} + 2p) + 1$$

showing that also m^2 is odd.

This proves Step 7: Indeed we know that m^2 is an even number from Step 6. If m was odd, then m^2 would be odd. Hence m must be even as well.

8. If we substitute (1.8) in (1.7) we get

$$m^2 = 2n^2 \implies (2p)^2 = 2n^2 \implies 4p^2 = 2n^2$$

Dividing both terms by 2, we obtain

$$n^2 = 2p^2. (1.9)$$

- 9. We now make a series of observations:
 - Equation (1.9) says that n^2 is even.
 - The same argument in Step 7 guarantees that also *n* is even.
 - We have already seen that *m* is even.
 - Therefore *n* and *m* are both even.
 - Hence *n* and *m* have 2 as common factor.
 - But Step 5 says that *n* and *m* have no common factors.
 - CONTRADICTION
- Our reasoning has run into a contradiction, stemming from assumption (1.4). Therefore (1.4) is FALSE, and so

$$\sqrt{2} \notin \mathbb{Q}$$

ending the proof.

Seeing that $\sqrt{2} \notin \mathbb{Q}$, we might be tempted to just fill in the gap by adding $\sqrt{2}$ to \mathbb{Q} . However, with analogous proof to Theorem 1.6, we can prove that

 $\sqrt{p} \notin \mathbb{Q}$

for each prime number p. As there are infinite prime numbers, this means that \mathbb{Q} has infinite gaps. Then we might attempt to fill in these gaps via the extension

 $\tilde{\mathbb{Q}} := \mathbb{Q} \cup \{\sqrt{p} : p \text{ prime}\}.$

However even this is not enough, as we would still have numbers which are not contained in \tilde{Q} , for example

$$\sqrt{2} + \sqrt{3}, \pi, \pi + \sqrt{2} \notin \tilde{\mathbb{Q}}$$
.

Conclusion: It is now intuitive to think that there is no straightforward way to fill the gaps of \mathbb{Q} by adding numbers by hand.

Remark 1.7	
Proving that $\sqrt{2} + \sqrt{3} \notin \mathbb{Q}$	
is relatively easy, and will be left as an exercise . Instead, proving that	
$\pi \notin \mathbb{Q}$	
is way more complicated. There are several proofs of the fact, all requiring mathematics which is more advanced than the one presented in this course. For some proofs, see this Wikipedia page.	

The reality of things is that to **complete** Q and make it into a **continuous line** we have to add a lot of points. Indeed, we need to add way more points than the ones already contained in Q.

Definition 1.8

Such extension of $\mathbb Q$ will be called $\mathbb R,$ the set of real numbers.

The inclusions will therefore be

$$\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}.$$

The set \mathbb{R} is not at all trivial to construct. In fact, explicit constructions are so involved and technical, that they are best left for future reading. It is customary instead to proceed as follows:

- We will assume that $\mathbb R \texttt{ exists}$ and satisfies a set of axioms.
- One of the axioms states that $\mathbb R$ fills **all the gaps** that $\mathbb Q$ has. Therefore $\mathbb R$ can be thought as a **continuous** line.
- We will study the $\mathbf{properties}$ of $\mathbb R$ which descend from such $\mathbf{axioms}.$

For example one of the properties of $\mathbb R$ will be the following:

Theorem 1.9: We will prove this in the future

 \mathbb{R} contains all the square roots. This means that for every $x \in \mathbb{R}$ with $x \ge 0$, we have

 $\sqrt{x} \in \mathbb{R}$.

A concrete **model** for the real numbers \mathbb{R} can be constructed using **Dedekind cuts**. The interested reader can refer to the Appendix in Chapter 1 of [3], or to the beautifully written Chapter 8.6 in [1]. Such model of \mathbb{R} can be used to prove the following Theorem:

Theorem 1.10: Existence of the Real Numbers

There exists a set \mathbb{R} , called the set of Real Numbers, which has the following properties:

• \mathbb{R} extends \mathbb{Q} , that is,

 $\mathbb{Q} \subset \mathbb{R}$.

- \mathbbm{R} satisfies a certain set of axioms.
- ${\mathbb R}$ fills all the gaps that ${\mathbb Q}$ has. In particular ${\mathbb R}$ can be represented by a continuous line.

2 Preliminaries

Before introducing \mathbb{R} we want to make sure that we cover all the basics needed for the task.

2.1 Sets

A set is a **collection** of objects. These objects are called **elements** of the set. For example in the previous section we mentioned the following sets:

- $\mathbb N$ the set of natural numbers
- \mathbb{Z} the set of integers
- Q the set of rational numbers
- \mathbb{R} the set of real numbers

Definition 2.1

Let A be a set.

- 1. We write $x \in A$ if the element x belongs to the set A.
- 2. We write $x \notin A$ if the element *x* does not belong to the set *A*.

Remark 2.2

A set can contain all sorts of elements. For example the students in a classroom can be modelled by a set *S*. The elements of the set are the students. For example

 $S = \{Alice, Olivia, Jake, Sahab\}$

In this case we have

```
Alice \in S
```

but instead

Silvio $\notin S$.

2.2 Logic

In this section we introduce some basic logic symbols. Suppose that you are given two statements, say α and β . The formula $\alpha \implies \beta$

means that
$$\alpha$$
 implies β . In other words, if α is true then also β is true.
The formula
 $\alpha \leftarrow \beta$

means that α is implied by β : if β is true then also α is true. When we write

$$\alpha \iff \beta \tag{2.1}$$

we mean that α and β are **equivalent**. We also say that α is true **if and only if** β is true. Note that (2.1) is equivalent to

$$\alpha \implies \beta \text{ and } \beta \implies \alpha$$
.

Example 2.3We have that $x > 0 \implies x > -100$,andcontradiction $\leftarrow \sqrt{2} \in \mathbb{Q}$.Concerning \leftrightarrow we have $x^2 < 2 \iff -\sqrt{2} < x < \sqrt{2}$.

We now introduce logic **quantifiers**. These are

- ∀ which reads **for all**
- 3 which reads **exists**
- ∃! which reads **exists unique**
- ∄ which reads **does not exists**

These work in the following way. Suppose that you are given a statement $\alpha(x)$ which depends on the point $x \in \mathbb{R}$. Then we say

• $\alpha(x)$ is satisfied for all $x \in A$ with A some collection of numbers. This translates to the symbols

$$\alpha(x)$$
 is true $\forall x \in A$,

• There exists some x in \mathbb{R} such that $\alpha(x)$ is satisfied: in symbols

 $\exists x \in \mathbb{R}$ such that $\alpha(x)$ is true,

- There exists a unique x_0 in \mathbb{R} such that $\alpha(x)$ is satisfied: in symbols

 $\exists x_0 \in \mathbb{R}$ such that $\alpha(x_0)$ is true,

• $\alpha(x)$ is never satisfied:

 $\nexists x \in \mathbb{R}$ such that $\alpha(x)$ is true.

Example 2.4

Let us make concrete examples:

• The expression x^2 is always non-negative. Thus we can say

 $x^2 \ge 0$ for all $x \in \mathbb{R}$.

• The equation $x^2 = 1$ has two solutions x = 1 and x = -1. Therefore we can say

 $\exists x \in \mathbb{R}$ such that $x^2 = 1$.

• The equation $x^3 = 1$ has a unique solution x = 1. Thus

```
\exists ! x \in \mathbb{R} such that x^3 = 1.
```

• We know that the equation $x^2 = 2$ has no solutions in Q. Then

 $\nexists x \in \mathbb{Q}$ such that $x^2 = 2$.

2.3 Operations on sets

Definition 2.5: Inclusion and equality

Given two sets A and B, we say that A is **contained** in B, in symbols

 $A \subseteq B$,

if all the elements of A are also contained in B. Two sets A and B are **equal**, in symbols

A = B,

if they contain the same elements.

Remark 2.6

The inclusion $A \subseteq B$ is equivalent to the implication:

 $x \in A \implies x \in B$

for all $x \in A$. The symbol \implies reads **implies**, and denotes the fact that the first condition implies the second.

Definition 2.7: Union and Intersection

For two sets *A* and *B* we define their **union** as the set

$$A \cup B := \{x : x \in A \text{ or } x \in B\}.$$

The **intersection** of *A* and *B* is defined by

$$A \cap B := \{x : x \in A \text{ and } x \in B\}.$$

We denote the **empty set** by the symbol Ø. Two sets are **disjoint** if

 $A \cap B = \emptyset$.

Example 2.8

Given two sets *A* and *B* we always have

$$(A \cap B) \subseteq A, \ (A \cap B) \subseteq B, \tag{2.2}$$

 $A \subseteq (A \cup B), \ B \subseteq (A \cup B).$ (2.3)

Example 2.9

Define the subset of rational numbers

$$S := \left\{ x \in \mathbb{Q} : 0 < x < \frac{5}{2} \right\}.$$

Then we have

 $\mathbb{N} \cap S = \{1, 2\}.$

We can also define the sets of **even** and **odd** numbers by

$$E := \{2n : n \in \mathbb{N}\},$$
(2.4)

$$O := \{2n+1 : n \in \mathbb{N}\}.$$
 (2.5)

Then we have

$$\mathbb{N} \cap E = E, \ \mathbb{N} \cap O = O, \tag{2.6}$$

$$O \cup E = \mathbb{N}, \ O \cap D = \emptyset.$$
(2.7)

Example 2.10

1. The sets

 $A = \{1, 2, 3\}, \quad B = \{3, 1, 2\}$

are equal, that is A = B. This is because they contain exactly the same elements: **order** does not matter when talking about sets.

2. Consider the sets

 $A = \{1, 2\}, \quad B = \{1, 2, 5\}.$

Then *A* is contained in *B*, but *A* is not equal to *B*. Therefore we write $A \subseteq B$ or $A \neq B$.

The next proposition is very useful when we need to **prove** that two sets are equal: rather than showing directly that A = B, we can prove that $A \subseteq B$ and $B \subseteq A$.

Proposition 2.11

Let A and B be sets. Then

 $A = B \iff A \subseteq B \text{ and } B \subseteq A$.

Proof

The proof is almost trivial. However it is a good exercise in basic logic, so let us do it.

1. First implication \implies : Suppose that A = B. Let us show that $A \subseteq B$. Since A = B, this means that all the elements of A are also contained in B. Therefore if we take $x \in A$ we have

 $x \in A \implies x \in B$.

This shows $A \subseteq B$. The proof of $B \subseteq A$ is similar.

2. Second implication \leftarrow :

Suppose that $A \subseteq B$ and $B \subseteq A$. We need to show A = B, that is, A and B have the same elements. To this end let $x \in A$. Since $A \subseteq B$ then we have $x \in B$. Thus B contains all the elements of A. Since we are also assuming $B \subseteq A$, this means that A contains all the elements of B. Hence A and B contain the same elements, and A = B.

2.3.1 Infinite union and intersection

Definition 2.12: Infinite union and intersection

Let Ω be a set, and $A_n \subseteq \Omega$ a family of subsets, where $n \in \mathbb{N}$.

1. The **infinte union** of the A_n is the set

$$\bigcup_{n \in \mathbb{N}} A_n := \{ x \in \Omega : x \in A_n \text{ for at least one } n \in \mathbb{N} \}.$$

2. The **infinte intersection** of the A_n is the set

$$\bigcap_{n \in \mathbb{N}} A_n := \{ x \in \Omega : x \in A_n \text{ for all } n \in \mathbb{N} \}.$$

Example 2.13

Question. Define $\Omega := \mathbb{N}$ and a family A_n by

$$A_n = \{n, n+1, n+2, n+3, ...\}, \quad n \in \mathbb{N}.$$

1. Prove that

$$\bigcup_{n\in\mathbb{N}}A_n=\mathbb{N}\,.\tag{2.8}$$

2. Prove that

$$\bigcap_{n \in \mathbb{N}} A_n = \emptyset.$$
(2.9)

Solution.

1. Assume that $m \in \bigcup_n A_n$. Then $m \in A_n$ for at least one $n \in \mathbb{N}$. Since $A_n \subseteq \mathbb{N}$, we conclude that $m \in \mathbb{N}$. This shows

$$\bigcup_{n\in\mathbb{N}}A_n\subseteq\mathbb{N}\,.$$

Conversely, suppose that $m \in \mathbb{N}$. By definition $m \in A_m$. Hence there exists at least one index n, n = m in this case, such that $m \in A_n$. Then by definition $m \in \bigcup_{n \in \mathbb{N}} A_n$, showing that

$$\mathbb{N} \subseteq \bigcup_{n \in \mathbb{N}} A_n \, .$$

This proves (2.8).

2. Suppose that (2.9) is false, i.e.,

$$\bigcap_{n\in\mathbb{N}}A_n\neq\emptyset.$$

This means there exists some $m \in \mathbb{N}$ such that $m \in \bigcap_{n \in \mathbb{N}} A_n$. Hence, by definition, $m \in A_n$ for all $n \in \mathbb{N}$. However $m \notin A_{m+1}$, yielding a contradiction. Thus (2.9) holds.

2.3.2 Complement

Definition 2.14: Complement

Let $A, B \subseteq \Omega$. The **complement** of *A* with respect to *B* is the set of elements of *B* which do not belong to *A*, that is

$$B \setminus A := \{x \in \Omega : x \in B \text{ and } x \notin A\}.$$

In particular, the complement of *A* with respect to Ω is denoted by

 $A^c := \Omega \setminus A := \{ x \in \Omega : x \notin A \}.$

Remark 2.15

Suppose that $A \subseteq \Omega$. Then *A* and A^c form a **partition** of Ω , in the sense that

 $A \cup A^c = \Omega$ and $A \cap A^c = \emptyset$.

Example 2.16

Question. Suppose $A, B \subseteq \Omega$. Prove that

$$A \subseteq B \iff B^c \subseteq A^c$$
.

Solution. Let us prove the above claim:

- First implication ⇒ : Suppose that A ⊆ B. We need to show that B^c ⊆ A^c. Hence, assume x ∈ B^c. By definition this means that x ∉ B. Now notice that we cannot have that x ∈ A. Indeed, assume x ∈ A. By assumption we have A ⊆ B, hence x ∈ B. But we had assumed x ∈ B, contradiction. Therefore it must be that x ∉ A. Thus B^c ⊆ A^c.
- Second implication ⇐ : Note that, for any set,

$$(A^c)^c = A.$$

Hence, by the first implication,

 $B^c \subseteq A^c \implies (A^c)^c \subseteq (B^c)^c \implies A \subseteq B.$

We conclude by stating the De Morgan's Laws. The proof will be left as an exercise.

Proposition 2.17: De Morgan's Laws

Suppose $A, B \subseteq \Omega$. Then

$$(A \cap B)^c = A^c \cup B^c$$
, $(A \cup B)^c = A^c \cap B^c$.

2.3.3 Power set

Definition 2.18: Power set

Let Ω be a set. The **power set** of Ω is

$$\mathscr{P}(\Omega) := \{A : A \subseteq \Omega\}.$$

Hence, the power set of Ω is the set of all subsets of Ω .

Remark 2.19

It holds that:

1. $\mathscr{P}(\Omega)$ is always non-empty, since we have that

$$\emptyset \in \mathscr{P}(\Omega), \quad \Omega \in \mathscr{P}(\Omega).$$

2. Given $A, B \in \mathcal{P}(\Omega)$, then the sets

 $A \cup B$, $A \cap B$, A^c , $B \smallsetminus A$

are all elements of $\mathscr{P}(\Omega)$.

3. Suppose Ω is **discrete** and **finite**, that is,

$$\Omega = \{x_1, \dots, x_m\}$$

for some $m \in \mathbb{N}$. Then $\mathscr{P}(\Omega)$ contains 2^m elements.

Indeed, suppose that we want to define a subset $A \subset \Omega$. Then for each element $x_i \in \Omega$ there are two choices: either we include $x_i \in A$, or we do not include $x_i \in A$. Therefore there are

$$\underbrace{2 \cdot 2 \cdot \ldots \cdot 2}_{m \text{ times}} = 2^m$$

ways of constructing A. It follows that Ω possesses exactly 2^m subsets, so that $\mathscr{P}(\Omega)$ contains 2^m elements.

Example 2.20

Question. Compute the power set of

 $\Omega = \{x, y, z\}.$

Solution. $\mathscr{P}(\Omega)$ has $2^3 = 8$, and

$$\mathcal{P}(\Omega) = \{\emptyset, \{x\}, \{y\}, \{z\}, \{x, y\} \\ \{x, z\}, \{y, z\}, \{x, y, z\}\}$$

2.3.4 Product of sets

Definition 2.21: Product of sets

Let A, B be sets. The **product** of A and B is the set of pairs

 $A \times B := \{(a, b) : a \in A, b \in B\}.$

By definition, two elements in $A \times B$ are the same, in symbols

 $(a,b) = (\tilde{a},\tilde{b})$

if and only if they are equal component-by-componenent, that is

 $a = \tilde{a}$, $b = \tilde{b}$.

2.4 Equivalence relation

Definition 2.22: Binary relation

Suppose A is a set. A **binary relation** R on A is a subset

 $R\subseteq A\times A\,.$

Definition 2.23: Equivalence relation

A binary relation R is called an **equivalence relation** if it satisfies the following properties:

1. **Reflexive**: For each $x \in A$ one has

 $(x,x)\in R$,

This is saying that all the elements in A must be related to themselves

2. Symmetric: We have

 $(x, y) \in R \implies (y, x) \in R$

If x is related to y, then y is related to x

3. Transitive: We have

$$(x, y) \in R, (y, z) \in R \implies (x, z) \in R$$

If x is related to y, and y is related to z, then x must be related to z

If $(x, y) \in R$ we write

 $x \sim y$

and we say that *x* and *y* are **equivalent**.

Definition 2.24: Equivalence classes

Suppose *R* is an **equivalence relation** on *A*. The **equivalence class** of an element $x \in A$ is the set

$$[x] := \{ y \in A : y \sim x \}.$$

The set of equivalence classes of elements of A with respect to the equivalence relation R is denoted by

$$A/R := A/\sim := \{[x] : x \in A\}.$$

In order for the definition of [x] to be *well-posed* we need to check that:

- 1. [x] is non-empty.
- 2. [*x*] does not depend on the representative *x*: we need to check that

$$x \sim y \quad \Longleftrightarrow \quad [x] = [y]$$

This is shown in the following proposition.

Proposition 2.25: Well-posedness of Definition 2.24

Let ~ be an equivalence relation on A. Then

- 1. For each $x \in A$ we have $[x] \neq \emptyset$.
- 2. For all $x, y \in A$ it holds

 $x \sim y \quad \Longleftrightarrow \quad [x] = [y].$

Proof

Proof of Point 1: By the *reflexive property* we have $x \sim x$. By definition of equivalence class we conclude that $x \in [x]$. This shows

 $[x] \neq \emptyset$

Proof of Point 2: We need to prove a double implication. It is convenient to divide the proof into two parts.

Part 1: $x \sim y \implies [x] = [y].$

Assume $x \sim y$. By the definition of an equivalence class

 $[x] = \{z \in A \mid z \sim x\}$

is the set of all elements in A that are related to x. Similarly,

$$[y] = \{z \in A \mid z \sim y\}.$$

We need to show that

$$[x] = [y],$$

meaning that every element in [x] is also in [y], and vice versa.

- 1. First, take an arbitrary element $z \in [x]$.
 - By definition, $z \sim x$.
 - Since ~ is an equivalence relation, it satisfies the *transitive property*.
 - Therefore, from $z \sim x$ and $x \sim y$, we can conclude that $z \sim y$
 - Hence, $z \in [y]$.
 - This shows that $[x] \subseteq [y]$.

2. Now, take an arbitrary element $z \in [y]$.

- By definition, $z \sim y$.
- Since ~ is an equivalence relation, it satisfies the symmetric property.
- Therefore, from $x \sim y$, we also have $y \sim x$.
- By the *transitive property*, from $z \sim y$ and $y \sim x$, we can conclude that $z \sim x$.
- Hence, $z \in [x]$.
- This shows that $[y] \subseteq [x]$.

Since $[x] \subseteq [y]$ and $[y] \subseteq [x]$, it follows that [x] = [y], as required. Thus, we have shown that $x \sim y \implies [x] = [y]$. *Part 2:* $[x] = [y] \implies x \sim y$.

Assume [x] = [y]. This means that the equivalence classes of *x* and *y* are the same.

- By point (i) in the Proposition, we have $x \in [x]$ and $y \in [y]$.
- Since [x] = [y], we have $x \in [y]$.
- By the definition of [y], this means $x \sim y$.

Thus, we have shown that $[x] = [y] \implies x \sim y$. *Conclusion:* Since we have proven both directions:

• $x \sim y \implies [x] = [y]$ • $[x] = [y] \implies x \sim y$

we conclude that

$$x \sim y \iff [x] = [y].$$

This completes the proof.

The prototypical (and trivial) example of equivalence relation is the **equality** over Q.

Example 2.26: Equality is an equivalence relation

Question. The equality defines a **binary relation** on $\mathbb{Q} \times \mathbb{Q}$, via

$$R := \{(x, y) \in \mathbb{Q} \times \mathbb{Q} : x = y\}.$$

- 1. Prove that *R* is an **equivalence relation**.
- 2. Prove that $[x] = \{x\}$ and compute \mathbb{Q}/R .

Solution.

- 1. We need to check that *R* satisfies the 3 properties of an equivalence relation:
 - Reflexive: It holds, since x = x for all $x \in \mathbb{Q}$,
 - Symmetric: Again x = y if and only if y = x,
 - Transitive: If x = y and y = z then x = z.

Therefore, R is an equivalence relation.

2. The class of equivalence of $x \in \mathbb{Q}$ is given by

 $[x] = \{x\},\$

that is, this relation is quite trivial, given that each element of $\mathbb Q$ can only be related to itself. The quotient space is then

$$\mathbb{Q}/R = \{ [x] : x \in \mathbb{Q} \} = \{ \{x\} : x \in \mathbb{Q} \}.$$

We now give an example of a non-trivial equivalence relation over Q.

Example 2.27

Question. Let R be the binary relation on the set \mathbb{Q} of rational numbers defined by

 $x \sim y \iff x - y \in \mathbb{Z}$.

- 1. Prove that R is an equivalence relation on \mathbb{Q} .
- 2. Compute [x] for each $x \in \mathbb{Q}$.
- 3. Compute \mathbb{Q}/R .

Solution.

- 1. We have:
 - Reflexive: Let $x \in \mathbb{Q}$. Then x x = 0 and $0 \in \mathbb{Z}$. Thus $x \sim x$.
 - Symmetric: If $x \sim y$ then $x y \in \mathbb{Z}$. But then also

$$-(x-y) = y - x \in \mathbb{Z}$$

and so $y \sim x$.

• Transitive: Suppose $x \sim y$ and $y \sim z$. Then

 $x - y \in \mathbb{Z}$ and $y - z \in \mathbb{Z}$.

Thus, we have

$$x - z = (x - y) + (y - z) \in \mathbb{Z}$$

showing that $x \sim z$.

Thus, we have shown that R is an equivalence relation on \mathbb{Q} .

2. Note that

 $x \sim y \quad \iff \quad \exists n \in \mathbb{Z} \text{ s.t. } y = x + n.$

Therefore the equivalence classes with respect to ~ are

 $[x] = \{x + n : n \in \mathbb{Z}\}.$

Each equivalence class has exactly one element in $[0, 1) \cap \mathbb{Q}$, meaning that:

$$\forall x \in \mathbb{Q}, \exists ! q \in \mathbb{Q} \text{ s.t. } 0 \le q < 1 \text{ and } q \in [x].$$
(2.10)

Condition (2.10) is illustrated in Figure 2.1. Indeed: take $x \in \mathbb{Q}$ arbitrary. Then $x \in [n, n+1)$ for some $n \in \mathbb{Z}$. Setting q := x - n we obtain that

$$x=q+n, \qquad q\in [0,1),$$

proving (2.10). In particular (2.10) implies that for each $x \in \mathbb{Q}$ there exists $q \in [0, 1) \cap \mathbb{Q}$ such that

$$[x] = [q]$$

3. From Point 2 we conclude that

 $\mathbb{Q}/R = \{ [x] : x \in \mathbb{Q} \} = \{ q \in \mathbb{Q} : 0 \le q < 1 \}.$



Figure 2.1: For each $x \in \mathbb{Q}$ there exist unique $q \in [0, 1) \cap \mathbb{Q}$ such that x = q + n. In particular [x] = [q].

2.5 Order relation

Order relations are defined similarly to **equivalence relations**. However notice that *symmetry* is replaced by *antisymmetry*.

Definition 2.28: Partial order

A binary relation R on A is called a **partial order** if it satisfies the following properties:

1. **Reflexive**: For each $x \in A$ one has

 $(x, x) \in R$,

2. Antisymmetric: We have

 $(x, y) \in R$ and $(y, x) \in R \implies x = y$

3. Transitive: We have

 $(x, y) \in R, (y, z) \in R \implies (x, z) \in R$

Remark 2.29

Notice that conditions 1 and 3 are the same for equivalence and order relations. Instead condition 2 changes: it is symmetry for equivalence relation, and antisymmetry for order relations.

We say that a partial order is total if all the elements in the set are related.

Definition 2.30: Total order

A binary relation R on A is called a **total order relation** if it satisfies the following properties:

- 1. **Partial order**: *R* is a partial order on *A*.
- 2. **Total**: For each $x, y \in A$ we have

 $(x, y) \in R$ or $(y, x) \in R$.

The operation of **set inclusion** is a partial order on $P(\Omega)$ but **not** a total order.

Example 2.31: Set inclusion is a partial order but not total order

Question. Let Ω be a non-empty set and consider its **power set**

$$\mathscr{P}(\Omega) = \{A : A \subseteq \Omega\}.$$

The inclusion defines **binary relation** on $\mathscr{P}(\Omega) \times \mathscr{P}(\Omega)$, via

$$R := \{ (A, B) \in \mathscr{P}(\Omega) \times \mathscr{P}(\Omega) : A \subseteq B \}.$$

- 1. Prove that *R* is an **order relation**.
- 2. Prove that R is **not a total order**.

Solution.

- 1. Check that *R* is a partial order relation on $\mathscr{P}(\Omega)$:
 - Reflexive: It holds, since $A \subseteq A$ for all $A \in \mathcal{P}(\Omega)$.
 - Antisymmetric: If $A \subseteq B$ and $B \subseteq A$, then A = B.
 - Transitive: If $A \subseteq B$ and $B \subseteq C$, then, by definition of inclusion, $A \subseteq C$.
- 2. In general, *R* is **not** a total order. For example consider

$$\Omega = \{x, y\}.$$

Thus

$$\mathscr{P}(\Omega) = \{\emptyset, \{x\}, \{y\}, \{x, y\}\}.$$

If we pick $A = \{x\}$ and $B = \{y\}$ then $A \cap B = \emptyset$, meaning that

 $A \not\subseteq B$, $B \not\subseteq A$.

This shows R is not a total order.

The **inequality** on Q is an example of total order.

Example 2.32: Inequality is a total order

Question. Consider the binary relation induced by the inequality on $Q \times Q$:

$$R := \{(x, y) \in \mathbb{Q} \times \mathbb{Q} : x \le y\}.$$

Prove that *R* is a **total order relation**.

Solution. We need to check that:

- 1. Reflexive: It holds, since $x \le x$ for all $x \in \mathbb{Q}$,
- 2. Antisymmetric: If $x \le y$ and $y \le x$ then x = y.
- 3. Transitive: If $x \le y$ and $y \le z$ then $x \le z$.

Finally, we halso have that *R* is a **total order** on \mathbb{Q} , since for all $x, y \in \mathbb{Q}$ we have

 $x \le y$ or $y \le x$.

Notation 2.33

If *A* is a set and *R* is a total order on *A*, we write

$$(x, y) \in R \iff x \leq y$$
.

Therefore the symbol \leq will always denote a total order relation.

2.6 Intervals

In this section we assume to have available the set \mathbb{R} of **real numbers**, which we recall is an extension of \mathbb{Q} . We now introduce the concept of **interval**.

Definition 2.34

Let $a, b \in \mathbb{R}$ with a < b. We define the **open interval** (a, b) as the set

 $(a,b) := \{x \in \mathbb{R} : a < x < b\}.$

We define the **close interval** [a, b] as the set

$$[a,b] := \{x \in \mathbb{R} : a \le x \le b\}.$$

In general we also define the intervals

- $[a,b) := \{ x \in \mathbb{R} : a \le x < b \},$ (2.11)
- $(a,b] := \{ x \in \mathbb{R} : a \le x \le b \},$ (2.12)
- $(a, \infty) := \{ x \in \mathbb{R} : x > a \},$ (2.13)
- $[a,\infty) := \{ x \in \mathbb{R} : x \ge a \}, \tag{2.14}$
- $(-\infty, b) := \{ x \in \mathbb{R} : x < b \},$ (2.15)
- $(-\infty, b] := \{ x \in \mathbb{R} : x \le b \}.$ (2.16)

Some of the above intervals are depicted in Figure 2.2, Figure 2.3, Figure 2.4, Figure 2.5 below.



Figure 2.2: Interval (a, b)

 $a \xrightarrow{\bullet} \mathbb{R}$

Figure 2.3: Interval [a, b]

2.7 Functions

Definition 2.35: Functions

Let *A* and *B* be sets. A **function** from *A* to *B* is a rule which associates **at each** element $x \in A$ a **single** element $y \in B$. Notations:

• We write

$$f : A \to B$$

to indicate such rule,

• For $x \in A$, we denote by

$$y := f(x) \in B$$

the element associated with x by f.

- We will often denote the map f also by

 $x \mapsto f(x)$.

In addition:

- The set A is called the **domain** of f,
- The **range** or **image** of *f* is the set

 $f(A) := \{y \in B : y = f(x) \text{ for some } x \in A\} \subseteq B.$

Warning

We want to stress the importance of the first two sentences in Definition 2.35. Assume that $f: A \rightarrow B$ is a function. Then:

- To each element $x \in A$ we can **only** associate **one** element $f(x) \in B$,
- Every element $x \in A$ has to be associated to an element $f(x) \in B$.

ā

Figure 2.4: Interval (a, ∞)

 $b \longrightarrow \mathbb{R}$

Figure 2.5: Interval $(-\infty, b]$

Example 2.36

Assume given the two sets

$$A = \{a_1, a_2\}, \quad B = \{b_1, b_2, b_3\}.$$

Let us see a few examples:

1. Define $f : A \rightarrow B$ by setting

$$f(a_1) = b_1$$
, $f(a_2) = b_1$.

In this way f is a function, with domain A and range

$$f(A) = \{b_1\} \subseteq B.$$

2. Define $g : A \rightarrow B$ by setting

$$g(a_1) = b_2$$
, $g(a_1) = b_3$, $g(a_2) = b_3$

Then g is **NOT** a function, since the element a_1 has two elements associated.

3. Define $h : A \rightarrow B$ by setting

$$h(a_1) = b_1$$
.

Then g is **NOT** a function, since the element a_2 has no element associated.



Figure 2.6: Schematic picture of the function f



Figure 2.7: Schematic picture of the function *g*



Figure 2.8: Schematic picture of the function h

Example 2.37

Let us make two examples of functions on $\mathbb{R}:$

1. Define $f : \mathbb{R} \to \mathbb{R}$ by

$$f(x) = x^2.$$

Note that the domain of f is given by \mathbb{R} , while the range is

$$f(\mathbb{R}) = [0, \infty).$$

2. Define $g : \mathbb{R} \to \mathbb{R}$ as the logarithm:

$$g(x) = \log(x).$$

This time the domain is $(0, \infty)$, while the range is $g(\mathbb{R}) = \mathbb{R}$.



Figure 2.10: Plot of function $g(x) = \log(x)$

2.8 Absolute value or Modulus

In this section we assume to have available the set $\mathbb R$ of real numbers, which we recall is an extension of $\mathbb Q.$

Definition 2.38: Absolute value

Let $x \in \mathbb{R}$. The **absolute value** of x is

$$x| = \begin{cases} x & \text{if } x \ge 0\\ -x & \text{if } x < 0 \end{cases}$$

Example 2.39

By definition one has |x| = x if $x \ge 0$. For example

$$|\pi| = \pi$$
, $|\sqrt{2}| = \sqrt{2}$, $|0| = 0$.

Instead |x| = -x if x < 0. For example

$$|-\pi| = \pi$$
, $|-\sqrt{2}| = \sqrt{2}$, $|-10| = 10$.

Let us highlight some basic properties of the absolute value.

Proposition 2.40: Properties of absolute value

For all $x \in \mathbb{R}$ they hold:

1. $|x| \ge 0$. 2. |x| = 0 if and only if x = 0. 3. |x| = |-x|.

We can use the definition of absolute value to define the **absolute value function**. This is the function

 $f : \mathbb{R} \to \mathbb{R}, \quad f(x) := |x|.$

You might be familiar with the graph associated to f, as seen below.



Figure 2.11: Plot of the absolute value function f(x) = |x|

It is also useful to understand the absolute value in a geometric way.

```
Remark 2.41: Geometric interpretation of |x|
```

A number $x \in \mathbb{R}$ can be represented with a point on the real line \mathbb{R} . The non-negative number |x| represents the **distance** of x from the origin 0. Notice that this works for both positive and negative numbers x_1 and x_2 respectively, as shown in Figure 2.12 below.



Figure 2.12: Geometric interpretation of |x|

Remark 2.42: Geometric interpretation of |x - y|

If $x, y \in \mathbb{R}$ then the number |x - y| represents the distance between x and y on the real line, as shown in Figure 2.13 below. Note that by Point 3 in Proposition 2.40 we have

$$|x-y| = |y-x|.$$

$$y = |y - x|$$

Figure 2.13: Geometric interpretation of |x - y|
In the next Lemma we show a fundamental equivalence regarding the absolute value.

Lemma 2.43	
Let $x, y \in \mathbb{R}$. Then	
	$ x < y \iff -y < x < y$.

The geometric meaning of the above statement is clear: the distance of *x* from the origin is less than *y*, in formulae $|x| \le y$,

if and only if *x* belongs to the interval [-y, y], in formulae

 $-y \le x \le y$.

A sketch of this explanation is seen in Figure 2.14 below.



Figure 2.14: Geometric meaning of Lemma 2.43

Proof: Proof of Lemma 2.43

Step 1: First implication. Suppose first that

$$|x| \le y \,. \tag{2.17}$$

Recalling that the absolute value is non-negative, from (2.17) we deduce that $0 \le |x| \le y$. In particular it holds

$$y \ge 0. \tag{2.18}$$

We make separate arguments for the cases $x \ge 0$ and x < 0:

• Case 1: $x \ge 0$. From (2.17), (2.18) and from $x \ge 0$ we have

$$-y \le 0 \le x = |x| \le y$$

which shows

$$-y \le x \le y$$

• Case 2: x < 0. From (2.17), (2.18) and from x < 0 we have

 $-y \le 0 < -x = |x| \le y$

which shows

$$-y \le -x \le y \,.$$

Multiplying the above inequalities by -1 yields

 $-y \le x \le y$.

Step 2: Second implication. Suppose now that

 $-y \le x \le y \,. \tag{2.19}$

We make separate arguments for the cases $x \ge 0$ and x < 0:

• Case 1: $x \ge 0$. Since $x \ge 0$, from (2.19) we get

$$|x| = x \le y$$

showing that

 $|x| \leq y$.

• Case 2: x < 0. Since x < 0, from (2.19) we have

 $-y \le x = -|x|.$

Multiplying the above inequality by -1 yields

 $|x| \leq y$.

With the same arguments, just replacing \leq with <, one can also show the following.

Corollary 2.44

Let $x, y \in \mathbb{R}$. Then

```
|x| < y \iff -y < x < y.
```

2.9 Triangle inequality

The triangle inequality relates the absolute value to the sum operation. It is a very important inequality, which we will use a lot in the future.

Theorem 2.45: Triangle inequality For every $x, y \in \mathbb{R}$ we have $||x| - |y|| \le |x + y| \le |x| + |y|$. (2.20)

Before proceeding with the proof, let us discuss the geometric meaning of the triangle inequality.

Remark 2.46: Geometric meaning of triangle inequality

The notion of absolute value can be extended also to vectors in the plane. Suppose that x and y are two vectors in the plane, as in Figure 2.15 below. Then |x| and |y| can be interpreted as the **lengths** of these vectors.

Using the rule of sum of vectors, we can draw x + y, as shown in Figure 2.16 below. From the picture it is evident that

$$|x+y| \le |x|+|y|, \tag{2.21}$$

that is, the length of each side of a triangle does not exceed the sum of the lengths of the two remaining sides. Note that (2.21) is exactly the second inequality in (2.20). This is why (2.20) is called triangle inequality.



Figure 2.15: Vectors x and y



Figure 2.16: Summing the vectors x and y. The triangle inequality relates the length of x + y to the length of x and y

Proof: Proof of Theorem 2.45

Assume that $x, y \in \mathbb{R}$. We prove the two inequalities in (2.20) individually. Step 1. Proof of the second inequality in (2.20). Trivially we have

 $|x| \leq |x|$.

Therefore we can apply Lemma 2.43 and infer

$$-|x| \le x \le |x|. \tag{2.22}$$

Similarly we have that $|y| \le |y|$, and so Lemma 2.43 implies

$$-|y| \le y \le |y|. \tag{2.23}$$

Summing (2.22) and (2.23) we get

 $-(|x| + |y|) \le x + y \le |x| + |y|.$

We can now again apply Lemma 2.43 to get

$$|x+y| \le |x|+|y|, \qquad (2.24)$$

which is the second inequality in (2.20). *Step 2. Proof of the first inequality in (2.20).* Note that the trivial identity

$$x = x + y - y$$

always holds. We then have

$$|x| = |x + y - y| \tag{2.25}$$

$$= |(x + y) + (-y)|$$
(2.26)

$$= |a+b| \tag{2.27}$$

with a = x + y and b = -y. We can now apply (2.24) to *a* and *b* to obtain

$$|\mathbf{x}| = |\mathbf{a} + \mathbf{b}| \tag{2.28}$$

$$\leq |a| + |b| \tag{2.29}$$

$$= |x + y| + |-y|$$
(2.30)

$$= |x + y| + |y|$$
(2.31)

Therefore

$$|x| - |y| \le |x + y|. \tag{2.32}$$

We can now swap x and y in (2.32) to get

 $|y| - |x| \le |x + y|.$

By rearranging the above inequality we obtain

$$-|x+y| \le |x| - |y|.$$
(2.33)

Putting together (2.32) and (2.33) yields

$$-|x + y| \le |x| - |y| \le |x + y|$$
.

By Lemma 2.43 the above is equivalent to

 $||x| - |y|| \le |x + y|,$

which is the first inequality in (2.20).

An immediate consequence of the triangle inequality are the following inequalities, which are left as an exercise.

Proposition 2.47

For any $x, y \in \mathbb{R}$ it holds

 $||x| - |y|| \le |x - y| \le |x| + |y|.$ (2.34)

Moreover for any $x, y, z \in \mathbb{R}$ it holds

 $|x - y| \le |x - z| + |z - y|$.

Notice that the inequality in (2.34) differs from the triangle inequality (2.20) by a sign. Indeed it can be shown tha (2.20) and (2.34) are equivalent.

2.10 Proofs in Mathematics

In a mathematical proof one needs to show that

$$\alpha \implies \beta$$
 (2.35)

where

- α is a given set of assumptions, or **Hypothesis**
- β is a conclusion, or **Thesis**

Proving (2.35) means convincing ourselves that β follows from α . Common strategies to prove (2.35) are:

1. **Contradiction**: Assume that the thesis is **false**, and hope to reach a contradiction: that is, prove that

 $\neg \beta \implies$ contradiction

where $\neg \beta$ is the **negation** of β .

For example we already proved by contradiction that

Definition of
$$\mathbb{Q} \implies \sqrt{2} \notin \mathbb{Q}$$
,

In the above statement

$$\begin{aligned} \alpha &= (\text{Definition of } \mathbb{Q}) \ . \\ \beta &= \left(\sqrt{2} \notin \mathbb{Q}\right) \ . \end{aligned}$$

Therefore

$$\neg \beta = \left(\sqrt{2} \in \mathbb{Q}\right) \,.$$

- 2. **Direct**: Sometimes proofs will also need **direct** arguments, meaning that one need to show directly that (2.35) holds.
- 3. Contrapositive: The statement (2.35) is equivalent to

$$\neg \beta \implies \neg \alpha \,. \tag{2.36}$$

Thus, instead of proving (2.35), one could show (2.36). The statement (2.36) is called the **contrapositive** of (2.35).

Let us make an example.

Proposition 2.48

Two real numbers *a*, *b* are equal if and only if for every real number $\varepsilon > 0$ it follows that $|a-b| < \varepsilon$.

Before proceeding with the proof, note that the above stetement is just saying that:

Two numbers are equal if and only if they are **arbitrarily** close

By arbitrarily close we mean that they are as close as you want the to be.

```
Proof: of Proposition 2.48
```

Let us first rephrase the statement using mathematical symbols:

Let $a, b \in \mathbb{R}$. Then it holds:

 $a=b \iff |a-b|<\varepsilon\,, \ \forall\,\varepsilon>0\,.$

Setting

$$\begin{aligned} \alpha &= (a = b) \\ \beta &= (|a - b| < \varepsilon, \ \forall \varepsilon > 0) \end{aligned}$$
 (2.37)

the statement is equivalent to

 $\alpha \iff \beta$.

To show the above, it is sufficient to show that

 $\alpha \implies \beta \quad \text{and} \quad \beta \implies \alpha$.

Step 1. Proof that $\alpha \implies \beta$. This proof can be carried out by a **direct** argument. Since we are assuming α , this means

a = b.

We want to see that β holds. Therefore fix an arbitrary $\varepsilon > 0$. This means that ε can be **any** positive number, as long as you fix it. Clearly

$$|a-b| = |0| = 0 < \varepsilon$$

since a = b, |0| = 0, and $\varepsilon > 0$. The above shows that

 $|a-b|<\varepsilon\,.$

As $\varepsilon > 0$ was arbitrary, we have just proven that

 $|a-b|<\varepsilon\,,\,\,\forall\,\varepsilon>0\,,$

meaning that β holds and the proof is concluded. Step 2. Proof that $\beta \implies \alpha$. Let us prove this implication by showing the **contrapositive**

 $\neg \alpha \implies \neg \beta$.

So let us assume $\neg \alpha$ is true. This means that

 $a \neq b$.

We have to see that $\neg \beta$ holds. But $\neg \beta$ means that

 $\exists \varepsilon_0 > 0 \text{ s.t. } |a-b| \ge \varepsilon_0$.

The above is satisfied by choosing

 $\varepsilon_0 := |a-b|,$

since $\varepsilon_0 > 0$ given that $a \neq b$.

2.11 Induction

Another technique for carrying out proofs is **induction**, which we take as an axiom.

Axiom 2.49: Principle of Induction

Let $S \subseteq \mathbb{N}$. Suppose that

- 1. We have $1 \in S$, and
- 2. Whenever $n \in S$, then $(n + 1) \in S$.

Then we have

 $S = \mathbb{N}$.

Important

The above is an **axiom**, meaning that we do not prove it, but rather we just **assume it holds**.

Remark 2.50

It would be possible to prove the Principle of Induction starting from elementary axioms for \mathbb{N} , called the **Peano Axioms**, see the Wikipedia page.

However, in justifying basic principles of mathematics, one at some point needs to draw a line. This means that something which looks elementary needs to be assumed to hold, in order to have a starting point for proving deeper statements.

In the case of the Principle of Induction, the intuition is clear:

The Principle of Induction is just describing the **domino effect**: *If one tile falls, then the next one will fall as well.* Therefore if the *first tile falls, all the tiles will fall.*

It seems reasonable to assume such evident principle.

The Principle of Induction can be used to prove statements which depend on some index $n \in \mathbb{N}$. Precisely, the following statement holds.

Corollary 2.51: Principle of Inducion - Alternative formulation

Let $\alpha(n)$ be a statement which depends on $n \in \mathbb{N}$. Suppose that

- 1. $\alpha(1)$ is true, and
- 2. Whenever $\alpha(n)$ is true, then $\alpha(n + 1)$ is true.

Then $\alpha(n)$ is true for all $n \in \mathbb{N}$.

Numbers Sequences and Series

Proof

Define the set

 $S := \{n \in \mathbb{N} \text{ s.t. } \alpha(n) \text{ is true} \}.$

Then

- 1. We have $1 \in S$, since $\alpha(1)$ is true.
- 2. If $n \in S$ then $\alpha(n)$ is true. By assumption this implies that $\alpha(n + 1)$ is true. Therefore $(n + 1) \in S$.

Therefore S satisfies the assumptions of the Induction Principle and we conclude that

 $S = \mathbb{N}$.

By definition this means that $\alpha(n)$ is true for all $n \in \mathbb{N}$.

Example 2.52: Formula for summing first *n* natural numbers

Question. Prove by induction that the following formula holds for all $n \in \mathbb{N}$:

$$1 + 2 + 3 + \dots + (n - 1) + n = \frac{n(n + 1)}{2}.$$
 (2.39)

Solution. To be really precise, consider the statement

 $\alpha(n)$:= the above formula is true for n.

In order to apply induction, we need to show that

- 1. $\alpha(1)$ is true,
- 2. If $\alpha(n)$ is true then $\alpha(n+1)$ is true.

Let us proceed: Define

 $S(n) = 1 + 2 + \ldots + n.$

This way the formula at (2.39) is equivalent to

$$S(n) = \frac{n(n+1)}{2}, \quad \forall n \in \mathbb{N}.$$

1. It is immediate to check that (2.39) holds for n = 1.

2. Suppose (2.39) holds for n = k. Then

$$S(k+1) = 1 + \dots + k + (k+1)$$
(2.40)

$$= S(k) + (k+1)$$
(2.41)

$$=\frac{k(k+1)}{2} + (k+1) \tag{2.42}$$

$$=\frac{k(k+1)+2(k+1)}{2}$$
(2.43)

$$=\frac{(k+1)(k+2)}{2}$$
(2.44)

where in the first equality we used that (2.39) holds for n = k. We have proven that

$$S(k+1) = \frac{(k+1)(k+2)}{2}$$

The RHS in the above expression is exactly the RHS of (2.39) computed at n = k + 1. Therefore, we have shown that formula (2.39) holds for n = k + 1.

By the Principle of Induction, we then conclude that $\alpha(n)$ is true for all $n \in \mathbb{N}$, which means that (2.39) holds for all $n \in \mathbb{N}$.

Example 2.53: Statements about sequences of numbers

Suppose you are given a collection of numbers

$$\{x_n \text{ s.t. } n \in \mathbb{N}\}.$$

Such collection of numbers is called sequence. Assume that

$$x_1 := 1$$
 (2.45)

$$x_{n+1} := \frac{x_n}{2} + 1.$$
 (2.46)

A sequence defined as above is called **recurrence sequence**. Using the above rule we can compute all the terms of x_n .

Let us compute the first 4 elements:

$$x_1 = 1$$
 (2.47)

$$x_2 = \frac{x_1}{2} + 1 = 1 + \frac{1}{2} = x_1 + \frac{1}{2}$$
(2.48)

$$x_3 = \frac{x_2}{2} + 1 = 1 + \frac{1}{2} + \frac{1}{4} = x_2 + \frac{1}{4}$$
(2.49)

$$x_4 = \frac{x_3}{2} + 1 = 1 + \frac{1}{2} + \frac{1}{4} + \frac{1}{8} = x_3 + \frac{1}{8}$$
(2.50)

(2.51)

We see a pattern: The successive term is obtained by adding a power of 1/2 to the previous term. We therefore conjecture that

$$x_{n+1} = x_n + \frac{1}{2^n} \tag{2.52}$$

We want to prove our conjecture rigorously, i.e. using induction. **Claim**. (2.52) holds for all $n \in \mathbb{N}$. **Proof of Claim.** We argue by induction:

1. We see that

$$x_2 = 1 + \frac{1}{2} = x_1 + \frac{1}{2}$$

proving (2.52) for n = 1.

2. Suppose now that (2.52) holds for n = k. We need to prove that (2.52) holds for n = k + 1, that is,

$$x_{k+2} = x_{k+1} + \frac{1}{2^{k+1}}.$$
 (2.53)

Starting with the definition of x_{k+2} we see that

 $x_{k+2} = \frac{x_{k+1}}{2} + 1$ (definition of x_{k+2}) $= \frac{x_k + \frac{1}{2^k}}{2} + 1$ (inductive hypothesis) $= \frac{x_k}{2} + 1 + \frac{1}{2^{k+1}}$ (just calculations) $= x_{k+1} + \frac{1}{2^{k+1}}$ (definition of x_{k+1})

which proves (2.53).

Therefore the assumptions of the Induction Principle are satisfied, and (2.52) follows.

Example 2.54: Bernoulli's inequality

Question. Let $x \in \mathbb{R}$ with x > -1. Bernoulli's inequality states that

$$(1+x)^n \ge 1 + nx, \quad \forall n \in \mathbb{N}.$$
(2.54)

Prove Bernoulli's inequality by induction.

Solution. Let $x \in \mathbb{R}$, x > -1. We prove the statement by induction:

- Base case: (6.9) holds with equality when n = 1.
- Induction hypothesis: Let $k \in \mathbb{N}$ and suppose that (6.9) holds for n = k, i.e.,

$$(1+x)^k \ge 1+kx.$$

Then

$$(1+x)^{k+1} = (1+x)^k (1+x)$$

$$\ge (1+kx)(1+x)$$

$$= 1+kx+x+kx^2$$

$$\ge 1+(k+1)x,$$

where we used that $kx^2 \ge 0$. Then (6.9) holds for n = k + 1.

By induction we conclude (6.9).

3 Real Numbers

In this chapter we introduce the system of Real Numbers \mathbb{R} and study some of its properties. We will follow quite an abstract approach, which requires the definition of ordered field. The fact that \mathbb{R} is a continuum with no gaps (unlike \mathbb{Q}) will be consequence of the Axiom of Completeness.

Therefore \mathbb{R} will be defined as an ordered field in which the Axiom of Completeness holds.

3.1 Fields

In order to introduce \mathbb{R} , we need the concepts of **binary operation** and **field**. We proceed in a general setting, starting from a set *K*. We will assume that there is an **equivalence relation** on *K* denoted by =.

Definition 3.1: Binary operation

A binary operation on a set *K* is a function

 $\circ \; : \; K \times K \to K$

which maps the ordered pair (x, y) into $x \circ y$.

Notation 3.2

We use the special symbols of + and \cdot to refer to addition and multiplication.

• Addition: The addition, or sum of $x, y \in K$ is denoted by

x + y

• **Multiplication**: The multiplication, or **product** of $x, y \in K$ is denoted by

 $x \cdot y$ or xy

Warning

When we talk about + and \cdot in a set *K* we are **not necessarily** thinking about the usual sum and multiplication in \mathbb{Q} . This is the case in the next Example, where we define

$$1 + 1 = 0$$

Example 3.3: An example of binary operation

Let $K = \{0, 1\}$. We can for example define operations of sum and product on K according to the tables

+	0	1	·	0	1
0	0	1	0	0	0
1	1	0	1	0	1

The above mean that

 $0 + 0 = 1 + 1 = 0, \quad 0 + 1 = 1 + 0 = 1,$ $0 \cdot 0 = 0 \cdot 1 = 1 \cdot 0 = 0, \quad 1 \cdot 1 = 1.$ (3.1)

Note that we have defined (rather controversially!)

$$1 + 1 = 0$$

The other option would have been to define

1 + 1 = 1

This would have been absolutely fine. However we could not have defined

$$1 + 1 = 2$$

because $2 \notin K$.

Note: The operations in (3.1) look more natural if you identify 0 with the even numbers and 1 with the odd numbers. In that case 1 + 1 = 0 reads

Odd + Odd = Even

Indeed all the definitions given in (3.1) agree with such identification, see tables below:

+	Even	Odd	•	Even	Odd
Even	Even	Odd	Even	Even	Even
Odd	Odd	Even	Odd	Even	Odd

Binary operations take ordered pairs of elements of K as input. Therefore the operation

 $x \circ y \circ z$

does not make sense, since we do not know which one between

$$x \circ y$$
 or $y \circ z$

has to be performed first. Moreover the outcome of an operation depends on order:

 $x \circ y \neq y \circ x$.

This motivates the following definition.

Definition 3.4: Properties of binary operations

Let *K* be a set and \circ : $K \times K \rightarrow K$ be a binary operation on *K*. We say that:

1. • is commutative if

$$x \circ y = y \circ x, \quad \forall \, x, \, y \in K$$

2. • is associative if

$$(x \circ y) \circ z = x \circ (y \circ z), \quad \forall x, y, z \in K$$

3. An element $e \in K$ is called **neutral element** of \circ if

$$x \circ e = e \circ x = x, \quad \forall x \in K$$

4. Let *e* be a neutral element of \circ and let $x \in K$. An element $y \in K$ is called an **inverse** of x with respect to \circ if

 $x \circ y = y \circ x = e.$

Example 3.5

Consider Q with the usual operations of sum and multiplication

- We already know that + and \cdot are commutative and associative
- The neutral element of the sum is e = 0
- The neutral element of the product is e = 1
- The additive inverse of x is -x
- The multiplicative inverse of $x \neq 0$ is 1/x.

Example 3.6

Let *K* with + and \cdot be as in Example 3.3. The sum satisfies:

• + is commutative, since

0 + 1 = 1 + 0 = 0.

• + is associative, since for example

$$(0+1)+1 = 1+1 = 0$$
, $0+(1+1) = 0+0 = 0$,

and therefore

$$(0+1)+1=0+(1+1).$$

In general one can show that + is associative by checking all the other permutations. • The neutral element of + is 0, since

$$0 + 0 = 0$$
, $1 + 0 = 0 + 1 = 1$.

• Every element has an inverse. Indeed, the inverse of 0 is 0, since

$$0 + 0 = 0$$
,

while the inverse of 1 is 1, since

$$1 + 1 = 1 + 1 = 0$$
.

The multiplication satisfies:

- \cdot is commutative, since

$$1\cdot 0=0\cdot 1=0\,.$$

• · is associative, since for example

$$(0 \cdot 1) \cdot 1 = 0 \cdot 1 = 0, \quad 0 \cdot (1 \cdot 1) = 0 \cdot 1 = 0,$$

and therefore

$$(0\cdot 1)\cdot 1=0\cdot (1\cdot 1).$$

By checking all the other permutations one can show that \cdot is associative.

- The neutral element of \cdot is 1, since

$$0 \cdot 1 = 1 \cdot 0 = 0$$
, $1 \cdot 1 = 1$.

• The element 0 has no inverse, since

$$0 \cdot 0 = 0 \cdot 1 = 1 \cdot 0 = 0,$$

and thus we never obtain the neutral element 1. The inverse of 1 is given by 1, since

 $1\cdot 1=1\,.$

Example 3.7

Question. Let $K = \{0, 1\}$ be a set with binary operation \circ defined by the table

1. Is • commutative? Justify your answer.

2. Is • associative? Justify your answer.

Solution.

1. The operation • is not commutative, since

 $0 \circ 1 = 1 \neq 0 = 1 \circ 0.$

2. The operation \circ is not associative, since

 $(0 \circ 1) \circ 1 = 1 \circ 1 = 0,$

while

 $0 \circ (1 \circ 1) = 0 \circ 0 = 1,$

so that

$$(0 \circ 1) \circ 1 \neq 0 \circ (1 \circ 1).$$

We are ready to define fields.

Definition 3.8: Field

Let *K* be a set with binary operations of **addition**

 $+ : K \times K \to K, \quad (x, y) \mapsto x + y$

and multiplication

 $\cdot : K \times K \to K, \quad (x, y) \mapsto x \cdot y = xy.$

We call the triple $(K, +, \cdot)$ a **field** if:

- 1. The addition + satisfies: $\forall x, y, z \in K$
 - (A1) Commutativity and Associativity:

$$x + y = y + x$$
$$(x + y) + z = x + (y + z)$$

• (A2) Additive Identity: There exists a neutral element in *K* for +, which we call 0. It holds:

$$x + 0 = 0 + x = x$$

• (A₃) **Additive Inverse**: There exists an **inverse** of *x* with respect to +. We call this element the **additive inverse** of *x* and denote it by -x. It holds

$$x + (-x) = (-x) + x = 0$$

- 2. The multiplication \cdot satisifes: $\forall x, y, z \in K$
 - (M1) Commutativity and Associativity:

$$x \cdot y = y \cdot x$$
$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

• (M2) **Multiplicative Identity**: There exists a **neutral element** in K for \cdot , which we call 1. It holds:

$$x \cdot 1 = 1 \cdot x = x$$

• (M₃) **Multiplicative Inverse**: If $x \neq 0$ there exists an **inverse** of x with respect to \cdot . We call this element the **multiplicative inverse** of x and denote it by x^{-1} . It holds

$$x \cdot x^{-1} = x^{-1} \cdot x = 1$$

- 3. The operations + and \cdot are related by
 - (AM) **Distributive Property**: $\forall x, y, z \in K$

$$x \cdot (y+z) = (x \cdot y) + (y \cdot z).$$

Remark 3.9

Since fields are an abstract model for \mathbb{Q} , we have chosen to denote the two field operations with + and \cdot . We could have denoted them with other symbols, e.g. \clubsuit and \heartsuit . In this case (AM) would read

$$x \clubsuit (y \heartsuit z) = (x \clubsuit y) \heartsuit (y \clubsuit z)$$

This would of course still make sense. It would just be a lot harder to write calculations and to remember which operation is which!

Warning

It might be tempting to dismiss the definition of field, and to think that all fields look like Q. This is not the case, as seen in the following Example.

Example 3.10

Question. Let *K* with + and \cdot be as in Example 3.3, that is, *K* = {0, 1} with operations defined by

+	0	1	·	0	1
0	0	1	0	0	0
1	1	0	1	0	1

Prove that $(K, +, \cdot)$ is a field. **Solution.** We have already shown in Example 3.6 that:

- (A1) and (M1) hold,
- (A2) holds with neutral element 0,
- (M2) holds with neutral element 1,
- (A₃) every element has an additive inverse, with

$$-0 = 0$$
, $-1 = 1$,

• (M3) every element which is not 0 a multiplicative inverse, with

$$1^{-1} = 1$$

We are left to show the Distributive Property (AM). Indeed:

• (AM) For all $y, z \in K$ we have

$$0 \cdot (y+z) = 0, \quad (0 \cdot y) + (0 \cdot z) = 0 + 0 = 0,$$

and also

$$1 \cdot (y+z) = y+z$$
, $(1 \cdot y) + (1 \cdot z) = y+z$.

Thus (AM) holds.

Definition 3.11: Subtraction and division

Let $(K, +, \cdot)$ be a field. We define:

1. **Subtraction** as the operation – defined by

$$x-y := x + (-y), \quad \forall x, y \in K,$$

where -y is the additive inverse of y.

2. Division as the operation / defined by

$$x/y := x \cdot y^{-1}, \quad \forall x, y \in K, \ y \neq 0,$$

where y^{-1} is the multiplicative inverse of *y*.

Warning

One might wonder why we are defining subtraction and division. Are they not already defined in \mathbb{Q} ? Yes they are. But they make no sense in a general field $(K, +, \cdot)$, unless we define them explicitly.

Using the field axioms we can prove the following properties.

Proposition 3.12: Uniqueness of neutral elements and inverses

Let $(K, +, \cdot)$ be a field. Then

- 1. There is a unique element in K with the property of 0,
- 2. There is a unique element in *K* with the property of 1,
- 3. For all $x \in K$ there is a unique additive inverse -x,
- 4. For all $x \in K$, $x \neq 0$, there is a unique multiplicative inverse x^{-1} .

Proof

1. Suppose that $0 \in K$ and $\tilde{0} \in K$ are both neutral element of +, that is, they both satisfy (A2). Then

 $0 + \tilde{0} = 0$

since $\tilde{0}$ is a neutral element for +. Moreover

$$\tilde{0} = 0 + \tilde{0}$$

since 0 is a neutral element for +. By commutativity of +, see property (A1), we have

$$0 = 0 + \tilde{0} = \tilde{0} + 0 = \tilde{0}$$
,

showing that $0 = \tilde{0}$. Hence the neutral element for + is unique.

- 2. Exercise.
- 3. Let $x \in K$ and suppose that $y, \tilde{y} \in K$ are both additive inverses of x, that is, they both satisfy (A₃). Therefore

$$x + y = 0$$

since y is an additive inverse of x and

$$x + \tilde{y} = 0$$

since \tilde{y} is an additive inverse of *x*. Therefore we can use commutativity and associativity and of +, see property (A1), and the fact that 0 is the neutral element of +, to infer

$$y = y + 0 = y + (x + \tilde{y})$$

= $(y + x) + \tilde{y} = (x + y) + \tilde{y}$
= $0 + \tilde{y} = \tilde{y}$,

concluding that $y = \tilde{y}$. Thus there is a unique additive inverse of *x*, and

 $y=\tilde{y}=-x\,,$

with -x the element from property (A₃).

4. Exercise.

Using the properties of field we can also show that the usual properties of sum, subtraction, multiplication and division still hold in any field. We list such properties in the following proposition.

Proposition 3.13: Properties of field operations

Let $(K, +, \cdot)$ be a field. Then for all $x, y, z \in K$,

```
• x + y = x + z \implies y = z

• x \cdot y = x \cdot z and x \neq 0 \implies y = z

• -0 = 0

• 1^{-1} = 1

• x \cdot 0 = 0

• -1 \cdot x = -x

• -(-x) = x

• (x^{-1})^{-1} = x if x \neq 0

• (x \cdot y)^{-1} = x^{-1} \cdot y^{-1}
```

The above properties can be all proven with elementary use of the field properties (A1)-(A3), (M1)-(M3) and (AM). This is an exercise in patience, and is left to the reader.

Let us conclude with examining the sets of numbers introduced in Chapter 1.

Theorem 3.14

Consider the sets $\mathbb{N}, \mathbb{Z}, \mathbb{Q}$ with the usual operations + and $\cdot.$ We have:

- $(\mathbb{N}, +, \cdot)$ is not a field.
- $(\mathbb{Z}, +, \cdot)$ is not a field.
- $(\mathbb{Q}, +, \cdot)$ is a field.

Proof

- (N, +, ·) is not a field: It satisfies properties (A1), (A2), (M1), (M2), (AM) of fields. It is missing properties (A3) and (M3), the additive and multiplicative inverse properties, respectively.
- (ℤ, +, ·) is not a field: It satisfies properties (A1), (A2), (A3), (M1), (M2), (AM) of fields. Thus it is only missing (M3), the multiplicative inverse property.
- $(Q, +, \cdot)$ is a field. This is clear: The field axioms are introduced exactly to model Q. Therefore, Q is trivially a field.

3.2 Ordered fields

Definition 3.15: Ordered field

Let *K* be a set with binary operations + and \cdot , and with an order relation \leq . We call $(K, +, \cdot, \leq)$ an **ordered field** if:

- 1. $(K, +, \cdot)$ is a field
- 2. There \leq is of **total order** on *K*: $\forall x, y, z \in K$
 - (O1) Reflexivity:

 $x \le x$

• (O2) Antisymmetry:

 $x \le y$ and $y \le x \implies x = y$

• (O₃) Transitivity:

 $x \leq y$ and $y \leq z \implies x = z$

• (O₄) Total order:

- $x \le y$ or $y \le x$
- 3. The operations + and ·, and the total order \leq , are related by the following properties: $\forall x, y, z \in K$
 - (AM) Distributive: Relates addition and multiplication via

 $x \cdot (y+z) = x \cdot y + x \cdot z$

• (AO) Relates addition and order with the requirement:

 $x \le y \implies x+z \le y+z$

• (MO) Relates multiplication and order with the requirement:

$$x \ge 0, \ y \ge 0 \implies x \cdot y \ge 0$$

Theorem 3.16

 $(\mathbb{Q}, +, \cdot, \leq)$ is an **ordered field**.

3.3 Cut Property

We have just introduced the notion of **fields** and **ordered fields**. We noted that the set of rational numbers with the usual operations of sum and multiplication

 $(\mathbb{Q}, +, \cdot)$

is a field. Moreover it is an ordered field with the usual order $\leq.$

We now need to address the key issue we proved in Chapter 1, namely, the fact that

 $\sqrt{2} \notin \mathbb{Q}$.

Intuitively, this means that Q has **gaps**, and cannot be represented as a **continuous** line. The rigorous definition of **lack of gaps** needs the concept of **cut** of a set. This, in turn, needs the concept of **partition**.

Definition 3.17: Partition of a set

Let *S* be a non-empty set. The pair (A, B) is a **partition** of *S* if

$$A, B \subseteq S, \quad A \neq \emptyset, \quad B \neq \emptyset,$$

and

 $S=A\cup B\,,\quad A\cap B=\emptyset\,.$



Figure 3.1: Schematic picture of a partition (A, B) of the set K.

Definition 3.18: Cut of a set

Let *S* be a non-empty set with a total order relation \leq . The pair (*A*, *B*) is a **cut** of *S* if

- 1. (*A*, *B*) is a **partition** of *S*,
- 2. We have

 $a \leq b$, $\forall a \in A$, $\forall b \in B$.

The **cut** of a set is often called **Dedekind cut**, named after **Richard Dedekind**, who used cuts to give an explicit construction of the real numbers **R**, see Wikipedia page.

Definition 3.19: Cut property

Let *S* be a non-empty set with a total order relation \leq . We say that *S* has the **cut property** if for every cut (*A*, *B*) of *S* there exists some $s \in S$ such that

 $a \leq s \leq b$, $\forall a \in A$, $\forall b \in B$.

We call *s* the **separator** of the cut (*A*, *B*).

Example 3.20

Let $S = \mathbb{Q}$ and consider the sets

$$A = (-\infty, s] \cap \mathbb{Q}, \quad B = (s, \infty) \cap \mathbb{Q}.$$

for some $s \in \mathbb{Q}$. Then the pair (A, B) is a cut of \mathbb{Q} , and s is the separator.

$$A = (-\infty, s] \cap \mathbb{Q} \qquad \qquad B = (s, \infty) \cap \mathbb{Q}$$



Question 3.21

Do all ordered fields have the Cut Property? Does Q have the Cut Property?

The answer to the above question is NO. For example the pair

$$A = (-\infty, \sqrt{2}) \cap \mathbb{Q}, \quad B = (\sqrt{2}, \infty) \cap \mathbb{Q}.$$
(3.2)

is a cut of \mathbb{Q} , since $\sqrt{2} \notin \mathbb{Q}$. However what is the separator? It should be $s = \sqrt{2}$, given that clearly

$$a \le \sqrt{2} \le b$$
, $\forall a \in A, \forall b \in B$.

However $\sqrt{2} \notin \mathbb{Q}$, so we are **NOT ALLOWED** to take it as separator. Indeed, we can show that (A, B) defined as in (3.2) has no separator.

$$\begin{array}{c|c} A=(-\infty,\sqrt{2})\cap\mathbb{Q} & B=(\sqrt{2},\infty)\cap\mathbb{Q} \\ \\ \hline & & \\ \sqrt{2}\notin\mathbb{Q} \end{array}$$

Figure 3.3: (A, B) is a cut of \mathbb{Q} which has no separator

Theorem 3.22: Q does not have the cut property.

 \mathbb{Q} does not have the cut property. More explicitly, there exist a cut (*A*, *B*) of \mathbb{Q} which has no separator.

Remark 3.23: Ideas for the proof of Theorem 3.22

Before proceeding with the proof, let us summarize the ideas behind it: We will consider the cut (A, B) in (3.2). We then assume by contradiction that (A, B) admits a separator $L \in \mathbb{Q}$, so that

 $a \le L \le b, \quad \forall a \in A, \ \forall b \in B.$ (3.3)

Since (A, B) is a partition of \mathbb{Q} , then either $L \in A$ or $L \in B$. These will both lead to a contradiction:

• If $L \in A$, by definition of A we have

 $L < \sqrt{2}$.

We want to contradict the fact that *L* is a separator for the cut (*A*, *B*). The idea is that, since $\sqrt{2} \notin \mathbb{Q}$, it is possible to squeeze a rational number $\tilde{L} \in \mathbb{Q}$ in between *L* and $\sqrt{2}$, i.e.

 $L < \tilde{L} < \sqrt{2}$.

How do we find such \tilde{L} in practice? We look for a number \tilde{L}_n of the form

$$\tilde{L}_n = L + \frac{1}{n}$$

for some $n \in \mathbb{N}$ to be suitably chosen later. Clearly $\tilde{L}_n \in \mathbb{Q}$ and

 $L < \tilde{L}_n$

for all $n \in \mathbb{N}$. We will then be able to find $n_0 \in \mathbb{N}$ such that

$$L < \tilde{L}_{n_0} < \sqrt{2} \,. \tag{3.4}$$

Now comes the contradiction: From (3.4) we see that $\tilde{L}_{n_0} \in A$. However *L* is a separator, and so from (3.3) we have

$$\tilde{L}_{n_0} \leq L$$
,

which contradicts (3.4).

• If $L \in B$, by definition of B we have

 $\sqrt{2} < L$.

The idea is the same as above: Since $\sqrt{2} \notin \mathbb{Q}$, we can squeeze a rational number $\tilde{L} \in \mathbb{Q}$ between $\sqrt{2}$ and L, i.e.,

 $\sqrt{2} < \tilde{L} < L \,.$

Since we want \tilde{L} to be a rational number smaller than *L*, we look for \tilde{L} of the form

$$\tilde{L}_n := L - \frac{1}{n}$$

for a suitable $n \in \mathbb{N}$. Clearly $\tilde{L}_n \in \mathbb{Q}$ and

 $\tilde{L}_n < L$,

for all $n \in \mathbb{N}$. We will be able to find $n_0 \in \mathbb{N}$ such that

$$\sqrt{2} < \tilde{L}_{n_0} < L. \tag{3.5}$$

Therefore $\tilde{L}_{n_0} \in B$. Again, contradiction: *L* is a separator and so

 $L \leq \tilde{L}_{n_0}$,

which contradicts (3.5).

Both cases $L \in A$ or $L \in B$ lead to a contradiction. Since these are all the possibilities, we conclude that the cut (A, B) has no separator in \mathbb{Q} .

Time to make the ideas in the above remark rigorous. Two main issues need fixing:

1. $\sqrt{2}$ is just a symbol for a number x such that $x^2 = 2$. As $\sqrt{2} \notin \mathbb{Q}$, what is the meaning of the expression

 $a < \sqrt{2} < b$

when $a, b \in \mathbb{Q}$? We can only compare rational numbers with rational numbers, so the above inequalities are meaningless. We need a more clever way to write down the sets *A* and *B* so that they make sense as objects in \mathbb{Q} .

2. We said it is possible to find $n_0 \in \mathbb{N}$ such that

 $L < \tilde{L}_{n_0} < \sqrt{2} \quad \text{ or } \quad \sqrt{2} < \tilde{L}_{n_0} < L$

We need to prove it!

Proof: Proof of Theorem 3.22

Let A and B be the sets defined in (3.2). It is useful to rewrite A and B in the form

 $A = A_1 \cup A_2,$

where

$$A_1 = \{ q \in \mathbb{Q} : q < 0 \},\$$
$$A_2 = \{ q \in \mathbb{Q} : q \ge 0, q^2 < 2 \}$$

and

$$B = \{q \in \mathbb{Q} : q > 0, q^2 > 2\}.$$

Step 1. (*A*, *B*) *is a cut of* Q: We need to prove the following:

1. (A, B) is a partition of \mathbb{Q} . This is because $A, B \subseteq \mathbb{Q}$ with $A \neq \emptyset$ and $B \neq \emptyset$. Moreover $A \cap B = \emptyset$ and

$$A \cup B = \mathbb{Q}$$
,

given that $\sqrt{2} \notin \mathbb{Q}$, and so there is no element $q \in \mathbb{Q}$ such that $q^2 = 2$. 2. It holds

$$a \leq b$$
, $\forall a \in A$, $\forall b \in B$

Indeed, suppose that $a \in A$ and $b \in B$. We have two cases:

• $a \in A_1$: Therefore a < 0. In particular

$$a < 0 < b$$

given that b > 0 for all $b \in B$. Thus a < b.

• $a \in A_2$: Therefore $a \ge 0$ and $a^2 < 2$. In particular

 $a^2 < 2 < b^2$,

since $b^2 > 2$ for all $b \in B$. In particular

 $a^2 < b^2 \, .$

Since
$$b > 0$$
 for all $b \in B$, from the above inequality we infer $a < b$, concluding.

Step 2. (*A*, *B*) *has no separator*: Suppose by contradiction that (*A*, *B*) admits a separator

$$L \in \mathbb{Q}$$
.

By definition this means

 $a \le L \le b$, $\forall a \in A, \forall b \in B$. (3.6)

Since

$$L \in \mathbb{Q}, \quad \mathbb{Q} = A \cup B, \quad A \cap B = \emptyset$$

then either $L \in A$ or $L \in B$. We will see that both these possibilities lead to a contradiction: *Case 1:* $L \in A$. By (3.6) we know that

$$a \le L, \quad \forall a \in A.$$
 (3.7)

In particular the above implies

 $L \ge 0 \tag{3.8}$

since $0 \in A$. Therefore we must have $L \in A_2$, that is,

$$L \ge 0 \text{ and } L^2 < 2.$$
 (3.9)

Set

$$\tilde{L} := L + \frac{1}{n}$$

for $n \in \mathbb{N}$, $n \neq 0$ to be chosen later. Clearly we have

$$\tilde{L} \in \mathbb{Q} \quad \text{and} \quad L < \tilde{L} \,.$$
 (3.10)

From
$$(3.8)$$
 and (3.10) we have also

$$\tilde{L} > 0. \tag{3.11}$$

We now want to show that there is a choice of *n* such that $\tilde{L}^2 < 2$, which will lead to a contradiction. Indeed, we can estimate

$$\begin{split} \tilde{L}^2 &= \left(L + \frac{1}{n}\right)^2 \\ &= L^2 + \frac{1}{n^2} + 2\frac{L}{n} \\ &< L^2 + \frac{1}{n} + 2\frac{L}{n} \qquad \left(\text{using } \frac{1}{n} < \frac{1}{n^2}\right) \\ &= L^2 + \frac{2L + 1}{n} \,. \end{split}$$

If we now impose that

$$L^2 + \frac{2L+1}{n} < 2\,,$$

we can rearrange the above and obtain

$$n(2-L^2) > 2L+1$$
.

Now note that $L^2 < 2$ by assumption (3.9). Thus we can divived by $(2 - L^2)$ and obtain

$$n > \frac{2L+1}{2-L^2}$$
.

Therefore we have just shown that

$$n > \frac{2L+1}{2-L^2} \implies \tilde{L}^2 < 2.$$

Together with (3.11) this implies $\tilde{L} \in A$. Therefore we have

 $\tilde{L} \leq L$

by (3.7). On the other hand it also holds

 $\tilde{L} > L$

by (3.10), and therefore we have a contradiction. Thus $L \notin A$. *Case 2:* $L \in B$. As $L \in B$, we have by definition

$$L > 0, \quad L^2 > 2.$$
 (3.12)

Moreover since L is a separator, see (3.6), in particular

$$L \le b , \ \forall b \in B. \tag{3.13}$$

Define now

$$\tilde{L} := L - \frac{1}{n}$$

with $n \in \mathbb{N}$, $n \neq 0$ to be chosen later. Clearly we have

$$\tilde{L} \in \mathbb{Q}, \quad \tilde{L} < L.$$
 (3.14)

We now show that *n* can be chosen so that $\tilde{L} \in B$. Indeed

$$\begin{split} \tilde{L}^2 &= \left(L - \frac{1}{n}\right)^2 \\ &= L^2 + \frac{1}{n^2} - 2\frac{L}{n} \\ &> L^2 - \frac{1}{n^2} - 2\frac{L}{n} \qquad \left(\text{using } \frac{1}{n^2} > -\frac{1}{n^2}\right) \\ &> L^2 - \frac{1}{n} - 2\frac{L}{n} \qquad \left(\text{using } -\frac{1}{n^2} > -\frac{1}{n}\right) \\ &= L^2 - \frac{1 + 2L}{n} \,. \end{split}$$

Now we impose

$$L^2 - \frac{1+2L}{n} > 2$$

which is equivalent to

$$n(L^2-2) > 1+2L$$
.

Since we are assuming $L \in B$, then $L^2 > 2$, see (3.12). Therefore we can divide by $(L^2 - 2)$ and get

$$n > \frac{1+2L}{L^2-2}$$

In total, we have just shown that

$$n > \frac{1+2L}{L^2-2} \quad \Longrightarrow \quad \tilde{L}^2 > 2 \,,$$

proving that $\tilde{L} \in B$. Therefore by (3.13) we get

 $L\leq \tilde{L}\,.$

This contradicts (3.14).

Conclusion:

We have seen that assuming that (A, B) has a separator $L \in \mathbb{Q}$ leads to a contradiction. Thus the cut (A, B) has no separator.

Remark 3.24

The above proof is also suggesting the following observations:

• The set

$$A = (-\infty, \sqrt{2}) \cap \mathbb{Q}$$

does not admit a largest element in \mathbbm{Q}

• The set

 $B = (\sqrt{2}, \infty) \cap \mathbb{Q}$

does not admit a **lowest element** in Q.

It turns out that the **largest** and **lowest** element play a crucial role in analysis. We will give precise definitions in the next section.

3.4 Supremum and infimum

A crucial definition in Analysis is the one of supremum or infimum of a set. This is also another way of studying the **gaps** of Q.

Example 3.25: Intuition about supremum and infimum

Consider the set

 $A = [0, 1) \cap \mathbb{Q}.$

Intuitively, we understand that A is bounded, i.e. not infinite. We also see that

- 0 is the lowest element of A
- 1 is the highest element of A

However we see that $0 \in A$ while $1 \notin A$. We will see that

- 0 can be defined as the **infimum** and **minimum** of *A*.
- 1 can be defined as the **supremum**, but not **maximum**, of *A*.

3.4.1 Upper bound, supremum, maximum

We start by defining the supremum. First we need the notion of upper bound of a set. In the following we assume that $(K, +, \cdot, \leq)$ is an ordered field.

Definition 3.26: Upper bound and bounded above

Let $A \subseteq K$:

1. We say that $b \in K$ is an **upper bound** for A if

 $a \leq b$, $\forall a \in A$.

2. We say that A is **bounded above** if there exists and upper bound $b \in K$ for A.

Definition 3.27: Supremum

Let $A \subseteq K$. A number $s \in K$ is called **least upper bound** or **supremum** of A if:

- 1. s is an upper bound for A,
- 2. *s* is the smallest upper bound of *A*, that is,

If $b \in K$ is upper bound for A then $s \leq b$.

If it exists, the supremum is denoted by

 $s := \sup A$.

Remark 3.28

Note that if a set $A \subseteq K$ in **NOT** bounded above, then the supremum does not exist, as there are no upper bounds of A.

Proposition 3.29: Uniqueness of the supremum

Let $A \subseteq K$. If sup *A* exists, then it is unique.

Proof

Suppose there exist $s_1, s_2 \in K$ such that

$$s_1 = \sup A, \ s_2 = \sup A.$$

Then:

• Since $s_2 = \sup A$, in particular s_2 is an upper bound for A. Since $s_1 = \sup A$ then s_1 is the lowest upper bound. Thus we get

 $s_1 \leq s_2$.

• Exchanging the roles s_1 and s_2 in the above reasoning we also get

 $s_2 \leq s_1$.

This shows $s_1 = s_2$.

Warning

It could happen that:

- A set can have infinite upper bounds,
- The supremum does not belong to the set.

For example

$$A = [0, 1) \cap \mathbb{Q}$$

has for upper bounds all the numbers $b \in \mathbb{Q}$ with b > 1. Moreover one can show that

$$\sup A = 1,$$

and so

 $\sup A \notin A$.

Warning

The supremum does not exist in general. For example let

 $A = [0, \sqrt{2}) \cap \mathbb{Q}.$

We will show that $\sup A$ does not exist in Q. Indeed we will have that

 $\sup A = \sqrt{2} \in \mathbb{R}.$



Figure 3.4: Supremum and upper bounds of a set A in the field K

Definition 3.30: Maximum

Let $A \subseteq K$. A number $M \in K$ is called the **maximum** of A if:

 $M \in A$ and $a \leq M$, $\forall a \in A$.

If it exists, we denote the maximum by

 $M = \max A$.

Proposition 3.31: Relationship between Max and Sup

Let $A \subseteq K$. If the maximum of A exists, then also the supremum exists, and

 $\sup A = \max A$.

Proof

Let

 $M = \max A$.

Then:

• By definition we have $M \in A$ and

 $a \leq M$, $\forall a \in A$.

In particular the above tells us that M is an upper bound of A.

• We claim that M is the least upper bound. Indeed, suppose b is an upper bound of A, that is,

$$a \leq b$$
, $\forall a \in A$.

In particular, since $M \in A$, by the above condition we have

 $M \leq b$.

Therefore *M* is the least upper bound of *A*, meaning that $M = \sup A$.

Warning

The converse of the above statement is false: In general the sup might exist while the max does not. For example

$$A = [0, 1) \cap \mathbb{Q}$$

is such that

 $\sup A = 1$

but max A does not exist. Instead for the set

 $B = [0, 1] \cap \mathbb{Q}$

we have that

 $\max A = \sup A = 1.$

3.4.2 Lower bound, infimum, minimum

We now introduce the definitions of lower bound, infimum, minimum. These are the counterpart of upper bound, supremum and maximum, respectively. In the following $(K, +, \cdot, \leq)$ is an ordered field.

Definition 3.32: Lower bound, bounded below, infimum, minimum

Let $A \subseteq K$:

1. We say that $l \in K$ is a **lower bound** for A if

 $l \leq a$, $\forall a \in A$.

- 2. We say that *A* is **bounded below** if there exists a lower bound $l \in K$ for *A*.
- 3. We say that $i \in K$ is the **greatest lower bound** or **infimum** of A if:

- *i* is a lower bound for *A*,
- *i* is the largest lower bound of *A*, that is,

If $l \in K$ is a lower bound for A then $l \leq i$.

If it exists, the infimum is denoted by

 $i = \inf A$.

4. We say that $m \in K$ is the **minimum** of A if:

 $m \in A$ and $m \leq a$, $\forall a \in A$.

If it exists, we denote the minimum by

$$m = \min A$$
.





Proposition 3.33

Let $A \subseteq K$:

- 1. If inf *A* exists, then it is unique.
- 2. If the minimum of A exists, then also the infimum exists, and

 $\inf A = \min A \,.$

The proof is similar to the ones of Propositions 3.29 and 2.42, and is left to the reader as an exercise.

Warning

It could happen that:

- A set has infinite lower bounds,
- The infimum does not belong to the set.

For example

 $A = (0,1) \cap \mathbb{Q}$

has for lower bounds all the numbers $b \in \mathbb{Q}$ with b < 1. Moreover we will show that

 $\inf A = 0$,

and so

 $\inf A \notin A$.

Warning

The infimum does not exist in general. For example let

 $A = (\sqrt{2}, 5] \cap \mathbb{Q}.$

We will show that $\inf A$ does not exist in Q. Indeed we will have that

 $\inf A = \sqrt{2} \in \mathbb{R}$.

Warning

In general the inf might exist while the min does not. For example

 $A = (0, 1) \cap \mathbb{Q}$

is such that

 $\inf A = 0$

but min A does not exist. Instead for the set

 $B = [0, 1] \cap \mathbb{Q}$

we have that

```
\inf A = \min A = 0.
```

Proposition 3.34

Let $A \subseteq K$. If $\inf A$ and $\sup A$ exist, then

 $\inf A \le a \le \sup A \,, \quad \forall \, a \in A \,.$

The proof is trivial, and is left as an exercise. We now have a complete picture about supremum and infimum, see Figure 3.6.

We conclude with another simple, but useful proposition. The proof is again left to the reader.


Figure 3.6: Supremum, upper bounds, infimum and lower bounds of a set A in K

Proposition 3.35: Relationship between sup and inf

Let $A \subseteq K$. Define

 $-A := \{-a : a \in A\}.$

They hold:

1. If $\sup A$ exists, then $\inf A$ exists and

 $\inf(-A) = -\sup A.$

2. If $\inf A$ exists, then $\sup A$ exists and

 $\sup(-A) = -\inf A.$

3.5 Completeness

In this section $(K, +, \cdot, \leq)$ denotes an ordered field.

Question 3.36

Let $A \subseteq K$ be non-empty and bounded above. Does

 $\sup A$

always exist?

The answer to the above question is **NO**. Like we did with the Cut Property, the counterexample can be found in the set of rational numbers \mathbb{Q} . A set bounded above for which the supremum does nor exist is, for example,

$$A = [0, \sqrt{2}) \cap \mathbb{Q}. \tag{3.15}$$

Theorem 3.37

There exists a set $A \subseteq \mathbb{Q}$ such that

• A is non-empty,

- A is bounded above,
- $\sup A$ does not exist in \mathbb{Q} .

The proof uses similar ideas to those employed in demonstrating that Q does not satisfy the Cut Property, see the proof of Theorem 3.22.

Proof

Define the set

$$A = \{q \in \mathbb{Q} : q \ge 0, q^2 < 2\}.$$

Notice that *A* formally coincides with the set in (3.15). *Step 1. A is bounded above.* Take b := 3. Then *b* is an upper bound for *A*. Indeed by definition

 $q^2 < 2\,, \ q \ge 0\,, \ \forall \, q \in A\,.$

Therefore

$$q^2 < 2 < 9 \implies q^2 < 9 \implies q < 3 = b$$

for each $q \in A$, showing that b = 3 is an upper bound for A. Step 2. The supremum of A does not exist in \mathbb{Q} . Assume by contradiction that there exists

$$s = \sup A \in \mathbb{Q}$$

By definition it holds

$$q \le s, \quad \forall q \in A \tag{3.16}$$

$$q \le b, \forall q \in A \implies s \le b$$
 (3.17)

There are two possibilities: $s \in A$ or $s \notin A$. *Case 1.* If $s \in A$, by definition of A we have

$$s \ge 0, \quad s^2 < 2.$$
 (3.18)

Define

$$\tilde{s} := s + \frac{1}{n}$$

with $n \in \mathbb{N}$, $n \neq 0$ to be chosen later. Then

$$\tilde{s}^{2} = \left(s + \frac{1}{n}\right)^{2}$$

= $s^{2} + \frac{1}{n^{2}} + 2\frac{s}{n}$
< $s^{2} + \frac{1}{n} + 2\frac{s}{n}$ (using $\frac{1}{n} < \frac{1}{n^{2}}$)
= $s^{2} + \frac{2s+1}{n}$.

If we now impose that

$$s^2 + \frac{2s+1}{n} < 2\,,$$

we can rearrange the above and obtain

$$n(2-s^2) > 2s+1$$
.

Now note that $s^2 < 2$ by assumption (3.18). Thus we can divide by $(2 - s^2)$ and obtain

$$n > \frac{2s+1}{2-s^2}$$

To summarize, we have just shown that

$$n > \frac{2s+1}{2-s^2} \implies \tilde{s}^2 < 2.$$

 $\tilde{s} > s > 0$,

 $\tilde{s} \in A$.

 $\tilde{s} \leq s$.

Moreover $\tilde{s} := (s + 1/n) \in \mathbb{Q}$ and

showing that

Since $s = \sup A$, we then have

However

$$\tilde{s} := s + \frac{1}{n} > s,$$

yielding a contradiction. Thus $s \in A$ is not possible. *Case 2.* Assume $s \notin A$. Since $s = \sup A$ and $0 \in A$, we deduce that

s > 0.

In particular $s \notin A$ implies that

 $s^2 > 2$.

(3.19)

Define

 $\tilde{s} := s - \frac{1}{n}$.

We have

$$\tilde{s}^{2} = \left(s - \frac{1}{n}\right)^{2}$$

$$= s^{2} + \frac{1}{n^{2}} - 2\frac{s}{n}$$

$$> s^{2} - \frac{1}{n^{2}} - 2\frac{s}{n} \qquad \left(\text{using } \frac{1}{n^{2}} > -\frac{1}{n^{2}}\right)$$

$$> s^{2} - \frac{1}{n} - 2\frac{s}{n} \qquad \left(\text{using } -\frac{1}{n^{2}} > -\frac{1}{n}\right)$$

$$= s^{2} - \frac{1 + 2s}{n}.$$

Now we impose

$$s^2 - \frac{1+2s}{n} > 2$$

which is equivalent to

$$n(s^2-2) > 1+2s$$
.

By (3.19) we have $s^2 > 2$. Therefore we can divide by $(s^2 - 2)$ and get

$$n > \frac{1+2s}{s^2-2}$$

In summary, we have just shown that

$$n > \frac{1+2s}{s^2-2} \implies \tilde{s}^2 > 2$$
.

We also want to choose n in such a way that

 $\tilde{s} > 0$

This means

$$0 < \tilde{s} = s - \frac{1}{n} \quad \iff \quad n > \frac{1}{s}$$

Therefore

$$n > \min\left\{\frac{1+2s}{s^2-2}, \frac{1}{s}\right\} \quad \Longrightarrow \quad \tilde{s} > 0, \quad \tilde{s}^2 > 2$$

In particular $\tilde{s} \notin A$, and by definition of A we have

 $\tilde{s} \ge q$, $\forall q \in A$.

Indeed, assume $\tilde{s} < q$ for some $q \in A$. Since $\tilde{s} > 0$, we can take the squares and deduce that $\tilde{s}^2 < q^2 < 2$, which is a contradiction.

By definition we have $\tilde{s} = (s - 1/n) \in \mathbb{Q}$. Therefore \tilde{s} is an upper bound for A in \mathbb{Q} . Since $s = \sup A$ is the smallest upper bound, see (3.17), it follows that

$$s \leq \tilde{s}$$
.

However

$$\tilde{s} := s - \frac{1}{n} < s \, ,$$

obtaining a contradiction. Then $s \notin A$.

Conclusion. We have assumed by contradiction that $s = \sup A$ exists in \mathbb{Q} . In this case either $s \in A$ or $s \notin A$. In both instances we found a contradiction. Therefore $\sup A$ does not exist in \mathbb{Q} .

The above theorem shows that the supremum does not necessarily exist. What about the infimum?

Question 3.38

Let $A \subseteq K$ be non-empty and bounded below. Does

 $\inf A$

always exist?

The answer to the above question is again **NO**. A set bounded below for which the infimum does nor exist is, for example,

$$A = (\sqrt{2}, 10] \cap \mathbb{Q}$$

The proof of this fact is, of course, very similar to the one of Theorem 3.37, and is therefore omitted.

In conclusion, infimum and supremum do not exist in general. The fields for which all bounded sets admit supremum or infimum are called **complete**.

Definition 3.39: Completeness

Let $(K, +, \cdot, \leq)$ be an ordered field. We say that *K* is **complete** if the following property holds:

• (AC) For every $A \subseteq K$ non-empty and bounded above

 $\sup A \in K$.

In particular, Theorem 3.37 can be re-stated as follows.

Theorem 3.40

 \mathbbm{Q} is not complete. In particular, there exists a set $A\subseteq \mathbbm{Q}$ such that

- A is non-empty,
- A is bounded above,
- $\sup A$ does not exist in \mathbb{Q} .

One of such sets is, for example,

```
A = \{q \in \mathbb{Q} \ : \ q \geq 0 \, , \ q^2 < 2 \} \, .
```

Notation 3.41

- 1. Property (AC) is called Axiom of Completeness
- 2. If K is an ordered field in which (AC) holds, then K is called a **complete ordered field**

Notice that if the **Axiom of Completeness** holds, then also the infimum exists. This is shown in the following proposition.

Proposition 3.42

Let $(K, +, \cdot, \leq)$ be a complete ordered field. Suppose that $A \subseteq K$ is non-empty and bounded below. Then

 $\inf A \in K$.

Proof

Suppose that $A \subseteq K$ is non-empty and bounded below. Then

$$-A := \{-a : a \in A\}$$

is non-empty and bounded above. By completeness we have that sup(-A) exists in *K*. From Proposition 2.46 we deduce that inf *A* exists in *K*, with

 $\inf A = -\sup(-A).$

Example 3.43

We have seen that $(\mathbb{Q}, +, \cdot, \leq)$ is an ordered field. However \mathbb{Q} is **not complete**, as proven in Theorem 3.37.

3.6 Equivalence of Completeness and Cut Property

We can show that **Completeness** is equivalent to the **Cut Property**.

Theorem 3.44: Equivalence of Cut Property and Completeness

Let $(K, +, \cdot, \leq)$ be an ordered field. They are equivalent:

- 1. *K* has the **Cut Property**
- 2. K is Complete

Remark 3.45: Ideas for proving Theorem 3.44

The proof of Theorem 3.44 is rather long, but the ideas are simple: Step 1. Cut Property \implies Completeness. Suppose *K* has the Cut Property. To prove that *K* is Complete, we need to:

- 1. Consider an arbitrary set $A \subseteq K$ such that $A \neq \emptyset$ and A is bounded above.
- 2. Show that *A* has a supremum.

To achieve this, consider the set of upper bounds of A

$$B := \{ b \in K : b \ge a, \forall a \in A \},\$$

We can show that the pair

 (B^c, B)

is a Cut of *K*. As *K* has the Cut Property, then there exists $s \in K$ separator of (B^c, B) . We will show that such separator *s* is the supremum of *A*, i.e.,

 $s = \sup A$.

This proves completeness of *K*. See Figure 3.7 for a schematic picture of the above construction. *Step 2. Completeness* \implies *Cut Property.*

Conversely, suppose that *K* is Complete. To prove that *K* has the Cut Property, we need to:

1. Consider a cut (A, B) of K.

2. Show that (A, B) has a separator $s \in K$.

This implication is easier. Indeed, since A is non-empty and bounded above (by the elements of B), by Completeness there exists

$$s := \sup A \in K$$
.

We will show that *s* is a separator for the cut (A, B), and therefore *K* has the Cut Property. See Figure 3.8 for a schematic picture of the above construction.



Figure 3.7: Let *s* be the separator of the cut (B^c, B) , with *B* the set of upper bounds of *A*. Then $s = \sup A$.



Figure 3.8: Let (A, B) be a cut of K and let $s = \sup A$. Then s is the separator of the cut (A, B).

Keeping the above ideas in mind, let us proceed with the proof.

Proof: Proof of Theorem 3.44

Step 1. Cut Property \implies Completeness.

We need to prove that *K* is complete. To this end, consider $A \subseteq K$ non-empty and bounded above. Define the set of upper bounds of *A*:

 $B := \{ b \in K : b \ge a, \forall a \in A \}.$

Claim. The pair (B^c, B) is a cut of K. Proof of Claim. We have to prove two points:

• (B^c, B) forms a partition of K.

Indeed, we have $B \neq \emptyset$, since A is bounded above. Further, we have $B^c \neq \emptyset$, since A is non-empty. Thus

 $K = B^c \cup B, \quad B^c \cap B = \emptyset.$

Then (B^c, B) is a partition of *K*.

• We have

 $x \le y, \quad \forall x \in B^c, \forall y \in B.$ (3.20)

To show the above, let $x \in B^c$ and $y \in B$. By definition of *B* we have that elements of B^c are not upper bounds of *A*. Therefore *x* is not an upper bound. This means there exists $\tilde{a} \in A$ which is larger than *x*, that is,

 $x \leq \tilde{a}$.

Since $y \in B$, then y is an upper bound for A, so that

 $a \leq y$, $\forall a \in A$.

Therefore

 $x \leq \tilde{a} \leq y$,

concluding (3.20).

Thus (B^c, B) is a cut of *K* and the claim is proven. Since (B^c, B) is a cut of *K*, by the Cut Property there exists a separator $s \in K$ such that

$$x \le s \le y, \quad \forall x \in B^c, \forall y \in B.$$
 (3.21)

Claim. s is an upper bound for A.

Proof of Claim.

Suppose by contradiction that *s* is not an upper bound for *A*. Therefore by definition of upper bound, there exists $\tilde{a} \in A$ such that

 $s < \tilde{a}$.

Consider the mid-point between s and \tilde{a} , that is,

$$m := \frac{s + \tilde{a}}{2} \in K.$$

Since *m* is the mid-point between *s* and \tilde{a} , and $s < \tilde{a}$, it holds

$$s < m < \tilde{a}$$
.

Indeed, since $s < \tilde{a}$ then

$$s = \frac{2s}{2} < \frac{s+\tilde{a}}{2} < \frac{2\tilde{a}}{2} = \tilde{a}.$$

In particular the above tells us that *m* is not an upper bound for *A*, given that $\tilde{a} \in A$ and $m < \tilde{a}$. Therefore $m \in B^c$, by definition of B^c . Therefore(3.21) implies

 $m \leq s$,

which contradicts s < m. Hence s is an upper bound of A, concluding the proof of Claim. *Conclusion*. We have shown that s is an upper bound of A. Condition (3.21) tells us that

$$s \leq y, \forall y \in B$$
.

Recalling that B is the set of upper bounds of A, this means that s is the smallest upper bound of A, that is,

$$s = \sup A \in K$$
.

Step 2. Completeness \implies Cut Property.

Suppose *K* is complete. We need to show that *K* has the Cut Property. Therefore assume (A, B) is a cut of *K*, that is,

$$A \neq \emptyset, \quad B \neq \emptyset,$$

$$K = A \cup B, \quad A \cap B = \emptyset,$$

$$a \le b, \quad \forall a \in A, \; \forall b \in B.$$
(3.22)

Since $B \neq \emptyset$, from (3.22) it follows that *A* is bounded above: indeed, every element of *B* is an upper bound for *A*, thanks to (3.22). Since $A \neq \emptyset$, by the Axiom of Completeness we have

$$s = \sup A \in K$$
.

In particular, by definition of supremum, we have

$$a \leq s$$
, $\forall a \in A$.

Let now $b \in B$ be arbitrary. From (3.22) we have that

$$a \le b , \ \forall a \in A . \tag{3.23}$$

Therefore *b* is an upper bound of *A*. Since $s = \sup A$, we have that *s* is the smallest upper bound, and so

 $s \leq b$.

Given that $b \in B$ was arbitrary, it actually holds

$$s \le b, \forall b \in B.$$
 (3.24)

From (3.23) and (3.24) we therefore have

$$a \leq s \leq b$$
, $\forall a \in A$, $\forall b \in B$,

showing that s is a separator of (A, B). Thus K has the Cut Property.

3.7 Axioms of Real Numbers

We now have all the key elements to introduce the Real Numbers R. These ingredients are:

- Definition of ordered field,
- The Cut Property or Axiom of Completeness.

The definition of \mathbb{R} is given in an axiomatic way.

Definition 3.46: System of Real Numbers ℝ

A system of Real Numbers is a set $\mathbb R$ with two operations + and $\cdot,$ and a total order relation $\leq,$ such that

- $(\mathbb{R}, +, \cdot, \leq)$ is an ordered field
- \mathbbm{R} sastisfies the Axiom of Completeness

For reader's convenience we explicitly state the above mentioned properties.

1. There is an operation + of **addition** on \mathbb{R}

 $+ : \mathbb{R} \times \mathbb{R} \to \mathbb{R}, \quad (x, y) \mapsto x + y$

The addition satisifes: $\forall x, y, z \in \mathbb{R}$

• (A1) Commutativity and Associativity:

$$x + y = y + x$$

$$(x+y) + z = x + (y+z)$$

• (A2) Additive Identity: $\exists 0 \in \mathbb{R}$ s.t.

$$x + 0 = 0 + x = x$$

• (A₃) Additive Inverse: $\exists (-x) \in \mathbb{R}$ s.t.

$$x + (-x) = (-x) + x = 0$$

2. There is an operation \cdot of **multiplication** on $\mathbb R$

$$\cdot : \mathbb{R} \times \mathbb{R} \to \mathbb{R}, \quad (x, y) \mapsto x \cdot y = xy$$

The multiplication satisifes: $\forall x, y, z \in \mathbb{R}$

• (M1) Commutativity and Associativity:

$$x \cdot y = y \cdot x$$
$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

• (M2) Multiplicative Identity: $\exists 1 \in \mathbb{R}$ s.t.

$$x \cdot 1 = 1 \cdot x = x$$

• (M₃) **Multiplicative Inverse**: If $x \neq 0$, $\exists x^{-1} \in \mathbb{R}$ s.t.

$$x \cdot x^{-1} = x^{-1} \cdot x = 1$$

- 3. There is a relation \leq of **total order** on \mathbb{R} . The order satisfies: $\forall x, y, z \in \mathbb{R}$
 - (O1) Reflexivity:

 $x \le x$

• (O2) Antisymmetry:

 $x \le y$ and $y \le x \implies x = y$

• (O₃) Transitivity:

 $x \le y$ and $y \le z \implies x = z$

• (O₄) Total order:

 $x \le y$ or $y \le x$

- 4. The operations + and ·, and the total order \leq , are related by the following properties: $\forall x, y, z \in \mathbb{R}$
 - (AM) Distributive: Relates addition and multiplication via

$$x \cdot (y+z) = x \cdot y + x \cdot z$$

• (AO) Relates addition and order with the requirement:

 $x \le y \implies x+z \le y+z$

• (MO) Relates multiplication and order with the requirement:

 $x \ge 0, \ y \ge 0 \implies x \cdot y \ge 0$

5. Axiom of Completeness:

• (AC) For every $A \subseteq \mathbb{R}$ non-empty and bounded above, there exists

 $\sup A \in \mathbb{R}$

Remark 3.47

Since **Axiom of Completeness** and **Cut Property** are equivalent by Theorem 3.44, one can replace the Axiom of Completeness in Definition 3.46 Point 5 with:

- 5. Cut Property holds:
 - (CP) Every cut (A, B) of \mathbb{R} admits a **separator** $s \in \mathbb{R}$ s.t.

 $a \leq s \leq b$, $\forall a \in A, \forall b \in B$

Notation 3.48

For $x \in \mathbb{R}$, $x \neq 0$, the multiplicative inverse is also denoted by

$$x^{-1} := \frac{1}{x}$$
.

Remark 3.49

Recall that

• $(K, +, \cdot)$ is a **field** if they hold:

(A1)-(A3), (M1)-(M3), (AM)

• $(K, +, \cdot, \leq)$ is an **ordered field** if they hold

(A1)-(A3), (M1)-(M3), (O1)-(O4) ,(AM), (AO), (MO)

In particular we have that

 $(\mathbb{R},+,\cdot,\leq)$

is a **complete ordered field**: that is, an ordered field in which the **Cut Property** (CP) or **Axiom of Completeness** (AC) hold.

Question 3.50

There are two main questions here:

- 1. We have only postulated the existence of R. Does such complete ordered field actually exist?
- 2. Is \mathbb{R} the only complete ordered field?

The answer to both questions is YES:

- There are several equivalent methods for explicitly constructing the system R. One such method uses **Dedekind cuts**. The interested reader can refer to the Appendix in Chapter 1 of [3], or Chapter 8.6 in [1].
- 2. It can be shown that $(\mathbb{R}, +, \cdot, \leq)$ is the **only** complete ordered field. Uniqueness is intended in the following sense: if $(K, +, \cdot, \leq)$ is another complete ordered field, then *K* looks like \mathbb{R} . Mathematically this means that there exists an invertible map $\Psi : \mathbb{R} \to K$, called isomorphism of fields, which preserves the operations $+, \cdot$ and the order \leq .

In particular the following Theorem can be proven.

Theorem 3.51: Existence of the Real Numbers

There exists a complete ordered field $(\mathbb{R}, +, \cdot, \leq)$ called the System of Real Numbers. Moreover \mathbb{R} is unique, up to isomorphism of fields.

3.8 Special subsets of ℝ

In Definition 3.46, we introduced \mathbb{R} as a complete ordered field, doing so axiomatically and in a nonconstructive manner. But what about the sets \mathbb{N} , \mathbb{Z} , \mathbb{Q} now? Are they still well-defined? Specifically, can we say that

$$\mathbb{N}, \mathbb{Z}, \mathbb{Q} \subseteq \mathbb{R}?$$

The definitions we provided in Chapter 1 for \mathbb{N} , \mathbb{Z} , and \mathbb{Q} are not directly linked to the real number system \mathbb{R} we just introduced. To address this issue, we will need to define new sets

$$\mathbb{N}_{\mathbb{R}}, \mathbb{Z}_{\mathbb{R}}, \mathbb{Q}_{\mathbb{R}}$$

from scratch, relying solely on the axioms of \mathbb{R} . The subscript \mathbb{R} is used here to clearly distinguish these new sets from the original ones.

3.8.1 Natural numbers

Let us start with the definition of $\mathbb{N}_{\mathbb{R}}.$ We would like $\mathbb{N}_{\mathbb{R}}$ to be

$$\mathbb{N}_{\mathbb{R}} = \{\mathbf{1}, \mathbf{2}, \mathbf{3}, \ldots\}.$$

We are denoting the above numbers with bold symbols in order to distinguish them from the elements of \mathbb{R} . The key property that we would like $\mathbb{N}_{\mathbb{R}}$ to have is the following:

Every $\mathbf{n} \in \mathbb{N}_{\mathbb{R}}$ has a successor $(\mathbf{n} + \mathbf{1}) \in \mathbb{N}_{\mathbb{R}}$.

How do we ensure this property? We could start by defining

1 := 1,

with 1 the neutral element of the multiplication in \mathbb{R} , which exists by the field axiom (M₂) in Definition 3.46. We could then define **2** by setting

 $\mathbf{2} := 1 + 1$.

We need a formal definition to capture this idea.

Definition 3.52: Inductive set

Let $S \subseteq \mathbb{R}$. We say that *S* is an inductive set if they are satisfied:

- 1 ∈ *S*,
- If $x \in S$, then $(x + 1) \in S$.

Note that in the above definition we just used:

- The existence of the neutral element 1, given by axiom (M2).
- The operation of sum in R, which is again given as an axiom.

Example 3.53

Question. Prove the following:

- 1. \mathbb{R} is an inductive set.
- 2. The set $A = \{0, 1\}$ is not an inductive set.

Solution.

- 1. We have that $1 \in \mathbb{R}$ by axiom (M2). Moreover $(x + 1) \in \mathbb{R}$ for every $x \in \mathbb{R}$, by definition of sum +.
- 2. We have $1 \in A$, but $(1 + 1) \notin A$, since $1 + 1 \neq 0$.

Since \mathbb{R} is itself an inductive set, it is clear that the definition of inductive set is not sufficient to fully describe our intuitive idea of $\mathbb{N}_{\mathbb{R}}$. The right way to define $\mathbb{N}_{\mathbb{R}}$ is as follows:

 $\mathbb{N}_{\mathbb{R}}$ is the smallest inductive subset of \mathbb{R} .

To make the above definition precise, we first need a Proposition.

Proposition 3.54

Let ${\mathscr M}$ be a collection of inductive subsets of ${\mathbb R}.$ Then

$$S := \bigcap_{M \in \mathcal{M}} M$$

is an inductive subset of $\mathbb R.$

Proof

We have to show that the two properties of inductive sets hold for S:

1. We have $1 \in M$ for every $M \in \mathcal{M}$, since these are inductive sets. Thus

$$\mathbf{I} \in \bigcap_{M \in \mathcal{M}} M = S.$$

2. Suppose that $x \in S$. By definition of *S* this implies that $x \in M$ for all $M \in \mathcal{M}$. Since *M* is an inductive set, then $(x + 1) \in M$. Therefore $(x + 1) \in M$ for all $M \in \mathcal{M}$, showing that $(x + 1) \in S$.

Therefore S is an inductive set.

We are now ready to define the natural numbers $\mathbb{N}_{\mathbb{R}}$.

Definition 3.55: Set of Natural Numbers

Let ${\mathscr M}$ be the collection of **all** inductive subsets of ${\mathbb R}.$ We define the set of natural numbers in ${\mathbb R}$ as

$$\mathbb{N}_{\mathbb{R}} := \bigcap_{M \in \mathscr{M}} M.$$

Therefore $\mathbb{N}_{\mathbb{R}}$ is the intersection of all the inductive subsets of \mathbb{R} . From this definition it follows that $\mathbb{N}_{\mathbb{R}}$ is the smallest inductive subset of \mathbb{R} , as shown in the following Proposition.

Proposition 3.56: $\mathbb{N}_{\mathbb{R}}$ is the smallest inductive subset of \mathbb{R}

Let $C\subseteq \mathbb{R}$ be an inductive subset. Then

 $\mathbb{N}_{\mathbb{R}} \subseteq C$.

In other words, $\mathbb{N}_{\mathbb{R}}$ is the smallest inductive set in $\mathbb{R}.$

Proof

Let \mathcal{M} be the collection of **all** inductive subsets of \mathbb{R} . By definition

$$\mathbb{N}_{\mathbb{R}} = \bigcap_{M \in \mathscr{M}} M.$$

Let $x \in \mathbb{N}_{\mathbb{R}}$. Then $x \in M$ for all $M \in \mathcal{M}$. Since $C \in \mathcal{M}$ then $x \in C$. This shows $\mathbb{N}_{\mathbb{R}} \subseteq C$.

The definition of $\mathbb{N}_{\mathbb{R}}$ guarantees that all numbers in $\mathbb{N}_{\mathbb{R}}$ are larger than 1.

Theorem 3.57 Let $x \in \mathbb{N}_{\mathbb{R}}$. Then

 $x \ge 1$.

Proof

Define the set

 $C := \{ x \in \mathbb{R} : x \ge 1 \}.$

We have that *C* is an inductive subset of \mathbb{R} .

By definition $1 \in C$. Suppose now that $x \in C$, so that $x \ge 1$. Since $1 \ge 0$ as a consequence of the field axioms, we deduce that

 $x + 1 \ge x + 0 = x \ge 1$,

showing that $x + 1 \ge 1$. Thus $(x + 1) \in C$.

By Proposition 3.56 we conclude that

 $\mathbb{N}_{\mathbb{R}} \subseteq C,$

showing that $x \ge 1$ for all $x \in \mathbb{N}_{\mathbb{R}}$.

Notation 3.58

We have just shown that all the numbers $x\in\mathbb{N}_{\mathbb{R}}$ satisfy

 $x \ge 1$.

Moreover, by the fact that $\mathbb{N}_{\mathbb{R}}$ is an inductive set, we know that

 $1+1\in\mathbb{N}_{\mathbb{R}}\,,$ since $1\in\mathbb{N}_{\mathbb{R}}.$ We denote $2\,:=\,1+1\,.$ Similarly, we will have that $2+1\in\mathbb{N}_{\mathbb{R}}\,,$ since $2\in\mathbb{N}_{\mathbb{R}}.$ We denote $3\,:=\,2+1\,.$ In this way we give a name to all the numbers in $\mathbb{N}_{\mathbb{R}}.$

3.8.2 Principle of induction

The Principle of Induction is a consequence of the definition of $\mathbb{N}_{\mathbb{R}}$, see Definition 3.55, and of the field axioms of \mathbb{R} in Definition 3.46.

Theorem 3.59: Principle of Induction

Let $\alpha(n)$ be a statement depending on $n \in \mathbb{N}_{\mathbb{R}}$. Assume that

- 1. $\alpha(1)$ is true.
- 2. If $\alpha(n)$ is true then also $\alpha(n + 1)$ is true.

Then $\alpha(n)$ is true for all $n \in \mathbb{N}_{\mathbb{R}}$.

Proof

Define the set

 $C := \{x \in \mathbb{N}_{\mathbb{R}} : \alpha(n) \text{ is true} \}.$

We have that *C* is an inductive subset of \mathbb{R} .

Indeed:

- $1 \in C$ since $\alpha(1)$ is true by assumption.
- If $n \in C$ then $\alpha(n)$ is true. By assumption $\alpha(n+1)$ is true. Therefore $(n+1) \in C$.

By Proposition 3.56 we conclude that

 $\mathbb{N}_{\mathbb{R}} \subseteq C$.

As by definition $C \subseteq \mathbb{N}_{\mathbb{R}}$, we have proven that

 $\mathbb{N}_{\mathbb{R}} = C$,

showing that $\alpha(n)$ is true for all $n \in \mathbb{N}_{\mathbb{R}}$.

As a consequence of the principle of induction, we can prove that $\mathbb{N}_{\mathbb{R}}$ is closed under the field operations of sum and multiplication.

Theorem 3.60

For all $n, m \in \mathbb{N}_{\mathbb{R}}$ we have:

1. $\mathbb{N}_{\mathbb{R}}$ is closed under addition, that is,

$$m+n\in\mathbb{N}_{\mathbb{R}}$$
.

2. $\mathbb{N}_{\mathbb{R}}$ is closed under multiplication, that is,

$$m \cdot n \in \mathbb{N}_{\mathbb{R}}$$
,

3. If m > n there exists $k \in \mathbb{N}_{\mathbb{R}}$ such that

$$m=n+k.$$

Proof

We only prove the first point, the other statements are left as an exercise. Fix $m \in \mathbb{N}_{\mathbb{R}}$. We prove that

$$m+n \in \mathbb{N}_{\mathbb{R}}, \quad \forall n \in \mathbb{N}_{\mathbb{R}},$$
 (3.25)

by using induction.

- Induction base: We have $m + 1 \in \mathbb{N}_{\mathbb{R}}$, since $m \in \mathbb{N}_{\mathbb{R}}$ and $\mathbb{N}_{\mathbb{R}}$ is an inductive set.
- Inductive hypothesis: Suppose $m + n \in \mathbb{N}_{\mathbb{R}}$. Since $\mathbb{N}_{\mathbb{R}}$ is an inductive set, we have $(m + n) + 1 \in \mathbb{N}_{\mathbb{R}}$. By according the sum according (A1) we get
 - n) + 1 $\in \mathbb{N}_{\mathbb{R}}$. By associativity of the sum, see axiom (A1), we get

$$m + (n + 1) = (m + n) + 1 \in \mathbb{N}_{\mathbb{R}}$$
,

which is the desired theis.

By the Induction Principle of Theorem 3.59 we conclude (3.25).

As a consequence of the above theorem, we see that the restriction of the operations of sum and multiplication to $\mathbb{N}_{\mathbb{R}}$ are still binary operations:

$$+ : \mathbb{N}_{\mathbb{R}} \times \mathbb{N}_{\mathbb{R}} \to \mathbb{N}_{\mathbb{R}}, \quad \cdot : \mathbb{N}_{\mathbb{R}} \times \mathbb{N}_{\mathbb{R}} \to \mathbb{N}_{\mathbb{R}}.$$

Equipped with the above operations, $\mathbb{N}_{\mathbb{R}}$ satisfies the following properties.

Theorem 3.61

 $(\mathbb{N}_{\mathbb{R}}, +, \cdot, \leq)$ satisfies the following axioms from Definition 3.46:

- (A1).
- (M1), (M2).
- (O1)-(O4).
- (AM), (AO), (MO).

The proof is trivial, as it follows immediately from the inclusion of $\mathbb{N}_{\mathbb{R}}$ in \mathbb{R} .

3.8.3 Integers

We have seen in Theorem 3.60 that $\mathbb{N}_{\mathbb{R}}$ is closed under addition. However $\mathbb{N}_{\mathbb{R}}$ is not closed under subtraction. We therefore define the set of **integers** $\mathbb{Z}_{\mathbb{R}}$ in a way that we can perform subtraction of any two natural numbers.

Definition 3.62: Set of Integers

The set of integers in $\mathbb R$ is defined by

$$\mathbb{Z}_{\mathbb{R}} := \{m - n : n, m \in \mathbb{N}_{\mathbb{R}}\}.$$

In the definition of $\mathbb{Z}_{\mathbb{R}}$ we denote by -n the inverse of n in \mathbb{R} , which exists by the field axiom (A₃) in Definition 3.46. The following characterization explains the relationship between $\mathbb{Z}_{\mathbb{R}}$ and $\mathbb{N}_{\mathbb{R}}$.

Theorem	3.63
---------	------

It holds

```
\mathbb{Z}_{\mathbb{R}} = \{-n : n \in \mathbb{N}_{\mathbb{R}}\} \cup \{0\} \cup \mathbb{N}_{\mathbb{R}}.
```

Proof

Define the set

$$M := \{-n : n \in \mathbb{N}_{\mathbb{R}}\} \cup \{0\} \cup \mathbb{N}_{\mathbb{R}}.$$

- $M \subseteq \mathbb{Z}_{\mathbb{R}}$: Suppose $m \in M$. We have 3 cases:
 - If $m \in \{-n : n \in \mathbb{N}_{\mathbb{R}}\}$ then there exists $n \in \mathbb{N}_{\mathbb{R}}$ such that m = -n. Thus

$$m=-n=1-(n+1)\in\mathbb{Z}_{\mathbb{R}},$$

since $1 \in \mathbb{N}_{\mathbb{R}}$ and $n + 1 \in \mathbb{N}_{\mathbb{R}}$ because $n \in \mathbb{N}_{\mathbb{R}}$.

- If m = 0 then

$$m=0=1-1\in\mathbb{Z}_{\mathbb{R}},$$

as $1 \in \mathbb{N}_{\mathbb{R}}$. - If $m \in \mathbb{N}_{\mathbb{R}}$ then

$$m=(m+1)-1\in\mathbb{Z}_{\mathbb{R}}\,,$$

since $1 \in \mathbb{N}_{\mathbb{R}}$ and $m + 1 \in \mathbb{N}_{\mathbb{R}}$, given that $m \in \mathbb{N}_{\mathbb{R}}$.

In all 3 cases we have shown that $m \in \mathbb{Z}_{\mathbb{R}}$, proving that $M \subseteq \mathbb{Z}_{\mathbb{R}}$.

• $\mathbb{Z}_{\mathbb{R}} \subseteq M$: Let $z \in \mathbb{Z}_{\mathbb{R}}$. Then z = m - n for some $n, m \in \mathbb{N}_{\mathbb{R}}$. We have 3 cases:

- If m = n then

$$z = m - n = m - m \stackrel{(A3)}{=} 0 \in M$$

− If m > n, by Theorem 3.60 there exists $k \in \mathbb{N}_{\mathbb{R}}$ such that m = k + n. Therefore

$$z = m - n = (k + n) - n$$

$$\stackrel{(A1)}{=} k + (n - n) \stackrel{(A3)}{=} k + 0$$

$$\stackrel{(A2)}{=} k \in M,$$

since $k \in \mathbb{N}_{\mathbb{R}}$.

− If *m* < *n*, by Theorem 3.60 there exists $k \in \mathbb{N}_{\mathbb{R}}$ such that n = k + m. Therefore

$$z=m-n=-k\in M\,,$$

since $k \in \mathbb{N}_{\mathbb{R}}$, where again we have used (implicitly) the field axioms (A1), (A2) and (A3).

Therefore $\mathbb{Z}_{\mathbb{R}} = M$.

Like we did with $\mathbb{N}_{\mathbb{R}}$, we can also show that $\mathbb{Z}_{\mathbb{R}}$ is closed under the operations of sum and multiplication.

Theorem 3.64

For all $n, m \in \mathbb{Z}_{\mathbb{R}}$ we have:

1. $\mathbb{Z}_{\mathbb{R}}$ is closed under addition, that is,

 $m+n\in\mathbb{Z}_{\mathbb{R}}$.

2. $\mathbb{Z}_{\mathbb{R}}$ is closed under multiplication, that is,

 $m \cdot n \in \mathbb{Z}_{\mathbb{R}}$,

The proof is left as an exercise. As a consequence of Theorem 3.64 we have that the restriction of

the operations of sum and multiplication to $\mathbb{Z}_{\mathbb{R}}$ are still binary operations:

 $+ \ : \ \mathbb{Z}_{\mathbb{R}} \times \mathbb{Z}_{\mathbb{R}} \to \mathbb{Z}_{\mathbb{R}} \,, \quad \cdot \ : \ \mathbb{Z}_{\mathbb{R}} \times \mathbb{Z}_{\mathbb{R}} \to \mathbb{Z}_{\mathbb{R}} \,.$

Equipped with the above operations, $\mathbb{Z}_{\mathbb{R}}$ satisfies the following properties.

Theorem 3.65

 $(\mathbb{Z}_{\mathbb{R}}, +, \cdot, \leq)$ satisfies the following axioms from Definition 3.46:

- (A1), (A2), (A3).
- (M1), (M2).
- (O1)-(O4).
- (AM), (AO), (MO).

Proof

The fact that

(A1), (A2), (M1), (M2), (O1)-(O4), (AM), (AO), (MO)

are satisfied descends immediately from the inclusion

 $\mathbb{Z}_{\mathbb{R}} \subseteq \mathbb{R}$.

We are left to prove (A₃). This is non-trivial because a priori the additive inverse -z of some $z \in \mathbb{Z}_{\mathbb{R}}$ belongs to \mathbb{R} . We need to check that $-z \in \mathbb{Z}_{\mathbb{R}}$. Indeed, since $z \in \mathbb{Z}_{\mathbb{R}}$, there exist $n, m \in \mathbb{N}_{\mathbb{R}}$ such that z = m - n. Define y := n - m. We have that $y \in \mathbb{Z}_{\mathbb{R}}$ and

$$z + y = (m - n) + (n - m) = (m - m) + (n - n) = 0.$$

Therefore *y* is the inverse of *z* and $y \in \mathbb{Z}_{\mathbb{R}}$, proving that the sum in $\mathbb{Z}_{\mathbb{R}}$ satisfies (A₃).

Remark 3.66

 $\mathbb{Z}_{\mathbb{R}}$ does not satisfy (M₃).

For example, let us show that $2 \in \mathbb{Z}_{\mathbb{R}}$ has no inverse in $\mathbb{Z}_{\mathbb{R}}$. Indeed, let $m \in \mathbb{Z}_{\mathbb{R}}$. By Theorem 3.63 we have 3 cases:

• $m \in \mathbb{N}_{\mathbb{R}}$: Since 2 > 1 we have

$$2 \cdot m > 1 \cdot m \ge 1$$

where in the last inequality we used that $m \ge 1$ for all $m \in \mathbb{N}_{\mathbb{R}}$, as shown in Theorem 3.57. The above shows that

$$2\cdot m>1\,,$$

and therefore *m* cannot be the inverse of 2.

- m = 0: Then $2 \cdot m = 0$, so that m cannot be the inverse of 2.
- m = -n with $n \in \mathbb{N}_{\mathbb{R}}$. Then

$$2 \cdot m = 2 \cdot (-n) < 0,$$

so that *m* cannot be the inverse of 2.

As we have exhausted all the possibilities, we conclude that 2 does not have a multiplicative inverse in $\mathbb{N}_{\mathbb{R}}.$

3.8.4 Rational numbers

In Theorem 3.65 and 3.66 we have seen that $\mathbb{Z}_{\mathbb{R}}$ satisfy all the field axiom, except for (M3). We therefore extend $\mathbb{Z}_{\mathbb{R}}$ in a way that the extension contains multiplicative inverses. The extension is the set of rational numbers $\mathbb{Q}_{\mathbb{R}}$.

Definition 3.67: Set of Rational Numbers

The set of rational numbers in $\mathbb R$ is

$$\mathbb{Q}_{\mathbb{R}} := \left\{ \frac{m}{n} : m \in \mathbb{Z}_{\mathbb{R}}, n \in \mathbb{N}_{\mathbb{R}} \right\}.$$

Notice that in the above definition we are just using the field axiom (M₃), with

$$\frac{m}{n} := m \cdot n^{-1} \, .$$

The inverse of *n* exists because we are assuming $n \in \mathbb{N}_{\mathbb{R}}$, and therefore *n* cannot be 0, as a consequence of Theorem 3.57.

The set $Q_{\mathbb{R}}$ is closed under addition and multiplication (exercise). Therefore they are well defined the operations:

$$+ : \mathbb{Q}_{\mathbb{R}} \times \mathbb{Q}_{\mathbb{R}} \to \mathbb{Q}_{\mathbb{R}}, \quad \cdot : \mathbb{Q}_{\mathbb{R}} \times \mathbb{Q}_{\mathbb{R}} \to \mathbb{Q}_{\mathbb{R}}.$$

Theorem 3.68

 $(\mathbb{Q}_{\mathbb{R}}, +, \cdot, \leq)$ is an ordered field.

Proof

All the field properties, except for (M3), follow from the inclusion

 $\mathbb{Q}_{\mathbb{R}} \subseteq \mathbb{R}$

and from the field properties of \mathbb{R} . To check (M₃), let $q \in \mathbb{Q}_{\mathbb{R}}$ with $q \neq 0$. Therefore q = m/n for $m \in \mathbb{Z}_{\mathbb{R}}$, $n \in \mathbb{N}_{\mathbb{R}}$. As $q \neq 0$ and $n \neq 0$, see Theorem 3.57, we deduce that $m \neq 0$. We have two cases:

• m > 0: In this case $m \in \mathbb{N}_{\mathbb{R}}$ by Theorem 3.63. Therefore

$$p = \frac{n}{m} \in \mathbb{Q}_{\mathbb{R}}$$

by definition, since $n, m \in \mathbb{N}_{\mathbb{R}}$. By commutativity we have

$$q \cdot p = \frac{m}{n} \cdot \frac{n}{m} = 1$$
.

• m < 0: Then m = -x with $x \in \mathbb{N}_{\mathbb{R}}$ by Theorem 3.63. Therefore

$$p = \frac{-n}{x} \in \mathbb{Q}_{\mathbb{R}}$$

by definition, since $-n \in \mathbb{Z}_{\mathbb{R}}$ and $x \in \mathbb{N}_{\mathbb{R}}$. By commutativity we have

$$q \cdot p = \frac{m}{n} \cdot \frac{-n}{x} = \frac{m}{n} \cdot \frac{-n}{-m} = 1.$$

Therefore *q* always admits a multiplicative inverse q^{-1} belonging to $Q_{\mathbb{R}}$, proving (M₃).

The set $\mathbb{Q}_{\mathbb{R}}$ does not have the Cut Property or the Axiom of Completeness.

Theorem 3.69

 $\mathbb{Q}_{\mathbb{R}}$ is not complete.

The proof of the above Theorem replicates exactly the proof of Theorem 3.37: This is because the proof of Theorem 3.37 only makes use of field axioms, and thus it applies to $\mathbb{Q}_{\mathbb{R}}$.

Notation 3.70

From now on we denote

$$\mathbb{N} \, := \mathbb{N}_{\mathbb{R}} \,, \quad \mathbb{Z} \, := \mathbb{Z}_{\mathbb{R}} \,, \quad \mathbb{Q} \, := \mathbb{Q}_{\mathbb{R}} \,,$$

dropping the subscript $\mathbb R.$

4 Properties of \mathbb{R}

Now that we established the axiomatic definition of the Real Numbers \mathbb{R} as a **complete ordered field**, let us investigate some of the properties of \mathbb{R} . These will be consequence of the axioms of the real numbers, particularly of the **Axiom of Completeness**.

4.1 Archimedean Property

The Archimedean property is one of the most useful properties of \mathbb{R} , and it essentially states that the set of natural numbers \mathbb{N} is not bounded above in \mathbb{R} .

More precisley, the Archimedean Property says two things:

1. For any $x \in \mathbb{R}$ we can always find a natural number $n \in \mathbb{N}$ such that

$$n > x$$
.

2. For any $x \in \mathbb{R}$ with x > 0, we can always find a natural number $m \in \mathbb{N}$ such that

$$0 < \frac{1}{m} < x \, .$$

The situation is depicted in Figure 4.2.

$$\begin{array}{c|c} \bullet & \bullet & \bullet \\ \hline 0 & \underline{1} & x & n \end{array} \rightarrow \mathbb{R}$$

Figure 4.1: For any x > 0 we can find $n, m \in \mathbb{N}$ such that 1/m < x < n.

Remark 4.1

The Archimedean property might sound trivial. However there are examples of ordered fields *K* that satisfy:

- 1. $\mathbb{N} \subseteq K$.
- 2. *K* does not have the Archimedean property.
- 3. In particular, \mathbb{N} is bounded above in *K*.

Of course such fields *K* cannot be complete.

If *K* is complete, then *K* is essentially \mathbb{R} , and we are going to prove the Archimedean

Property holds in $\mathbb R.$

Let us proceed with the precise statement of the Archimedean property in R.

Theorem 4.2: Archimedean Property

Let $x \in \mathbb{R}$ be given. Then:

1. There exists $n \in \mathbb{N}$ such that

n > x.

2. Suppose in addition that x > 0. There exists $n \in \mathbb{N}$ such that

$$\frac{1}{n} < x \, .$$

Proof

Part 1. Let $x \in \mathbb{R}$. Suppose by contradiction that there is no $n \in \mathbb{N}$ such that

n > x.

This means that

$$n \le x \quad \forall n \in \mathbb{N} \,. \tag{4.1}$$

The above is saying that the set $\mathbb N$ is bounded above. Since $\mathbb N$ is not empty, by the Axiom of Completeness there exists

 $\alpha = \sup \mathbb{N}$.

Claim: $(\alpha - 1)$ is not an upper bound for \mathbb{N} . Proof of Claim. Indeed, we have

 $(\alpha - 1) < \alpha \,. \tag{4.2}$

Therefore $\alpha - 1$ cannot be an upper bound for \mathbb{N} . Indeed, if by contradiction $\alpha - 1$ was an upper bound for \mathbb{N} , then we would have

$$\alpha \leq (\alpha - 1),$$

since α is the smallest upper bound for \mathbb{N} . This contradicts (4.2). Therefore $\alpha - 1$ is not an upper bound for \mathbb{N} .

Conclusion. Since $\alpha - 1$ is not an upper bound for \mathbb{N} , there exists $n_0 \in \mathbb{N}$ such that

$$\alpha - 1 < n_0 \, .$$

The above implies

 $\alpha < n_0 + 1.$

Since

$$(n_0+1)\in\mathbb{N}$$
,

we have obtained a contradiction, given that α was the supremum of N. Thus (4.1) is false, meaning that there exists $n \in \mathbb{N}$ such that

n > x.

Part 2. Suppose $x \in \mathbb{R}$ with x > 0. We can define

$$y := \frac{1}{x}.$$

By Part 1 there exists $n \in \mathbb{N}$ such that

 $n > y = \frac{1}{r}$.

Using that x > 0, we can rearrange the above inequlaity to obtain

$$\frac{1}{n} < x \,,$$

which is the desired thesis.

There is another formulation of the Archimedean Property which, depending on the situation, might be more useful. This formulation says the following: If $x, y \in \mathbb{R}$ are such that

 $0 < x < \gamma$,

then there exists $n \in \mathbb{N}$ such that

nx > y.

In other words, if one does n steps of size x in the positive numbers direction, then the resulting number *nx* will be larger than *y*. The situation is depicted in Figure 4.2.



Figure 4.2: For 0 < x < y there exists $n \in \mathbb{N}$ such that hat nx > y. In the picture n = 3.

Theorem 4.3: Archimedean Property (Alternative formulation)

Let $x, y \in \mathbb{R}$, with 0 < x < y. There exists $n \in \mathbb{N}$ such that

nx > y.

Proof

Suppose by contradiction there is no natural $n \in \mathbb{N}$ such that

nx > y.

$$nx \le y, \quad \forall n \in \mathbb{N}. \tag{4.3}$$

Define the set

 $A := \{nx : n \in \mathbb{N}\}.$

Condition (4.3) is saying that A is bounded above by y. Morever A is trivially non-empty. By the Axiom of Completeness there exists

$$\alpha = \sup A$$
.

Since α is the supremum of *A*, by definition of supremum and of the set *A*, we have

$$nx \le \alpha, \quad \forall n \in \mathbb{N}. \tag{4.4}$$

As (4.4) holds for every $n \in \mathbb{N}$, then it also holds for (n + 1), meaning that

 $(n+1)x \leq \alpha$.

The above implies

 $nx \leq \alpha - x$.

As *n* was arbitrary, we conclude that

 $nx \leq \alpha - x, \quad \forall n \in \mathbb{N}.$

The above is saying that $(\alpha - x)$ is an upper bound for *A*. Since α is the supremum of *A*, in particular α is the smallest upper bound. Thus it must hold

 $\alpha \leq \alpha - x$.

The above is equivalent to

 $x \leq 0$,

which contradicts our assumption of x > 0. Therefore (4.3) is false, and there exists $n \in \mathbb{N}$ such that

nx > y,

concluding the proof.

4.2 Nested Interval Property

Another consequence of the axiom of completeness is the **Nested Interval Property**. This is yet another way of saying the same thing: \mathbb{R} does not have **gaps**.

Let us look at a construction. Suppose given some closed intervals

$$I_n := [a_n, b_n] = \{x \in \mathbb{R} : a_n \le x \le b_n\},\$$

where the end points are ordered in the following way:

$$a_1 \leq a_2 \leq \ldots \leq a_n \leq \cdots \leq b_n \leq \ldots b_n \leq b_1$$

as shown in Figure 4.3.



Figure 4.3: Nested intervals $I_n = [a_n, b_n]$.

The intervals I_n are **nested**, meaning that

$$I_1 \supset I_2 \supset I_3 \supset \dots \supset I_n \supset \dots$$

For finite intersections we clearly have

$$\bigcap_{n=1}^k I_n = I_k \,,$$

that is, intersecting the first k intervals yields I_k , the smallest interval in the sequence.

Question 4.4

Consider the infinite intersection

$$\bigcap_{n=1}^{\infty} I_n := \left\{ x \in \mathbb{R} : x \in I_n , \forall n \in \mathbb{N} \right\}.$$

What can we say about it? Is it empty? Is it not empty?

The answer is that the infinite intersection is not empty, because \mathbb{R} was constructed in a way that it does not have gaps.

Theorem 4.5: Nested Interval Property

For each $n \in \mathbb{N}$ assume given a closed interval

$$I_n := [a_n, b_n] = \{x \in \mathbb{R} : a_n \le x \le b_n\}.$$

Suppose that the intervals are nested, that is,

$$I_n \supset I_{n+1}, \quad \forall n \in \mathbb{N}$$

Then

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset.$$
(4.5)

Proof

By definition we have

$$\bigcap_{n=1}^{\infty} I_n \, := \left\{ x \in \mathbb{R} \, : \, x \in I_n \, , \forall \, n \in \mathbb{N} \right\}.$$

We want to prove (4.5). This means we need to find a real number x such that

$$x \in I_n, \quad \forall n \in \mathbb{N}.$$
 (4.6)

Idea of the Proof: Condition (4.6) is saying that it should hold

 $a_n \leq x \leq b_n$, $\forall n \in \mathbb{N}$.

Since a_n is increasing and b_n is decreasing, the point x is being squeezed between these two sequences. This suggests x should be the largest (i.e. the supremum) of all a_ns and smallest (i.e. the infimum) of all the b_ns .

Define the set

 $A := \{a_n : n \in \mathbb{N}\}.$

The set A is non-empty and is bounded above by any of the b_n . Therefore there exists

```
x = \sup A.
```

By definition of supremum and definition of the set *A*, we have

$$a_n \leq x$$
, $\forall n \in \mathbb{N}$

On the other hand, consider an arbitrary number b_n . By construction we have

$$a_i \leq b_n$$
, $\forall i \in \mathbb{N}$.

Therefore b_n is an upper bound for A. Since the supremum is the smallest upper bound, we conclude that

 $x \leq b_n$.

The index n was chosen arbitrarily, and therefore

$$x \leq b_n$$
, $\forall n \in \mathbb{N}$.

In total we have

$$a_n \leq x \leq b_n$$
, $\forall n \in \mathbb{N}$,

showing that x satisfies (4.6). Therefore (4.5) holds and the proof is concluded.

Important

The assumption that I_n is **closed** is crucial in Theorem 2.59. Without such assumption the thesis of Theorem 2.59 does not hold in general, as seen in Example 4.6 below.

Example 4.6

Question. Consider the open intervals

$$I_n := \left(0, \frac{1}{n}\right).$$

These are clearly nested

$$I_n \supset I_{n+1}$$
, $\forall n \in \mathbb{N}$

Prove that

$$\bigcap_{n=1}^{\infty} I_n = \emptyset \,. \tag{4.7}$$

Solution. Suppose by contradiction that the intersection is non-empty. Then there exists $x \in \mathbb{N}$ such that

$$x \in I_n$$
, $\forall n \in \mathbb{N}$.

By definition of I_n the above reads

$$0 < x < \frac{1}{n}, \quad \forall n \in \mathbb{N}.$$
(4.8)

Since x > 0, by the Archimedean Property in Theorem 2.57 Point 2, there exists $n_0 \in \mathbb{N}$ such that

$$0 < \frac{1}{n_0} < x \, .$$

The above contradicts (4.8). Therefore (4.7) holds.

4.3 Revisiting Sup and inf

We now investigate some of the properties of supremum and infimum in \mathbb{R} . The first property is an alternative characterization of the supremum, which we will often use. A sketch of such characterization is in Figure 4.4 below.

Proposition 4.7: Characterization of Supremum

Let $A \subseteq \mathbb{R}$ be a non-empty set. Suppose that $s \in \mathbb{R}$ is an upper bound for A. They are equivalent:

- 1. $s = \sup A$
- 2. For every $\varepsilon > 0$ there exists $x \in A$ such that



Figure 4.4: Let $s = \sup A$. Then for every $\varepsilon > 0$ there exist $x \in A$ such that $s - \varepsilon < x$.

Proof: Proof of Proposition 2.61

Step 1. Assume that

 $s = \sup A$.

Let $\varepsilon > 0$ be arbitrary. We clearly have that

$$s - \varepsilon < s \,. \tag{4.9}$$

Therefore $(s - \varepsilon)$ cannot be an upper bound of *A*. Indeed, if by contradiction $(s - \varepsilon)$ was an upper bound, then we would have

$$s \leq (s - \varepsilon)$$
,

since *s* is the smallest upper bound. The above contradicts (4.9), and therefore $(s - \varepsilon)$ is not an upper bound for *A*. Hence there exists some $x \in A$ such that

$$s-\varepsilon < x$$
,

concluding.

Step 2. Assume that Point 2 in the statement of Proposition 2.61 holds. By assumption we have that *s* is an upper bound for *A*. Suppose by contradiction that

$$s \neq \sup A$$
.

This is equivalent to the statement

s is not the smallest upper bound of A. (4.10)

Hence there exists an upper bound b of A such that

b < s.

Let

 $\varepsilon := s - b$.

By assumption there exists $x \in A$ such that

 $s - \varepsilon < x$.

Substituting the definition of ε we get

 $s - s + b < x \implies b < x$.

Since *b* is an upper bound for *A* and $x \in A$, the above is a contradiction. Therefore (4.10) is false, and *s* is the smallest upper bound of *A*. Thus $s = \sup A$.

The analogue of Proposition 2.61 is as follows. The proof is left as an exercise.

Proposition 4.8: Characterization of Infimum

Let $A \subseteq \mathbb{R}$ be a non-empty set. Suppose that $i \in \mathbb{R}$ is a lower bound for A. They are equivalent:

i = inf A
 For every ε ∈ ℝ, with ε > 0, there exists x ∈ A such that

 $x < i + \varepsilon$.

A sketch of the characterization in Proposition 2.62 can be found in Figure 4.5 below.





With the above characterizations of supremum and infimum, it will be easier, in some cases, to compute infimum and supremum. We now characterize infimum and supremum of an open interval of \mathbb{R} .

Proposition 4.9	
Let $a, b \in \mathbb{R}$ with $a < b$. Let	
	$A := (a, b) = \{ x \in \mathbb{R} : a < x < b \}.$

1. We have that

$$\inf A = a, \quad \sup A = b.$$

2. $\min A$ and $\max A$ do not exist.

Proof

Part 1. We will only prove that

since the proof of

 $\sup A = b$

 $\inf A = a$,

is similar. By definition of *A*, we have that

a < x, $\forall x \in A$.

The above says that *a* is a lower bound for *A*. *Claim. a is the largest lower bound of A*. *Proof of Claim.* Let *L* be a lower bound for *A*, that is,

$$L \le x \,, \quad \forall \, x \in A \,.$$

We have to prove that

Suppose by contradiction that (4.11) does not hold, namely that

a < L .

We want to prove that L < b. To conclude this, first observe that the midpoint between *a* and *b* belongs to *A*:

$$\frac{a+b}{2} \in A.$$

Indeed, using that a < b, we have

$$\frac{a+b}{2} > \frac{2a}{2} = a,$$

and

$$\frac{a+b}{2} < \frac{2b}{b} = b\,,$$

showing that the midpoint between a and b belongs to A.

 $L \le a \,. \tag{4.11}$

Since *L* is a lower bound for *A*, and $\frac{a+b}{2} \in A$, we have

$$L \le \frac{a+b}{2} < b \,, \tag{4.12}$$

where in the second inequality we used that a < b. In particular, we have proven that a < L < b, which means that $L \in A$.

We now want to find an element of A which is less than L. A good candidate is the midpoint between a and L, that is,

$$M := \frac{a+L}{2}$$

Since a < L < b, we have that

 $M \in A$.

Indeed, using that a < L, we have

$$a = \frac{2a}{2} < \frac{a+L}{2} = M \le \frac{a+b}{2}$$
.

Using that L < b, we have

$$M = \frac{a+L}{2} \le \frac{a+b}{2} < b.$$

Thus a < M < b, showing that $M \in A$.

Moreover

M < L .

This is because
$$a < L$$
, and therefore

$$M = \frac{a+L}{2} < \frac{2L}{2} = L.$$

This is a contradiction, since *L* is a lower bound for *A* and $M \in A$. Therefore (4.11) holds, showing that *a* is the largest lower bound of *A*. We conclude that $a = \inf A$.

Part 2. Suppose by contradiction that min *A* exists. We have shown that if the minimum of a set exists, then it must be

 $\min A = \inf A.$

Since

 $\inf A = a$,

by Proposition 4.9, we would obtain that

$$\min A = a.$$

As min $A \in A$ by definition, we infer $a \in A$. This is contradiction. Then min A does not exist. The proof that max A does not exist is similar, and is left as an exercise.

We can also consider intervals for which one or both of the sides are closed.

Corollary 4.10	
Let $a, b \in \mathbb{R}$ with $a < b$. Let	
	$A := [a, b) = \{ x \in \mathbb{R} : a \le x < b \}.$
Then	$\min A = \inf A = a, \sup A = b,$
max A does not exist.	

The proof is very simple, and is left as an exercise. Let us now compute supremum and infimum of a set which is not an interval.

Proposition 4.11

Define the set

$$A := \left\{ \frac{1}{n} : n \in \mathbb{N} \right\}.$$

Then

 $\inf A = 0$, $\sup A = \max A = 1$.

Proof

Part 1. We have

$$\frac{1}{n} \leq 1$$
, $\forall n \in \mathbb{N}$.

Therefore 1 is an upper bound for A. Since $1 \in A$, by definition of maximum we conclude that

$$\max A = 1$$
.

Since the maximum exists, we conclude that also the supremum exists, and

$$\sup A = \max A = 1.$$

Part 2. We have

$$\frac{1}{n} > 0, \quad \forall n \in \mathbb{N},$$

showing that 0 is a lower bound for *A*. Suppose by contradiction that 0 is not the infimum. Therefore 0 is not the largest lower bound. Then there exists $\varepsilon \in \mathbb{R}$ such that:

• ε is a lower bound for *A*, that is,

$$\varepsilon \leq \frac{1}{n}, \quad \forall n \in \mathbb{N},$$
(4.13)

• ε is larger than 0:

 $0 < \varepsilon$.

As $\varepsilon > 0$, by the Archimedean Property there exists $n_0 \in \mathbb{N}$ such that

$$0 < \frac{1}{n_0} < \varepsilon \, .$$

This contradicts (4.13). Thus 0 is the largest lower bound of A, that is, $0 = \inf A$. *Part 3.* We have that min A does not exist. Indeed suppose by contradiction that min A exists. Then

 $\min A = \inf A.$

As $\inf A = 0$ by Part 2, we conclude $\min A = 0$. As $\min A \in A$, we obtain $0 \in A$, which is a contradiction.

4.4 Existence of *k*-th Roots

We have started our discussion in Chapter 1 by proving that

$$\sqrt{2} \notin \mathbb{Q}$$
. (4.14)

We have shown that (4.14) implies that the set

 $A \, := \{ q \in \mathbb{Q} \, : \, q^2 < 2 \}$

does not have a supremum in \mathbb{Q} . We then introduced the Real Numbers \mathbb{R} so that each non-empty and bounded above set would have a supremum. As the set *A* is non-empty and bounded above, there exists $\alpha \in \mathbb{R}$ such that

 $\alpha = \sup A.$

We are going to prove that

 $\alpha^2 = 2$,

which means that α is the square root of 2.

More in general, we can prove existence of *k*-th roots: Let $x \in \mathbb{R}$ with $x \ge 0$ and $k \in \mathbb{N}$. Define the set

$$A := \{t \in \mathbb{R} : t^k < x\}$$

We will see that A is non-empty and bounded above. By the axiom of completeness there exists $\alpha \in \mathbb{R}$ such that

$$\alpha = \sup A$$

In the following Theorem we will prove that

$$\alpha^k = x$$
,

meaning that α is the *k*-th root of *x*.
Theorem 4.12: Existence of *k*-th roots

Let $x \in \mathbb{R}$ with $x \ge 0$ and $k \in \mathbb{N}$. Define the set

$$A = \{ t \in \mathbb{R} : t^k < x \}.$$

There exists $\alpha \in \mathbb{R}$ such that

 $\alpha = \sup A$.

Moreover it holds that

 $\alpha^k = x$.

The proof of Theorem 4.12 rests on similar ideas to the ones used to prove that Q does not have the cut property.

Proof: Proof of Theorem 4.12

Part 1: Uniqueness. Suppose $\alpha_1, \alpha_2 \in \mathbb{R}$ are such that

$$\alpha_1^k = \alpha_2^k = x.$$

 $\alpha_1^k \neq \alpha_2^k$,

If $\alpha_1 \neq \alpha_2$, then

obtaining a contradiction. Therefore $\alpha_1 = \alpha_2$. *Part 2: Existence.* Let $x \in \mathbb{R}$ with $x \ge 0$. If x = 0 there is nothing to prove, as

$$0^k = 0$$
,

so that $\alpha = 0$. Therefore we can assume x > 0. Define the subset of \mathbb{R}

$$A := \{t \in \mathbb{R} : t^k < x\}.$$

Clearly A is non-empty and bounded above.

An upper bound is given, for example, by b := x+1. Indeed, since we are assuming $x \ge 0$, then $x + 1 \ge 1$. In particular we have

$$(x+1)^k \ge x+1.$$

Let $t \in A$. Then

$$t^k < x < x + 1 < (x + 1)^k,$$

showing that t < x + 1.

By the Axiom of Completeness of \mathbb{R} , there exists $\alpha \in \mathbb{R}$ such that

 $\alpha = \sup A$.

We claim that

$$\alpha^k = x \,. \tag{4.15}$$

Suppose by contradiction that (4.15) does not hold. We will need the formula: For all $a, b \in \mathbb{R}$ it holds

$$a^{k} - b^{k} = (a - b)(a^{k-1} + a^{k-2}b + a^{k-3}b^{k} + \dots + ab^{k-2} + b^{k-1}).$$
(4.16)

Formula (4.16) can be easily proven by induction on *k*. Since we are assuming that (4.15) does not hold, we have two cases:

• $\alpha^k < x$: We know that α is the supremum of *A*. We would like to violate this, by finding a number *L* which is larger than α , but still belongs to *A*. This means *L* has to satisfy

$$\alpha < L, \quad L^k < x.$$

We look for L of the form

$$L_n := \alpha + \frac{1}{n}$$

for $n \in \mathbb{N}$ to be chosen later. Clearly

$$\alpha < L_n \,, \tag{4.17}$$

for all $n \in \mathbb{N}$. We now search for $n_0 \in \mathbb{N}$ such that

$$L_{n_0}^k < x$$

Using formula (4.16) with $a = \alpha$ and $b = L_{n_0}$ we obtain

$$L_{n_0}^k - \alpha^k = \frac{1}{n_0} \left(L_{n_0}^{k-1} + L_{n_0}^{k-2} \alpha + \dots + L_{n_0} \alpha^{k-2} + \alpha^{k-1} \right) .$$
(4.18)

Now notice that (4.17) implies

$$\alpha^j < L_{n_0}^j$$

for all $j \in \mathbb{N}$. Using this estimate on all the terms α^j appearing in the RHS of (4.18) we obtain

$$\begin{split} L_{n_0}^k - \alpha^k &= \frac{1}{n_0} \left(L_{n_0}^{k-1} + L_{n_0}^{k-2} \alpha + \dots + L_{n_0} \alpha^{k-2} + \alpha^{k-1} \right) \\ &< \frac{1}{n_0} \left(L_{n_0}^{k-1} + L_{n_0}^{k-2} L_{n_0} + \dots + L_{n_0} L_{n_0}^{k-2} + L_{n_0}^{k-1} \right) \\ &= \frac{k}{n_0} L_{n_0}^{k-1} \end{split}$$

Rearranging the above we get

$$L_{n_0}^k < \frac{k}{n_0} L_{n_0}^{k-1} + \alpha^k \,. \tag{4.19}$$

Now note that

$$L_{n_0}=\alpha+\frac{1}{n_0}<\alpha+1.$$

Therefore

$$L_{n_0}^{k-1} < (\alpha + 1)^{k-1}$$
,

and from (4.19) we obtain

$$L_{n_0}^k < \frac{k}{n_0} (\alpha + 1)^{k-1} + \alpha^k$$
.

We wanted to find $n_0 \in \mathbb{N}$ so that $L_{n_0}^k < x$. Therefore we impose

$$\frac{k}{n_0} \left(\alpha + 1 \right)^{k-1} + \alpha^k < x \,,$$

and find that the above is satisfied for

$$n_0 > \frac{k(\alpha+1)^{k-1}}{x-\alpha^k} \,. \tag{4.20}$$

Notice that the RHS in (4.20) is a positive real number, since $\alpha^k < x$ by assumption. Therefore, by the Archimedean Property of Theorem 2.57 Point 1, there exists $n_0 \in \mathbb{N}$ satisfying (4.20).

We have therefore shown the existence of $n_0 \in \mathbb{N}$ such that

$$\alpha < L_{n_0}, \quad L_{n_0}^k < x.$$

The above says that $L_{n_0} \in A$ and that

$$L_{n_0} > \alpha = \sup A$$
,

which is a contradiction, as sup A is an upper bound for A.

• $\alpha^k > x$: We know that α is the supremum of *A*. We would like to find a contradiction, by finding an upper bound *L* for *A* which is smaller than α . This means *L* has to satisfy

$$L < \alpha$$
, $L^k > x$.

Such *L* is an upper bound for *A*: If $t \in A$ then

$$t^k < x < L^k \quad \Longrightarrow \quad t < L \,.$$

We therefore look for L of the form

$$L_n := \alpha - \frac{1}{n}$$

for $n \in \mathbb{N}$ to be chosen later. In this way

$$L_n < \alpha \,, \tag{4.21}$$

for all $n \in \mathbb{N}$. We now search for $n_0 \in \mathbb{N}$ such that

$$L_{n_0}^k > x \, .$$

Using formula (4.16) with $a = L_{n_0}$ and $b = \alpha$ we obtain

$$\alpha^{k} - L_{n_{0}}^{k} = \frac{1}{n_{0}} \left(\alpha^{k-1} + \alpha^{k-2} L_{n_{0}} + \dots + \alpha L_{n_{0}}^{k-2} + L_{n_{0}}^{k-1} \right) .$$
(4.22)

Now notice that (4.21) implies

 $L_{n_0}^j < \alpha^j$

for all $j \in \mathbb{N}$. Using this estimate on all the terms $L_{n_0}^j$ appearing in the RHS of (4.22) we obtain

$$\begin{aligned} \alpha^{k} - L_{n_{0}}^{k} &= \frac{1}{n_{0}} \left(\alpha^{k-1} + \alpha^{k-2} L_{n_{0}} + \dots + \alpha L_{n_{0}}^{k-2} + L_{n_{0}}^{k-1} \right) \\ &< \frac{1}{n_{0}} \left(\alpha^{k-1} + \alpha^{k-2} \alpha + \dots + \alpha \alpha^{k-2} + \alpha^{k-1} \right) \\ &= \frac{k}{n_{0}} \alpha^{k-1} \end{aligned}$$

Rearranging the above we get

$$L_{n_0}^k > \alpha^k - \frac{k}{n_0} \, \alpha^{k-1}$$

We wanted to find $n_0 \in \mathbb{N}$ so that $L_{n_0}^k > x$. Therefore we impose

$$\alpha^k - \frac{k}{n_0} \, \alpha^{k-1} > x \, ,$$

and find that the above is satisfied for

$$n_0 > \frac{k\alpha^{k-1}}{\alpha^k - x} \,. \tag{4.23}$$

Notice that the RHS in (4.23) is a positive real number, since $\alpha^k > x$ by assumption. Therefore, by the Archimedean Property of Theorem 2.57 Point 1, there exists $n_0 \in \mathbb{N}$ satisfying (4.23).

We have therefore shown the existence of $n_0 \in \mathbb{N}$ such that

$$L_{n_0} < \alpha \,, \quad L_{n_0}^k > x \,.$$

Condition $L_{n_0}^k > x$ says that L_{n_0} is an upper bound for *A*. At the same time it holds

$$L_{n_0} < \alpha = \sup A$$
,

which is a contradiction, as $\sup A$ is the smallest upper bound for A.

Therefore, both cases $\alpha^k > x$ and $\alpha^k < x$ lead to a contradiction. Hence $\alpha^k = x$, concluding.

Definition 4.13: *k*-th root of a number

Let $x \in \mathbb{R}$ with $x \ge 0$ and $k \in \mathbb{N}$. The real number α such that

 $\alpha^k = x$

is called the *k*-th root of *x*, and is denoted by

 $\sqrt[k]{x} := \alpha$.

4.5 Density of \mathbb{Q} in \mathbb{R}

A set *S* is dense in \mathbb{R} if the elements of *S* are arbitrarily close to the elements of \mathbb{R} .

Definition 4.14: Dense set

Let $S \subseteq \mathbb{R}$. We say that *S* is dense in \mathbb{R} if for every $x \in \mathbb{R}$ and $\varepsilon > 0$, there exist $s \in S$ such that

 $|x-s|<\varepsilon$.

In other words, the above definition is saying that S and \mathbb{R} are tightly knitted together. An equivalent definition of dense set is given below.

Remark 4.15

Let $S \subseteq \mathbb{R}$. They are equivalent:

- S is dense in \mathbb{R} .
- For every pair of numbers $x, y \in \mathbb{R}$ with x < y, there exists $s \in S$ such that

x < s < y.

We now prove that \mathbb{Q} is dense in \mathbb{R} .

Theorem 4.16: Density of \mathbb{Q} in \mathbb{R}

Let $x, y \in \mathbb{R}$, with x < y. There exists $q \in \mathbb{Q}$ such that



Figure 4.6: Let $n \in \mathbb{N}$ be such that 1/n < y - x. Then take *m* so that $m/n \in (x, y)$.

Proof

We need to find $q \in \mathbb{Q}$ such that

$$x < q < y \,. \tag{4.24}$$

By definition of \mathbb{Q} , we have that q has to be of the form q = m/n for $m \in \mathbb{Z}$ and $n \in \mathbb{N}$. Therefore (4.24) is equivalent to finding $m \in \mathbb{Z}$ and $n \in \mathbb{N}$ such that

$$x < \frac{m}{n} < y. \tag{4.25}$$

The idea is to proceed as in Figure 4.6: We take *n* such that 1/n is small enough, so that we can make *m* jumps of size 1/n and end up between *x* and *y*.

To this end, note that y - x > 0 by assumption. By the Archimedean Property in Theorem 2.57 Point 2, there exists $n \in \mathbb{N}$ such that

$$\frac{1}{n} < y - x \,. \tag{4.26}$$

Now, we would like to find $m \in \mathbb{Z}$ such that

$$x < \frac{m}{n}$$
.

However m cannot be too large, because we also want to have

$$\frac{m}{n} < y.$$

The right thing to do, is to take $m \in \mathbb{Z}$ such that

$$\frac{m-1}{n} \le x < \frac{m}{n} \,. \tag{4.27}$$

Why does such *m* exist? The inequality in (4.27) is equivalent to

$$m-1 \le nx < m.$$

As nx>0, by Archimedean Property in Theorem 2.57 Point 1, there exists $m\in\mathbb{Z}$ such that

nx < m

We can then choose m' to be the smallest element in \mathbb{Z} such that

$$nx < m'. \tag{4.28}$$

For such m', we have that

$$m'-1 \le nx < m' \ .$$

Indeed, if by contradiction

 $m'-1>nx\,,$

then m' - 1 would be another integer such that (4.28) holds. Since m' - 1 < m', this contradicts the minimality of m'.

The second inequality in (4.27) implies

$$x < \frac{m}{n}$$
,

which is the first inequality in (4.25). Now note that (4.26) is equivalent to

$$x < y - \frac{1}{n}.$$

We can use the above, and the first inequality in (4.27) to estimate

$$m \le 1 + nx$$

< $1 + n\left(y - \frac{1}{n}\right)$
= ny ,

which yields

 $\frac{m}{n} < y.$

Therefore the second inequality in (4.25) is proven, concluding the proof.

We have constructed the real numbers \mathbb{R} so that they would fill the **gaps** of \mathbb{Q} . Formally, these gaps are the numbers in $\mathbb{R} \setminus \mathbb{Q}$. Let us give a name to this set.

Definition 4.17: Irrational numbers

The set of $\mathbf{irrational}\ \mathbf{numbers}\ \mathbf{in}\ \mathbb{R}$ is

$$\mathscr{I} := \mathbb{R} \setminus \mathbb{Q}.$$

Question 4.18

How many gaps does Q have? In other words, how many irrational numbers are out there?

The answer is quite surprising, and is a corollary of the density result of Theorem 4.16: The irrational numbers are dense in \mathbb{R} .

Corollary 4.19

Let $x, y \in \mathbb{R}$, with x < y. There exists $t \in \mathcal{F}$ such that

x < t < y.

Proof

Consider

 $\tilde{x} := x - \sqrt{2}, \quad \tilde{y} := y - \sqrt{2}.$

Since x < y, we have

 $\tilde{x} < \tilde{y}$.

By Theorem 4.16 there exists $q \in \mathbb{Q}$ such that

 $\tilde{x} < q < \tilde{y}$.

```
Adding \sqrt{2} to the above inequalities we get
```

$$x < t < y, \quad t := q + \sqrt{2}.$$
 (4.29)

We claim that $t \in \mathcal{I}$. Indeed, suppose by contradiction $t \in \mathbb{Q}$. Then

$$\sqrt{2} = t - q \in \mathbb{Q},$$

since $t, q \in \mathbb{Q}$, and \mathbb{Q} is closed under summation. Since $\sqrt{2} \in \mathcal{I}$, we obtain a contradiction. Thus $t \in \mathcal{I}$ and (4.29) is our thesis.

4.6 Cardinality

We have proven that the sets or rational numbers $\mathbb Q$ and irrational numbers $\mathscr I$ are both dense in $\mathbb R$, with

$$\mathbb{R} = \mathbb{Q} \cup \mathcal{F}.$$

From this result we might think that \mathbb{R} is obtained by mixing \mathbb{Q} and \mathscr{I} in equal proportions. This is however false. We will see that \mathbb{R} has much more elements than \mathbb{Q} . Therefore also the set of irrational numbers \mathscr{I} is much larger than \mathbb{Q} .

To make the above discussion precise, we need to introduce the *size* of a set. For this, we need the concept of bijective function.

Definition 4.20: Bijective function

Let *X*, *Y* be sets and $f : X \rightarrow Y$ be a function. We say that:

1. *f* is **injective** if it holds:

$$f(x) = f(y) \implies x = y.$$

2. *f* is **surjective** if it holds:

```
\forall y \in Y, \exists x \in X \text{ s.t. } f(x) = y.
```

3. *f* is **bijective** if it is both **injective** and **surjective**.

In other words: A function $f : X \to Y$ is:

- 1. Injective if any two different elements in *X* are mapped into two different elements in *Y*.
- 2. Surjective if every element in Y has at least one element in X associated via f.
- 3. Bijective if to each element in X we associate one and only one element in Y via f.

Example 4.21: Injectivity

Consider the sets

$$X = \{1, 2, 3\}, \quad Y = \{a, b, c, d, e\}.$$

1. The function $f : X \to Y$ defined by

$$f(1) = c$$
, $f(2) = a$, $f(3) = e$,

is injective.

2. The function $g : X \to Y$ defined by

$$g(1) = c$$
, $g(2) = a$, $g(3) = c$,

is not injective, since

$$g(1) = g(3) = c$$
, $1 \neq 3$.

3. The function $h : \mathbb{R} \to \mathbb{R}$ defined by

$$h(x) = x^2$$

is not injective, since

h(1) = h(-1) = 1, $1 \neq -1$.

4. The function $l : \mathbb{R} \to \mathbb{R}$ defined by

$$l(x) = 2x$$

is injective, since

$$l(x) = l(y) \implies 2x = 2y \implies x = y.$$

Example 4.22: Surjectivity

Consider the sets

$$X = \{1, 2, 3, 4\}, \quad Y = \{a, b, c\}.$$

1. The function $f : X \to Y$ defined by

$$f(1) = c$$
, $f(2) = a$, $f(3) = a$, $f(4) = b$,

is surjective.

2. The function $g : X \to Y$ defined by

$$g(1) = a$$
, $g(2) = a$, $g(3) = c$, $g(4) = a$,

is not surjective, since there is no element $x \in X$ such that

$$g(x) = b$$
.

3. The function $h : \mathbb{R} \to \mathbb{R}$ defined by

$$h(x) = x^2$$

is not surjective, since there is no $x \in \mathbb{R}$ such that

$$h(x) = x^2 = -1.$$

4. The function $l : \mathbb{R} \to [0, \infty)$ defined by

$$l(x) = x^2$$

is surjective, since for every $y \ge 0$ there exists $x \in \mathbb{R}$ such that

$$l(x)=x^2=y\,.$$

This is true by Theorem 4.12.

Example 4.23: Bijectivity

1. Let $X = \{1, 2, 3\}, Y = \{a, b, c\}$. The function $f : X \rightarrow Y$ defined by

f(1) = c, f(2) = a, f(3) = b,

is bijective, since it is both injective and surjective.

2. Let $X = \{1, 2, 3\}, Y = \{a, b\}$. The function $g : X \rightarrow Y$ defined by

g(1) = a, g(2) = b, g(3) = b,

is not bijective, since it is not injective: we have

$$g(2) = g(3) = b$$
 $2 \neq 3$.

3. Let $X = \{1, 2\}, Y = \{a, b, c\}$. The function $h : \mathbb{R} \to \mathbb{R}$ defined by

$$h(1) = a, \quad h(2) = c,$$

is not bijective, since it is not surjective: there is no $x \in X$ such that

$$h(x)=b\,.$$

4. Let $X = \{1, 2, 3\}, Y = \{a, b, c\}$. The function $l : \mathbb{R} \rightarrow [0, \infty)$ defined by

$$l(1) = a$$
, $l(2) = a$, $l(3) = b$,

is not bijective, as it is neither injective nor surjective.

Definition 4.24: Composition of functions

Let X, Y, Z be sets and $f : X \to Y, g : Y \to Z$ functions. The composition of f with g is the function

$$g \circ f : X \to Z$$
, $(g \circ f)(x) := g(f(x))$.

Definition 4.25: Identity map

The identity map on a set X is the function

$$\operatorname{id}_X : X \to X$$
, $\operatorname{id}_X(x) = x$.

Definition 4.26: Invertibility

- Let *X*, *Y* be sets and $f : X \to Y$. We say that:
 - 1. A function $g: Y \to X$ is the **inverse** of f if

$$g \circ f = \operatorname{id}_X, \quad f \circ g = \operatorname{id}_Y.$$

2. f is **invertible** if it admits an inverse g.

Proposition 4.27

Let *X*, *Y* be sets and $f : X \to Y$.

1. If f is invertible, the inverse is unique. We denote it by

$$g := f^{-1}$$
.

- 2. They are equivalent:
 - *f* is invertible.
 - *f* is bijective.

We are ready to define the *size* of a set.

Definition 4.28: Cardinality, Finite, Countable, Uncountable

Let *X* be a set. The **cardinality** of *X* is the number of elements in *X*. We denote the cardinality of *X* by

|X| := # of elements in X.

Further, we say that:

1. *X* is **finite** if there exists a natural number $n \in \mathbb{N}$ and a bijection

$$f: \{1, 2, \dots, n\} \to X.$$

In particular

|X|=n.

2. X is **countable** if there exists a bijection

 $f: \mathbb{N} \to X.$

In this case we denote the cardinality of X by

 $|X| = |\mathbb{N}|.$

3. *X* is **uncountable** if *X* is neither finite, nor countable.

In other words:

1. X is finite if it can be listed as

$$X = \{x_1, \dots, x_n\}, \quad x_i := f(i)$$

for some $n \in \mathbb{N}$.

2. X is countable if it can be listed as

$$X = \{x_n : n \in \mathbb{N}\}, \quad x_n := f(n)$$

3. *X* is uncountable if it *X* cannot be listed.

Remark 4.29

The functions in Definition 4.28 are bijections. Therefore they are invertible, and points 1 and 2 are equivalent to requesting there exist bijections

$$f: X \to \{1, 2, \dots, n\}, \qquad f: X \to \mathbb{N},$$

respectively.

Question 4.30

Is there an intermediate cardinality between finite and countable?

The answer is no, as shown in the next proposition.

Proposition 4.31

Let *X* be a countable set and $A \subseteq X$. Then either *A* is finite or countable.

Proof

If A is finite, there is nothing to prove. Therefore suppose A is not finite. Since X is countable there exists a bijection

$$f: \mathbb{N} \to X.$$

This means X can be listed as

$$X = \{x_n : n \in \mathbb{N}\}, \quad x_n := f(n).$$

Our goal is to list the elements of A. Therefore, we consider the set of indices of elements in A:

$$I_1 := \{ n \in \mathbb{N} \text{ s.t. } f(n) \in A \} = \{ n \in \mathbb{N} : x_n \in A \}.$$

Clearly, $I_1 \neq \emptyset$, since f is bijective and A is a non-empty subset of X. Since I_1 is discrete and

bounded below, there exists $n_1 \in \mathbb{N}$ such that

$$n_1 = \min I_1.$$

This way, x_{n_1} is the first element of A that we encounter in the list of elements of X:

$$X = \{x_1, x_2, x_3, x_4, x_5, \ldots\}.$$

We define

$$g(1) := f(n_1) = x_{n_1}$$

Now, we want to find the second element of *A* in the list of elements of *X*. We need to make sure that we do not pick x_{n_1} again. Therefore, we define the set of indices of elemets in *A* wich are not x_{n_1} :

$$I_2 := \{ n \in \mathbb{N} : n > n_1, f(n) \in A \}$$

= $\{ n \in \mathbb{N} : n > n_1, x_n \in A \}.$

This way I_2 does not contain n_1 , but only successive indices. Note that I_2 is non-empty, since f is surjective and A is an infinite subset of X. Therefore it is well defined

$$n_2 := \min I_2.$$

Notice that by construction

 $n_1 < n_2, \quad x_{n_2} \in A.$

 $g(2) := f(n_2).$

Set

Iterating this procedure, we define

$$I_k := \{ n \in \mathbb{N} : n > n_{k-1}, f(n) \in A \}$$

= $\{ n \in \mathbb{N} : n > n_{k-1}, x_n \in A \}.$

and set

 $n_k = \min I_k, \quad g(k) := f(n_k).$

Note that, by construction, we have

$$n_k > n_{k-1}$$
, $x_{n_k} \in A$, $\forall k \in \mathbb{N}$.

In this way we have defined a function

 $g: \mathbb{N} \to A$.

We have:

- *g* is injective: Suppose g(k) = g(s). Then $f(n_k) = f(n_s)$. From injectivity of *f* we infer $n_k = n_s$. Since $n_k > n_{k-1}$ for all *k*, from the condition $n_k = n_s$ we conclude that k = s. This shows *g* is injective.
- g is surjective: If $x \in A$, by surjectivity of f there exists $\tilde{n} \in \mathbb{N}$ such that $f(\tilde{n}) = x$. Therefore

$$\tilde{n} \in \{n \in \mathbb{N} \text{ s.t. } f(n) \in A\} = I_1.$$

Hence, by construction, there is an index $k \in \mathbb{N}$ such that $n_k = \tilde{n}$. Thus $g(k) = f(n_k) = x$, showing that g is surjective.

Hence g is bijective, showing that A is countable.

Example 4.32

Question. Prove that $X = \{a, b, c\}$ is finite. **Solution.** Set $Y = \{1, 2, 3\}$. The function $f : X \to Y$ defined by

$$f(1) = a$$
, $f(2) = b$, $f(3) = c$,

is bijective. Therefore *X* is finite, with |X| = 3.

Example 4.33

Question. Prove that the set of natural numbers \mathbb{N} is countable. **Solution.** The function $f : X \to \mathbb{N}$ defined by

f(n) := n,

is bijective. Therefore $X = \mathbb{N}$ is countable.

Example 4.34

Question. Let *X* be the set of even numbers

 $X = \{2n : n \in \mathbb{N}\}.$

Prove that *X* is countable. **Solution.** Define the map $f : \mathbb{N} \to X$ by

f(n) := 2n.

We have that:

1. f is injective, because

$$f(m) = f(k) \implies 2m = 2k \quad m = k.$$

2. *f* is surjective: Suppose that $m \in X$. By definition of *X*, there exists $n \in \mathbb{N}$ such that m = 2n. Therefore, f(n) = m.

We have shown that f is bijective. Thus, X is countable.

Example 4.35

Question. Prove that the set of integers \mathbb{Z} is countable. **Solution.** Define $f : \mathbb{N} \to \mathbb{Z}$ by

$$f(n) := \begin{cases} \frac{n}{2} & \text{if } n \text{ even} \\ -\frac{n+1}{2} & \text{if } n \text{ odd} \end{cases}$$

For example

$$\begin{aligned} f(0) &= 0, \quad f(1) = -1, \quad f(2) = 1, \quad f(3) = -2, \\ f(4) &= 2, \quad f(5) = -3, \quad f(6) = 3, \quad f(7) = -4. \end{aligned}$$

We have:

1. *f* is injective: Indeed, suppose that $m \neq n$. If *n* and *m* are both even or both odd we have, respectively

$$f(m) = \frac{m}{2} \neq \frac{n}{2} = f(n)$$

$$f(m) = -\frac{m+1}{2} \neq -\frac{n+1}{2} = f(n)$$

If instead m is even and n is odd, we get

$$f(m) = \frac{m}{2} \neq -\frac{n+1}{2} = f(n),$$

since the LHS is positive and the RHS is negative. The case when m is odd and n even is similar.

2. *f* is surjective: Let $z \in \mathbb{Z}$. If $z \ge 0$, then m := 2z belongs to \mathbb{N} , is even, and

$$f(m)=f(2z)=z\,.$$

If instead z < 0, then m := -2z - 1 belongs to \mathbb{N} , is odd, and

$$f(m)=f(-2z-1)=z.$$

Therefore *f* is bijective, showing that \mathbb{Z} is countable.

We have seen that the sets \mathbb{N} and \mathbb{Z} are countable. What about \mathbb{Q} ? To study this case, we need the following result.

Proposition 4.36

Let the set A_n be countable for all $n \in \mathbb{N}$. Define

$$A=\bigcup_{n\in\mathbb{N}}A_n\,.$$

Then A is countable.

Proof

Since each A_i is countable, we can list their elements as

$$A_i = \{a_k^i : k \in \mathbb{N}\} = \{a_1^i, a_2^i, a_3^i, a_4^i, \ldots\}.$$

The proof that *A* is countable is based on a diagonal argument by Georg Cantor, see Wikipedia page. The idea is that we can list the elements of the sets A_i in an infinite square: In the first row we put the elements of A_1 , in the second row the elements of A_2 , and so on. Therefore the *i*-th row contains the elements of A_i . This procedure is illustrated in Figure 4.7. Therefore this infinite square contains all the elements of A. We then list all the elements of the square by looking at the diagonals, as shown in Figure 4.7. This procedure defines a function $f : \mathbb{N} \to A$. For example the first few terms of f are

$$f(1) = a_1^1$$
, $f(2) = a_2^1$, $f(3) = a_1^2$, $f(4) = a_1^3$.

Since f is injective and surjective, we have that A is countable.



Figure 4.7: The i-th row contains all the elements $a_1^i, a_2^i, a_3^i, ...$ of the countable set A_i . We define the function $f : \mathbb{N} \to A$ by going through the square diagonally.

Theorem 4.37: Q is countable

The set of rational numbers Q is countable.

Proof

For $i \in \mathbb{N}$ define the sets

$$L_i := \left\{ \frac{m}{i} : m \in \mathbb{Z} \right\}.$$

We have that $f: L_i \to \mathbb{Z}$ defined by

$$f\left(\frac{m}{i}\right) := m$$

is a bijection. As \mathbb{Z} is countable, we deduce that L_i is countable. Therefore the set L defined by

$$L := \bigcup_{i \in \mathbb{N}} L_i$$

is countable by Proposition 4.36. Clearly, we have

 $\mathbb{Q} \subseteq L$.

Since *L* is countable, and \mathbb{Q} is not finite, by Proposition 4.31 we conclude that \mathbb{Q} is countable.

We have proven that the sets

 $\mathbb{N}, \mathbb{Z}, \mathbb{Q},$

are all countable. What about \mathbb{R} ?

Theorem 4.38: R is uncountable

The set of Real Numbers \mathbb{R} is **uncountable**.

Proof

Suppose by contradiction \mathbb{R} is countable. Then there exists a bijection $f: \mathbb{N} \to \mathbb{R}$, meaning that we can list the elements of \mathbb{R} as

$$\mathbb{R} = \{x_1, x_2, x_3, x_4, x_5, \ldots\}, \quad x_i := f(i).$$

Let I_1 be a closed interval such that

 $x_1 \not \in I_1 \, .$

Now, we have two possibilities:

1. If $x_2 \in I_1$, then we can find a closed interval I_2 such that

 $x_2 \notin I_2, \quad I_2 \subseteq I_1 \tag{4.30}$

2. If $x_2 \notin I_1$, we define the closed interval $I_2 := I_1$. By construction, (4.30) is satisfied.

To summarize, we have found closed intervals I_1 and I_2 such that

$$x_1 \notin I_1 \,, \quad x_2 \notin I_2 \,, \quad I_2 \subseteq I_1 \,.$$

We can iterate this procedure, and construct a sequence of closed nested intervals I_n such that

$$I_{n+1} \subseteq I_n$$
, $x_n \notin I_n$,

for all $n \in \mathbb{N}$, see Figure 4.8. Since $x_k \notin I_k$, we conclude that

$$x_k \notin \bigcap_{n=1}^{\infty} I_n, \quad \forall k \in \mathbb{N}.$$

As the points x_k are all the elements of \mathbb{R} , we conclude that

$$\bigcap_{n=1}^{\infty} I_n = \emptyset.$$

However, since each I_n is closed, the Nested Interval Property of Theorem 2.59 implies that

$$\bigcap_{n=1}^{\infty} I_n \neq \emptyset,$$

yielding a contradiction. Therefore $\mathbb R$ is uncountable.





As a corollary we obtain that also the irrational numbers are uncountable.

Theorem 4.39

The set of irrational numbers

$$\mathscr{I} := \mathbb{R} \smallsetminus \mathbb{Q}$$

is uncountable.

Proof

In Theorems 2.74, 2.75 we have shown that $\mathbb R$ in uncountable and $\mathbb Q$ is countable. Suppose by contradiction that $\mathcal F$ is countable. Then

 $\mathbb{Q}\cup\mathcal{I}$

is countable by Proposition 4.36, being union of countable sets. Since by definition

 $\mathbb{R} = \mathbb{Q} \cup \mathcal{I},$

we conclude that ${\mathbb R}$ is countable. Contradiction.

5 Complex Numbers

In Section 4.4, we have proven the existence of \sqrt{x} for all for $x \ge 0$. The number \sqrt{x} was defined as

$$\sqrt{x} := \alpha$$
, $\alpha := \sup\{t \in \mathbb{R} : t^2 < x\}$

We then proved that

$$\alpha^2 = x$$

This procedure is possible for any $x \ge 0$.

Question 5.1

Is there a number $\alpha \in \mathbb{R}$ such that

$$\alpha^2 = -1$$
 ?

(5.1)

The answer to the above question is no. This is because \mathbb{R} is an ordered field, and from axiom (MO) it follows that:

$$x^2 \ge 0$$
, $\forall x \in \mathbb{R}$.

However we would still like to solve equation (5.1) somehow. To do this, we introduce the **imaginary numbers** or **complex numbers**. We define *i* to be that *number* such that

$$i^2 = -1$$
.

Formally, we can also think of *i* as

$$i = \sqrt{-1}$$

We can use this speacial number to define the square root of a negative real number x < 0:

$$\sqrt{x} := i\sqrt{-x}$$
.

Note that $\sqrt{-x}$ is properly defined in R, because -x is positive when x is negative.

5.1 The field ℂ

We would like to be able to do calculations with the newly introduced complex numbers, and investigate their properties. We can introduce them rigorously as a field, as we did for \mathbb{R} .

Definition 5.2: Complex Numbers

The set of complex numbers $\mathbb C$ is defined as

$$\mathbb{C} := \mathbb{R} \oplus i\mathbb{R} := \{ x \oplus iy : x, y \in \mathbb{R} \}.$$

In the above the symbol \oplus is used to denote the pair

 $x \oplus iy = (x, y)$

with $x, y \in \mathbb{R}$. This means that x and y play different *roles*.

Definition 5.3

For a complex number

$$z = x \oplus iy \in \mathbb{C}$$

we say that

• *x* is the **real part** of *z*, and denote it by

 $x = \operatorname{Re}(z)$

• *y* is the **imaginary part** of *z*, and denote it by

 $y = \operatorname{Im}(z)$

We say that

- If $\operatorname{Re} z = 0$ then z is a **purely imaginary** number.
- If Im z = 0 then z is a **real** number.

In order to make the set \mathbb{C} into a field, we first have to define the two binary operations of addition + and multiplication \cdot ,

 $+, \cdot : \mathbb{C} \times \mathbb{C} \to \mathbb{C}$.

Then we need to prove that these operations satisfy all the field axioms.

Definition 5.4: Addition in \mathbb{C}

Let $z_1, z_2 \in \mathbb{C}$, so that

 $z_1 = x_1 \oplus i y_1 \ , \quad z_2 = x_2 \oplus i y_2 \ ,$

for some $x_1, x_2, y_1, y_2 \in \mathbb{R}$. We define the sum of z_1 and z_2 as

 $z_1 + z_2 = (x_1 \oplus iy_1) + (x_2 \oplus iy_2) := (x_1 + x_2) \oplus i(y_1 + y_2)$

where the + symbol on the right hand side is the addition operator in $\mathbb R.$

Clearly, $z_1 + z_2$ as defined above is an element of \mathbb{C} . Therefore + defines a binary operation over \mathbb{C} .

Notation 5.5

From the above definition, we have that, for all $x, y \in \mathbb{R}$,

 $(x \oplus i0) + (0 \oplus iy) = x \oplus iy.$

To simplify notation, we will write

 $x \oplus i0 = x$, $0 \oplus iy = iy$

and

 $x \oplus iy = x + iy$.

We will also often swap i and y, writing equivalently

x + iy = x + yi.

We now want to define multiplication between complex numbers.

Remark 5.6: Formal calculation for multiplication in \mathbb{C}

How to define multiplication in $\mathbb{C}?$ Whatever the definition may be, at least it has to give that that

 $i^2 = i \cdot i = -1 \, .$

Keeping the above in mind, let us do some formal calculations: For $z_1 = x_1 + iy_1$, $z_2 = x_2 + iy_2$ we have

$$z_1 \cdot z_2 = (x_1 + iy_1) \cdot (x_2 + iy_2)$$

= $x_1 \cdot x_2 + x_1 \cdot iy_2 + x_2 \cdot iy_1 + y_1 \cdot i^2 y_2$
= $(x_1 \cdot x_2 - y_1 \cdot y_2) + i(x_1 \cdot y_2 + x_2 \cdot y_1)$

Remark 5.6 motivates the following definition of multiplication.

Definition 5.7: Multiplication in \mathbb{C}

Let $z_1, z_2 \in \mathbb{C}$, so that

 $z_1 = x_1 \oplus iy_1 , \quad z_2 = x_2 \oplus iy_2 ,$

for some $x_1, x_2, y_1, y_2 \in \mathbb{R}$. Define the multiplication of z_1 and z_2 as

$$z_1 \cdot z_2 = (x_1 + iy_1) \cdot (x_2 + iy_2)$$

= $(x_1 \cdot x_2 - y_1 \cdot y_2) + i(x_1 \cdot y_2 + x_2 \cdot y_1),$

where the operations + and \cdot on the right hand side are the operations in \mathbb{R} .

Clearly, $z_1 \cdot z_2$ as defined above is an element of \mathbb{C} . Therefore \cdot defines a binary operation over \mathbb{C} .

Remark 5.8

To check that we have given a good definition of product, we should have that

 $i^2 = -1$,

as expected. Indeed:

$$i^{2} = (0 + 1i) \cdot (0 + 1i)$$

= (0 \cdot 0 - 1 \cdot 1) + (0 \cdot 1 + 0 \cdot 1)i = -1.

Important

In view of Remark 5.8, we see that he formal calculations in Remark 5.6 are compatible with the definition of multiplication of complex numbers. Therefore, it is not necessary to memorize the multiplication formula, but it suffices to carry out calculations as usual, and replace i^2 by -1.

Example 5.9

Question. Compute *zw*, where

$$z = -2 + 3i$$
, $w = 1 - i$.

Solution. Using the definition we compute

$$z \cdot w = (-2 + 3i) \cdot (1 - i)$$

= (-2 - (-3)) + (2 + 3)i
= 1 + 5i.

Alternatively, we can proceed formally as in Remark 5.6. We just need to recall that i^2 has to be replaced with -1:

$$z \cdot w = (-2 + 3i) \cdot (1 - i)$$

= -2 + 2i + 3i - 3i²
= (-2 + 3) + (2 + 3)i
= 1 + 5i.

We now want to check that

 $(\mathbb{C},+,\cdot)$

is a field. All the field axioms are trivial to check, except for the existence of additive and multiplicative inverses.

Proposition 5.10: Additive inverse in \mathbb{C}

The neutral element of addition in $\mathbb C$ is the number

0 := 0 + 0i.

For any $z = x + iy \in \mathbb{C}$, the unique additive inverse is given by

-z := -x - iy.

The proof is immediate and is left as an exercise. The multiplication requires more care.

Remark 5.11: Formal calculation for multiplicative inverse

Let us first carry our some formal calculations. Let

$$z = x + iy \in \mathbb{C}, \quad z \neq 0.$$

First, note that

$$z \cdot 1 = (x + iy) \cdot (1 + 0i) = x + iy = z$$

and therefore 1 is the neutral element of multiplication. Thus, the inverse of z should be a complex number $z^{-1} \in \mathbb{C}$ such that

 $z \cdot z^{-1} = 1 \,.$

We would like to define

$$z^{-1} = \frac{1}{x + iy} \, .$$

Such number does not belong to \mathbb{C} , as it is not of the form a + ib for some $a, b \in \mathbb{R}$. However it is what the inverse should look like. Proceeding formally:

$$\frac{1}{x+iy} = \frac{1}{x+iy} \cdot 1$$
$$= \frac{1}{x+iy} \cdot \frac{x-iy}{x-iy}$$
$$= \frac{x-iy}{x^2 - (iy)^2}$$
$$= \frac{x-iy}{x^2 + y^2}$$
$$= \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}.$$

The right hand side is an element of \mathbb{C} , and looks like a good candidate for z^{-1} .

Motivated by the above remark, we define inverses in $\mathbb C$ in the following way.

Proposition 5.12: Multiplicative inverse in \mathbb{C}

The neutral element of multiplication in $\mathbb C$ is the number

$$1 := 1 + 0i$$
.

For any $z = x + iy \in \mathbb{C}$, the unique multiplicative inverse is given by

$$z^{-1} := \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}.$$

Proof

It is immediate to check that 1 is the neutral element of multiplication in $\mathbb{C}.$ For the remaining part of the statement, set

$$w := \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}.$$

We need to check that $z \cdot w = 1$

$$z \cdot w = (x + iy) \cdot \left(\frac{x}{x^2 + y^2} + i\frac{-y}{x^2 + y^2}\right)$$
$$= \left(\frac{x^2}{x^2 + y^2} - \frac{y \cdot (-y)}{x^2 + y^2}\right) + i\left(\frac{x \cdot (-y)}{x^2 + y^2} + \frac{xy}{x^2 + y^2}\right)$$
$$= 1,$$

so indeed $z^{-1} = w$.

Important

It is not necessary to memorize the formula for z^{-1} . Indeed one can just remember the trick of multiplying by

$$1=\frac{x-iy}{x-iy},$$

and proceed formally, as done in Remark 5.11.

Example 5.13

Question. Let z = 3 + 2i. Compute z^{-1} . **Solution.** By the formula in Propostion 5.12 we immediately get

$$z^{-1} = \frac{3}{3^2 + 2^2} + \frac{-2}{3^2 + 2^2}i = \frac{3}{13} - \frac{2}{13}i$$

Alternatively, we can proceed formally as in Remark 5.11

$$(3+2i)^{-1} = \frac{1}{3+2i}$$
$$= \frac{1}{3+2i} \frac{3-2i}{3-2i}$$
$$= \frac{3-2i}{3^2+2^2}$$
$$= \frac{3}{13} - \frac{2}{13}i,$$

and obtain the same result.

We can now prove that \mathbb{C} is a field.

Theorem 5.14

 $(\mathbb{C}, +, \cdot)$ is a field.

Proof

We need to check that all field axioms hold. For the addition we have

• (A1) To show that + is commutative, note that

$$(x + iy) + (a + ib) = (x + a) + i(y + b)$$

= (a + x) + i(b + y)
= (a + ib) + (x + iy).

where we used Definition 2.78 in the first and last equality, and the commutative property of the real numbers (which holds since by definition \mathbb{R} is a field) in the second equality. Associativity can be checked in the same way.

- (A2) The neutral element of addition is 0, as stated in Proposition 2.80.
- (A₃) Existence of additive inverses is given by Proposition 2.80.

For multiplication we have:

- (M1) Commutativity and associativity of product in \mathbb{C} can be checked using Definition 5.7 and commutativity and associativity of sum and multiplication in \mathbb{R} .
- (M2) The neutral element of multiplication is 1, as stated in Propostion 5.12.
- (M₃) Existence of multiplicative inverses is guaranteed by Proposition 5.12.

Finally one should check the associative property (AM). This is left as an exercise.

5.1.1 Division in $\ensuremath{\mathbb{C}}$

Suppose we want to divide two complex numbers $w, z \in \mathbb{C}, z \neq 0$, with

$$z = x + iy$$
, $w = a + ib$.

We have two options:

1. Use the formula for the inverse from Proposition 5.12 and compute

$$z^{-1} := \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}.$$

Then we use the multiplication formula of Definition 5.7 to compute

$$\frac{w}{z} = w \cdot z^{-1}$$
$$= (a + ib) \cdot \left(\frac{x}{x^2 + y^2} + i\frac{-y}{x^2 + y^2}\right)$$
$$= \frac{(ax + by) + i(bx - ay)}{x^2 + y^2}$$

2. Proceed formally as in Remark 5.11, using the multiplication by 1 trick. We would have

$$\frac{w}{z} = \frac{a+ib}{x+iy}$$
$$= \frac{a+ib}{x+iy} \frac{x-iy}{x-iy}$$
$$= \frac{(ax+by)+i(bx-ay)}{x^2+y^2}$$

Example 5.15

Question. Let w = 1 + i and z = 3 - i. Compute $\frac{w}{z}$. **Solution.** We compute w/z using the two options we have:

1. Using the formula for the inverse from Proposition 5.12 we compute

$$z^{-1} = \frac{x}{x^2 + y^2} + i \frac{-y}{x^2 + y^2}$$
$$= \frac{3}{3^2 + 1^2} - i \frac{-1}{3^2 + 1^2}$$
$$= \frac{3}{10} + \frac{1}{10}i$$

and therefore

$$\frac{w}{z} = w \cdot z^{-1}$$

= (1+i) $\left(\frac{3}{10} + \frac{1}{10}i\right)$
= $\left(\frac{3}{10} - \frac{1}{10}\right) + \left(\frac{1}{10} + \frac{3}{10}\right)i$
= $\frac{2}{10} + \frac{4}{10}i$
= $\frac{1}{5} + \frac{2}{5}i$

2. We proceed formally, using the multiplication by 1 trick. We have

$$\frac{w}{z} = \frac{1+i}{3-i}$$

= $\frac{1+i}{3-i}\frac{3+i}{3+i}$
= $\frac{3-1+(3+1)i}{3^2+1^2}$
= $\frac{2}{10} + \frac{4}{10}i$
= $\frac{1}{5} + \frac{2}{5}i$

5.1.2 C is not ordered

We have seen that $(\mathbb{C}, +, \cdot)$ is a field. One might wonder whether \mathbb{C} is also an ordered field. It turns out that this is not the case.

Theorem 5.16

The field $(\mathbb{C}, +, \cdot)$ is not ordered.

Proof

Suppose that \mathbb{C} is an ordered field, that is, there exists an order relation \leq on \mathbb{C} compatible with the operations + and \cdot . By axiom (MO) it follows that for all elements $z \in \mathbb{C}$, $z \neq 0$, we have that $z^2 > 0$. But since $i^2 = -1 < 0$, we get a contradiction.

Hence, it is not possible to compare two complex numbers.

5.1.3 Completeness of $\mathbb C$

One might also wonder whether \mathbb{C} is complete. Our definition of completeness uses the notions of supremum and infimum, which require an order relation. However, \mathbb{C} is not ordered, as seen in Theorem 5.16. Therefore we cannot define infimum and supremum of subsets of \mathbb{C} , and we have to find an alternative way to discuss completeness.

The right way to introduce completeness in \mathbb{C} is by using the notion of **Cauchy sequence**. In ordered fields, this new definition of completeness is equivalent to the definition which uses the supremum.

5.2 Complex conjugates

When computing inverses, we used the trick to multiply by 1:

$$z^{-1} = \frac{1}{z} \cdot 1 = \frac{1}{x + iy} \cdot \frac{x - iy}{x - iy}.$$

The complex number x - iy is obtained by changing the sign to the imaginary part of z = x + iy. We give a name to this operation.

Definition 5.17: Complex conjugate

Let z = x + iy. We call the **complex conjugate** of *z*, denoted by \overline{z} , the complex number

 $\bar{z} = x - iy$.

Example 5.18

We have the following conjugates:

$$\overline{3+4i} = 3 - 4i, \qquad \overline{3-4i} = 3 + 4i,
\overline{-3+4i} = -3 - 4i, \qquad \overline{-3-4i} = -3 + 4i
\overline{3} = 3, \qquad \overline{4i} = -4i.$$

Complex conjugates have the following properties:

Theorem 5.19

For all $z_1, z_2 \in \mathbb{C}$ it holds:

- $\overline{z_1 + z_2} = \overline{z_1} + \overline{z_2}$
- $\overline{z_1 \cdot z_2} = \overline{z_1} \cdot \overline{z_2}$

Proof

Let $z_1, z_2 \in \mathbb{C}$. Then

 $z_1 = x_1 + iy_1$, $z_2 = x_2 + iy_2$,

for some $x_1, y_1, x_2, y_2 \in \mathbb{R}$.

- Using the definition of addition in $\mathbb C$ and of conjugate,

$$\overline{z_1 + z_2} = \overline{(x_1 + iy_1) + (x_2 + iy_2)}$$
$$= \overline{(x_1 + x_2) + i(y_1 + y_2)}$$
$$= (x_1 + x_2) - i(y_1 + y_2)$$
$$= (x_1 - iy_1) + (x_2 - iy_2)$$
$$= \overline{x_1 + iy_1} + \overline{x_2 + iy_2}$$
$$= \overline{z_1} + \overline{z_2}.$$

- Using the definition of multiplication in $\mathbb C$ and of conjugate,

$$\overline{z_1 \cdot z_2} = \overline{(x_1 + iy_1) \cdot (x_2 + iy_2)}$$
$$= \overline{(x_1 x_2 - y_1 y_2) + i(x_1 y_2 + x_2 y_1)}$$
$$= (x_1 x_2 - y_1 y_2) - i(x_1 y_2 + x_2 y_1)$$
$$= (x_1 - iy_1) \cdot (x_2 - iy_2)$$
$$= \overline{z_1} \cdot \overline{z_2}$$

Example 5.20

Let $z_1 = 3 - 4i$ and $z_2 = -2 + 5i$. Then

· Let us check that

$$\overline{z_1 + z_2} = \overline{z_1} + \overline{z_2}$$

Indeed, we have

$$z_1 + z_2 = 1 + i \quad \Longrightarrow \quad \overline{z_1 + z_2} = 1 - i.$$

On the other hand

$$\overline{z_1} = 3 + 4i, \quad \overline{z_2} = -2 - 5i \quad \Longrightarrow \quad \overline{z_1} + \overline{z_2} = 1 - i.$$

Let us check that

$$\overline{z_1 \cdot z_2} = \overline{z_1} \cdot \overline{z_2}$$

Indeed,

$$z_1 \cdot z_2 = (3+4i) \cdot (-2+5i)$$

= (-6+20) + (8+15)i
= 14+23i

so that

 $\overline{z_1 \cdot z_2} = 14 - 23i$

On the other hand:

$$\overline{z_1} \cdot \overline{z_2} = (3+4i) \cdot (-2-5i)$$
$$= (-6+20) + (-15-8)i$$
$$= 14-23i$$

5.3 The complex plane

We can represent a real number *x* as a point on the one-dimensional real line \mathbb{R} . The distance between two real numbers $x, y \in \mathbb{R}$ on the real line is given by |x - y|, see Figure 5.1.



Figure 5.1: Two points *x* and *y* on the real line \mathbb{R} . Their distance is |x - y|.

We would like to do something similar for the complex numbers, i.e. points

$$z = x + iy, \quad x, y \in \mathbb{R}$$

We represent z = x + iy in the two-dimensional plane at the point with (Cartesian) coordinates (x, y). This two-dimensional plane in which we can represent all complex numbers is called the **complex plane**. The origin of such plane, with coordinates (0, 0), corresponds to the complex number

$$0+0i=0,$$

see Figure 5.2.

5.3.1 Distance on C

The Cartesian representation allows us to introduce a distance between two complex numbers. Let us start with the distance between a complex number z = x + iy and 0. By Pythagoras Theorem this



Figure 5.2: A point $z = x+iy \in \mathbb{C}$ can be represented on the complex plane by the point of coordinates (x, y). The distance between z and 0 is given by $|z| = \sqrt{z^2 + y^2}$.

distance is given by

$$\sqrt{x^2 + y^2}$$
,

see Figure 5.2. We give a name to this quantity.

Definition 5.21: Modulus

The **modulus** of a complex number z = x + iy is defined by

$$|z| := \sqrt{x^2 + y^2}$$
.

Note that the distance between z and 0 is always a non-negative number.

Remark 5.22: Modulus of Real numbers

A real number $x \in \mathbb{R}$ can be written as

$$x = x + 0i \in \mathbb{C}.$$

Hence the modulus of x is given by

$$|x| = \sqrt{x^2 + 0^2} = \sqrt{x^2} \,.$$

The above coincides with the absolute value of x. This explains why the notation for modulus in \mathbb{C} is the same as the one for absolute value in \mathbb{R} .

We can now define the distance between two complex numbers.

Definition 5.23: Distance in \mathbb{C}

Given $z_1, z_2 \in \mathbb{C}$, we define the **distance** between z_1 and z_2 as the quantity

 $|z_1 - z_2|$.

The geometric intuition of why the quantity $|z_1 - z_2|$ is defined as the distance between z_1 and z_2 is given in Figure 5.3.



Figure 5.3: The difference $z_1 - z_2$ of the two points $z_1, z_2 \in \mathbb{C}$ is given by the magenta vector. We define $|z_1 - z_2|$ as the distance between z_1 and z_2 .

Theorem 5.24

Given $z_1, z_2 \in \mathbb{C}$, we have

$$|z_1 - z_2| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Proof

We have

$$z_1 - z_2 = (x_1 - x_2) + i(y_1 - y_2).$$

Therefore, by definition of modulus,

$$|z_1 - z_2| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

Example 5.25

Question. Compute the distance between

$$z = 2 - 4i$$
, $w = -5 + i$.

Solution. The distance is

$$|z - w| = |(2 - 4i) - (-5 + i)|$$

= |7 - 5i|
= $\sqrt{7^2 + (-5)^2}$
= $\sqrt{74}$

5.3.2 Properties of modulus

The modulus has the following properties.

Theorem 5.26

Let $z, z_1, z_2 \in \mathbb{C}$. Then 1. $|z_1 \cdot z_2| = |z_1| |z_2|$ 2. $|z^n| = |z|^n$ for all $n \in \mathbb{N}$ 3. $z \cdot \bar{z} = |z|^2$

Proof

Part 1. We have

$$z_1 \cdot z_2 = (x_1 + iy_1) \cdot (x_2 + iy_2)$$

= $(x_1 x_2 - y_1 y_2) + i(x_2 y_1 + x_1 y_2)$

and therefore

$$\begin{aligned} |z_1 \cdot z_2| &= \sqrt{(x_1 x_2 - y_1 y_2)^2 + (x_2 y_1 + x_1 y_2)^2} \\ &= \sqrt{x_1^2 x_2^2 + y_1^2 y_2^2 + x_2^2 y_1^2 + x_1^2 y_2^2} \,. \end{aligned}$$

On the other hand,

$$|z_1| = \sqrt{x_1^2 + y_1^2}, \quad |z_2| = \sqrt{x_2^2 + y_2^2}$$

so that

$$\begin{aligned} |z_1||z_2| &= \sqrt{x_1^2 + y_1^2} \sqrt{x_2^2 + y_2^2} \\ &= \sqrt{x_1^2 x_2^2 + y_1^2 y_2^2 + x_2^2 y_1^2 + x_1^2 y_2^2} \end{aligned}$$

proving that $|z_1 \cdot z_2| = |z_1| |z_2|$.

Part 2. Exercise. It easily follows from Point 1 and induction. *Part 3.* Let z = x + iy for some $x, y \in \mathbb{R}$. Then,

$$z \cdot \overline{z} = (x + iy)(x - iy)$$
$$= x^{2} - (iy)^{2}$$
$$= x^{2} + y^{2}$$
$$= |z|^{2}$$

The modulus in $\mathbb C$ satisfies the triangle inequality.

Theorem 5.27: Triangle inequality in ℂ

For all $x, y, z \in \mathbb{C}$, 1. $|x + y| \le |x| + |y|$

2. $|x - z| \le |x - y| + |y - z|$

Proof

Part 1. Suppose that x = a + ib and y = c + id for $a, b, c, d \in \mathbb{R}$. Then,

$$|x + y| = |(a + c) + i(b + d)| = \sqrt{(a + c)^2 + (b + d)^2}.$$

Therefore the inequality

$$|x + y| \le |x| + |y|$$
(5.2)

is equivalent to

$$\sqrt{(a+c)^2 + (b+d)^2} \le \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2} \,. \tag{5.3}$$

Now note that, for $A, B \in \mathbb{R}$, we have that

$$A^2 \le B^2 \quad \Longrightarrow \quad |A| \le |B| \,. \tag{5.4}$$
In the two reverse implications \leftarrow below we will use (5.4):

$$\begin{split} \sqrt{(a+c)^2 + (b+d)^2} &\leq \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2} \\ &\Leftarrow (a+c)^2 + (b+d)^2 \leq \left(\sqrt{a^2 + b^2} + \sqrt{c^2 + d^2}\right)^2 \\ &\Leftrightarrow a^2 + 2ac + c^2 + b^2 + 2bd + d^2 \leq a^2 + b^2 + 2\sqrt{a^2 + b^2}\sqrt{c^2 + d^2} + c^2 + d^2 \\ &\Leftrightarrow ac + bd \leq \sqrt{a^2 + b^2}\sqrt{c^2 + d^2} \\ &\Leftarrow (ac + bd)^2 \leq (a^2 + b^2)(c^2 + d^2) \\ &\Leftrightarrow a^2c^2 + 2abcd + b^2d^2 \leq a^2c^2 + a^2d^2 + b^2c^2 + b^2d^2 \\ &\Leftrightarrow a^2d^2 + b^2c^2 - 2abcd \geq 0 \\ &\Leftrightarrow (ad - bc)^2 \geq 0. \end{split}$$

This last statement is clearly true, since $ad - bc \in \mathbb{R}$. Therefore (5.3) holds, and so (5.2) follows. *Part 2.* Using (5.2) we estimate

$$|x - z| = |x - y + y - z| \le |x - y| + |y - z|.$$

Remark 5.28: Geometric interpretation of triangle inequality

We finally have a justification of why the inequality

$$|x-z| \le |x-y| + |y-z|$$

is called **triangle inequality**: By drawing three points $x, y, z \in \mathbb{C}$ in the complex plane, the distance between x and z is shorter than the distance to go from x to z via the point y, see Figure 5.4.

5.4 Polar coordinates

We have seen that we can identify a complex number z = x + iy by a point in the complex plane with Cartesian coordinates (x, y). We can also specify the point (x, y) by using the so-called polar coordinates (ρ, θ) , where

ρ is the distance between *z* and the origin

$$\rho = |z| = \sqrt{x^2 + y^2}$$

• θ is the angle between the line connecting the origin and z and the positive real axis, see Figure 5.5.



Figure 5.4: Let $x, y, z \in \mathbb{C}$. The distance between x and z is shorter than the distance to go from x to z via the point y.



Figure 5.5: Polar coordinates (ρ, θ) for the complex number $z \in \mathbb{C}$.

We give such angle a name.

Definition 5.29: Argument

Let $z \in \mathbb{C}$. The angle θ between the line connecting the origin and z and the positive real axis is called the **argument** of z, and is denoted by

 $\theta := \arg(z).$

Warning

We always use angles in radians, not degrees. Make sure your calculator is set to radians if you want to use it to compute angles.

Remark 5.30: Principal Value

The argument of a complex number is not uniquely defined. We can always add an integer number of times 2π to the argument to specify the same point. We usually use the convention to choose the argument in the interval $(-\pi, \pi]$. This is called the **principal value** of the argument function. Therefore the complex numbers in the upper half plane have a positive argument, and the ones in the lower half plane have a negative argument.

Example 5.31

We have the following arguments:

$$arg(1) = 0 arg(i) = \frac{\pi}{2}$$

$$arg(-1) = \pi arg(-i) = -\frac{\pi}{2}$$

$$arg(1+i) = \frac{1}{4}\pi arg(-1-i) = -\frac{3}{4}\pi$$

We can represent any non-zero complex number in polar coordinates.

Theorem 5.32: Polar coordinates

Let $z \in \mathbb{C}$ with z = x + iy and $z \neq 0$. Then

$$x = \rho \cos(\theta), \quad y = \rho \sin(\theta),$$

where

$$\rho := |z| = \sqrt{x^2 + y^2}, \quad \theta := \arg(z).$$

The proof of Theorem 2.95 is trivial, and is based on basic trigonometry and definition of $\arg(z)$.

Complex numbers in polar form can be useful. We give a name to such polar form.

Definition 5.33: Trigonometric form

Let $z \in \mathbb{C}$. The trigonometric form of z is

$$z = |z| \left[\cos(\theta) + i\sin(\theta)\right],$$

where $\theta = \arg(z)$.

Example 5.34

Question. Suppose that $z \in \mathbb{C}$ has polar coordinates

$$\rho = \sqrt{8}, \quad \theta = \frac{3}{4}\pi.$$

Therefore, the trigonometric form of z is

$$z = \sqrt{8} \left[\cos\left(\frac{3}{4}\pi\right) + i\sin\left(\frac{3}{4}\pi\right) \right] \,.$$

Write *z* in cartesian form. **Solution.** We have

$$x = \rho \cos(\theta) = \sqrt{8} \cos\left(\frac{3}{4}\pi\right) = -\sqrt{8} \cdot \frac{\sqrt{2}}{2} = -2$$
$$y = \rho \sin(\theta) = \sqrt{8} \sin\left(\frac{3}{4}\pi\right) = \sqrt{8} \cdot \frac{\sqrt{2}}{2} = 2.$$

Therefore, the cartesian form of z is

$$z = x + iy = -2 + 2i.$$

As a consequence of Theorem 2.95 we obtain a formula for computing the argument.

Corollary 5.35: Computing arg(z)

Let $z \in \mathbb{C}$ with z = x + iy and $z \neq 0$. Then

$$\arg(z) = \begin{cases} \arctan\left(\frac{y}{x}\right) & \text{if } x > 0\\ \arctan\left(\frac{y}{x}\right) + \pi & \text{if } x < 0 \text{ and } y \ge 0\\ \arctan\left(\frac{y}{x}\right) - \pi & \text{if } x < 0 \text{ and } y < 0\\ \frac{\pi}{2} & \text{if } x = 0 \text{ and } y > 0\\ -\frac{\pi}{2} & \text{if } x = 0 \text{ and } y < 0 \end{cases}$$

where arctan is the inverse of tan.

Proof

Using the polar coordinates formulas from Theorem 2.95, we get

$$\frac{y}{x} = \frac{\rho \sin(\theta)}{\rho \cos(\theta)} = \tan(\theta).$$

The thesis can be obtained by carefully inverting the tangent: With reference to Figure 5.6 and Figure 5.7, we have:

1. When x > 0: Then z = x + iy belongs to the 1st or 4th quadrant. Therefore the argument of z is $\theta \in (-\pi/2, \pi/2)$. Such interval coincides with the domain of tan, and therefore

$$\theta = \arctan\left(\frac{y}{x}\right)$$
.

2. When x < 0 and $y \ge 0$: Then z = x + iy lies in the 2nd quadrant. Thus, the argument of z is $\theta \in (\pi/2, \pi)$. Hence

$$\theta = \arctan\left(\frac{y}{x}\right) + \pi$$
.

3. When x < 0 and y < 0: Then z = x + iy lies in the 3rd quadrant. Thus, the argument of z is $\theta \in (-\pi, \pi/2)$. Hence

$$\theta = \arctan\left(\frac{y}{x}\right) - \pi$$
.

- 4. When x = 0 and y > 0: Then z = iy is imaginary with y > 0, meaning that $\theta = \pi/2$.
- 5. When x = 0 and y < 0: Then z = iy is imaginary with y < 0, meaning that $\theta = -\pi/2$.

This exhausts all the cases, and the proof is concluded.

Example 5.36

Question. Compute the arguments of the complex numbers

z = 3 + 4i, $\bar{z} = 3 - 4i$, $-\bar{z} = -3 + 4i$, -z = -3 - 4i.



Figure 5.6: The definition of arg(z) depends on the position of z in the complex plane.



Figure 5.7: Plot of the tangent function for θ between $-\frac{3\pi}{2}$ and $\frac{3\pi}{2}$.

Solution. Using the formula for arg in Corollary 2.98 we have

$$\arg(3+4i) = \arctan\left(\frac{4}{3}\right)$$
$$\arg(3-4i) = \arctan\left(-\frac{4}{3}\right) = -\arctan\left(\frac{4}{3}\right)$$
$$\arg(-3+4i) = \arctan\left(-\frac{4}{3}\right) + \pi = -\arctan\left(\frac{4}{3}\right) + \pi$$
$$\arg(-3-4i) = \arctan\left(\frac{4}{3}\right) - \pi$$

5.5 Exponential form

We have seen that we can represent complex numbers in

- Cartesian form
- Trigonometric form

We now introduce a third way of representing complex numbers: the exponential form. For this, we need Euler's identity:

Theorem 5.37: Euler's identity

For all $\theta \in \mathbb{R}$ it holds

$$e^{i\theta} = \cos(\theta) + i\sin(\theta)$$
.

Proof

The proof of this theorem uses complex power series. We have not yet introduced series. However, we carry out the proof formally, assuming that all the quantities below converge. We have the following Taylor series at $x_0 = 0$ (you might know them from calculus):

$$e^{x} = 1 + x + \frac{x^{2}}{2!} + \frac{x^{3}}{3!} + \frac{x^{4}}{4!} + \frac{x^{5}}{5!} + \frac{x^{6}}{6!} + \frac{x^{7}}{7!} + \dots$$

$$\sin(x) = \frac{x}{1!} - \frac{x^{3}}{3!} + \frac{x^{5}}{5!} - \frac{x^{7}}{7!} + \dots$$

$$\cos(x) = 1 - \frac{x^{2}}{2!} + \frac{x^{4}}{4!} - \frac{x^{6}}{6!} + \dots$$

The above identities also hold for $x \in \mathbb{C}$. Hence we can substitute $x = i\theta$ in the series for e^x to

obtain

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} \dots \\ &= 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} - \frac{\theta^6}{6!} - i\frac{\theta^7}{7!} + \dots \\ &= \cos(\theta) + i\sin(\theta), \end{aligned}$$

where in the second equality we used that $i^2 = -1$, and where the third equality follows by observing that all terms with an even power of θ are exactly the terms in the expansion of $\cos(\theta)$, and all terms with an odd power of θ are exactly the terms in the expansion of $\sin(\theta)$ multiplied by *i*.

Theorem 5.38

For all $\theta \in \mathbb{R}$ it holds

$$\left|e^{i\theta}\right|=1$$
.

Proof

From Euler's identity in Theorem 2.100 we get

$$|e^{i\theta}| = |\cos(\theta) + i\sin(\theta)| = \sqrt{\cos^2(\theta) + \sin^2(\theta)} = 1.$$

Theorem 5.39

Let $z \in \mathbb{C}$ with z = x + iy and $z \neq 0$. Then

$$z = \rho e^{i\theta}$$

where

$$ho := |z| = \sqrt{x^2 + y^2}, \qquad heta := \arg(z).$$

Proof

By Theorem 2.95 we have

$$x = \rho \cos(\theta), \quad y = \rho \sin(\theta)$$

Hence

$$z = x + iy$$

= $\rho \cos(\theta) + i\rho \sin(\theta)$
= $\rho e^{i\theta}$,

where in the last line we used Euler's identity in Theorem 2.100.

Definition 5.40: Exponential form

The **exponential form** of a complex number $z \in \mathbb{C}$ is

$$z = \rho e^{i\theta} = |z| e^{i \arg(z)}$$

Example 5.41

Question. Write the number

z = -2 + 2i

in exponential form.

Solution. From Example 5.34 we know that z = -2 + 2i can be written in trigonometric form as

$$z = \sqrt{8} \left[\cos\left(\frac{3}{4}\pi\right) + i \sin\left(\frac{3}{4}\pi\right) \right].$$

By Euler's identity we hence obtain the exponential form

 $z=\sqrt{8}e^{i\frac{3}{4}\pi}\,.$

Remark 5.42: Periodicity of exponential

For all $k \in \mathbb{Z}$ we have

$$e^{i\theta} = e^{i(\theta + 2\pi k)},\tag{5.5}$$

meaning that the complex exponential is 2π -periodic. As we did for the principal value of the argument, we select θ in (5.5) so that $\theta \in (-\pi, \pi]$.

Proof

Equation (5.5) follows immediately by Euler's identity and periodicity of cos and sin, since

$$e^{i(\theta+2\pi k)} = \cos(\theta+2\pi k) + i\sin(\theta+2\pi k)$$
$$= \cos(\theta) + i\sin(\theta) = e^{i\theta}.$$

The exponential form is very useful for computing products and powers of complex numbers.

Proposition 5.43

Let $z, z_1, z_2 \in \mathbb{C}$ and suppose that

$$z=
ho e^{i heta}\,,\quad z_1=
ho_1 e^{i heta_1}\,,\quad z_2=
ho_2 e^{i heta_2}\,.$$

We have

$$z_1 \cdot z_2 =
ho_1
ho_2 e^{i(heta_1 + heta_2)}, \quad z^n =
ho^n e^{in heta},$$

for all $n \in \mathbb{N}$.

The proof follows immediately by the properties of the exponential. Let us see some applications of Propostion 5.43.

Example 5.44

Question. Compute $(-2 + 2i)^4$. **Solution.** We have two possibilities:

1. Use the binomial theorem:

$$(-2+2i)^{4} = (-2)^{4} + \binom{4}{1}(-2)^{3} \cdot 2i + \binom{4}{2}(-2)^{2} \cdot (2i)^{2} + \binom{4}{3}(-2) \cdot (2i)^{3} + (2i)^{4}$$

= 16 - 4 \cdot 8 \cdot 2i - 6 \cdot 4 \cdot 4 + 4 \cdot 2 \cdot 8i + 16
= 16 - 64i - 96 + 64i + 16 = -64.

2. A much simpler calculation is possible by using the exponential form: We know that

$$-2 + 2i = \sqrt{8}e^{i\frac{3}{4}\pi}$$

by Example 5.41. Hence

$$(-2+2i)^4 = \left(\sqrt{8}e^{i\frac{3}{4}\pi}\right)^4 = 8^2e^{i3\pi} = -64,$$

where we used that

$$e^{i3\pi} = e^{i\pi} = \cos(\pi) + i\sin(\pi) = -1$$

by 2π periodicity of $e^{i\theta}$ and Euler's identity.

Definition 5.45: Complex exponential

The complex exponential of $z = a + ib \in \mathbb{C}$ is defined as

 $e^z = e^a e^{ib}$.

The complex exponential behaves exactly as exponentials should.

Theorem 5.46

Let $z, w \in \mathbb{C}$. Then

$$e^{z+w} = e^z e^w, \quad (e^z)^w = e^{zw}.$$
 (5.6)

We still do not have the technical means to prove this Theorem. The idea is to express e^z as

$$e^z = \sum_{k=0}^{\infty} \frac{z^n}{n!},$$
 (5.7)

where the convergence is intendend in the sense of complex series. The properties at (5.6) then follow from manipulating the series on the RHS of (5.7).

Example 5.47

Question. Compute i^i . **Solution.** We know that

$$|i|=1$$
, $\arg(i)=\frac{\pi}{2}$

Hence we can write i in exponential form

$$i = |i|e^{i \arg(i)} = e^{i\frac{\pi}{2}}.$$

Therefore

$$e^{i} = \left(e^{irac{\pi}{2}}
ight)^{i} = e^{i^{2}rac{\pi}{2}} = e^{-rac{\pi}{2}} \, .$$

5.6 Fundamental Theorem of Algebra

We started the introduction to complex numbers with the following question:

Question 5.48

Is there a number $x \in \mathbb{R}$ such that

$$x^2 = -1 ? (5.8)$$

The answer is no. For this reason we introduced the complex number *i*, which satisfies

$$i^2 = -1$$
.

Therefore (5.8) has solution in \mathbb{C} , with x = i. Moreover, note that

$$(-i)^2 = (-1)^2 i^2 = -1$$
.

Hence (5.8) has two solutions in \mathbb{C} , given by

$$x_1 = i, \quad x_2 = -i.$$

It turns out that the set \mathbb{C} is so large, that not only we are able to solve (5.8), but in fact any polynomial equation.

Theorem 5.49: Fundamental theorem of algebra

Let $p_n(z)$ be a polynomial of degree *n* with complex coefficients, i.e.,

$$p_n(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0,$$

for some coefficients $a_n, \ldots, a_0 \in \mathbb{C}$ with $a_n \neq 0$. There exist

$$z_1, \ldots, z_n \in \mathbb{C}$$

solutions to the polynomial equation

$$p_n(z) = a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0 = 0.$$
(5.9)

In particular, p_n factorizes as

$$p_n(z) = a_n (z - z_1) (z - z_2) \cdots (z - z_n) .$$
(5.10)

Theorem 2.111 says that equation (5.9) admits *n* complex solutions:

- We call these solutions zeros, or also roots.
- We call the expression (5.10) a *factorization* of the polynomial p_n .

Several proofs of Theorem 2.111 exist in the literature, but they all use mathematical tools which are out of reach for now. Therefore we will not show a proof. For example one can prove Theorem 2.111 by

- Liouville's theorem (complex analysis)
- Homotopy arguments (general topology)
- Fundamental Theorem of Galois Theory (algebra)

Example 5.50

Question. Find all the complex solutions to

$$^{2} = -1$$
 (5.11)

Solution. The equation $z^2 = -1$ is equivalent to

$$p(z) = 0$$
, $p(z) := z^2 + 1$.

z

Since *p* has degree n = 2, the Fundamental Theorem of Algebra tells us that there are two solutions to (5.11). We have already seen that these two solutions are z = i and z = -i. Then *p* factorizes as

$$p(z) = z^{2} + 1 = (z - i)(z + i).$$

Example 5.51

Question. Find all the complex solutions to

$$z^4 - 1 = 0. (5.12)$$

Solution The associated polynomial equation is

p(z) = 0, $p(z) := z^4 - 1$.

Since p has degree n = 4, the Fundamental Theorem of Algebra tells us that there are 4 solutions to (5.12). Let us find such solutions. We use the well known identity

$$a^2 - b^2 = (a+b)(a-b), \quad \forall a, b \in \mathbb{R},$$

to factorize p. We get:

$$p(z) = (z^4 - 1) = (z^2 + 1)(z^2 - 1).$$

We know that

 $z^2 + 1 = 0$

has solutions $z = \pm i$. Instead

 $z^2 - 1 = 0$

has solutions $x = \pm 1$. Hence, the four solutions of (5.12) are given by

z = 1, -1, i, -i,

and p factorizes as

$$p(z) = z^4 - 1 = (z - 1)(z + 1)(z - i)(z + i).$$

Definition 5.52: Multiplicity

Suppose that the polynomial p_n factorizes as

$$p_n(z) = a_n(z-z_1)^{k_1}(z-z_2)^{k_2}\cdots(z-z_m)^{k_m}$$

with $a_n \neq 0, z_1, \dots, z_m \in \mathbb{C}$ and $k_1, \dots, k_m \in \mathbb{N}, k_i \ge 1$. In this case p_n has degree

$$n = k_1 + \ldots + k_m = \sum_{i=1}^m k_i.$$

Note that z_i is solves the equation

 $p_n(z)=0$

exactly k_i times. We call k_i the **multiplicity** of the solution z_i .

Example 5.53

The equation

$$(z-1)(z-2)^2(z+i)^3 = 0$$

has 6 solutions:

- z = 1 with multiplicity 1
- z = 2 with multiplicity 2
- z = -i with multiplicity 3

5.7 Solving polynomial equations

The non-factorized version of the polynomial

$$p(z) = (z-1)(z-2)^2(z+i)^3$$
(5.13)

from Example 5.53 is

$$p(z) = z^{6} - (5 - 3i)z^{5} + (5 - 15i)z^{4} + (11 + 23i)z^{3} - (24 + 7i)z^{2} + (12 - 8i)z + 4i$$
(5.14)

Question: How do we figure out the factorization at (5.13) if we are given p in the form at (5.13)? More in general, consider the following question:

Question 5.54

The Fundamental Theorem of Algebra states that

$$p_n(z) = 0 \tag{5.15}$$

has n complex solutions. How do we find such solutions in practice?

Answer: There is no general formula to solve (5.15) when $n \ge 5$. This is the content of the Abel-Ruffini Theorem.

Theorem 5.55: Abel-Ruffini

There is no elementary solution formula to the polynomial equation

 $p_n(z)=0\,,$

with p_n polynomial of degree n, with $n \ge 5$.

Similarly to the Fundamental Theorem of Algebra, the proof of the Abel-Ruffini Theorem is out of reach for now. A proof can be carried out, for example, using Galois Theory.

There are however explicit formulas for solving (5.15) when p_n has degree n = 2, 3, 4.

5.7.1 Quadratic polynomials

Consider polynomial equations of order n = 2, that is, equations of the form

$$az^2 + bz + c = 0. (5.16)$$

When the coefficients *a*, *b*, *c* are real, the solutions are given by the well-known quadratic formula.

Proposition 5.56: Quadratic formula

Let $a, b, c \in \mathbb{R}, a \neq 0$ and consider the equation at (5.16). Define

$$\Delta := b^2 - 4ac \in \mathbb{R}.$$

The following hold:

1. If $\Delta > 0$ then (5.16) has two distinct real solutions $z_1, z_2 \in \mathbb{R}$ given by

$$z_1 = \frac{-b - \sqrt{\Delta}}{2a}, \quad z_2 = \frac{-b + \sqrt{\Delta}}{2a}$$

2. If $\Delta = 0$ then (5.16) has one real solution $z \in \mathbb{R}$ with multiplicity 2. Such solution is given by

$$z=z_1=z_2=\frac{-b}{2a}.$$

3. If $\Delta < 0$ then (5.16) has two distinct complex solutions $z_1, z_2 \in \mathbb{C}$ given by

$$z_1 = \frac{-b - i\sqrt{-\Delta}}{2a}$$
, $z_2 = \frac{-b + i\sqrt{-\Delta}}{2a}$

where $\sqrt{-\Delta} \in \mathbb{R}$, since $-\Delta > 0$.

In all cases, the polynomial at (5.16) factorizes as

$$az^{2} + bz + c = a(z - z_{1})(z - z_{2}).$$

Example 5.57

Question. Solve the following equations:

1. $3z^2 - 6z + 2 = 0$ 2. $4z^2 - 8z + 4 = 0$ 3. $z^2 + 2z + 3 = 0$

Solution.

1. We have that

$$\Delta = (-6)^2 - 4 \cdot 3 \cdot 2 = 12 > 0$$

Therefore the equation has two distinct real solutions, given by

$$z = \frac{-(-6) \pm \sqrt{12}}{2 \cdot 3} = \frac{6 \pm \sqrt{12}}{6} = 1 \pm \frac{\sqrt{3}}{3}$$

In particular we have the factorization

$$3z^{2} - 6z + 2 = 3\left(z - 1 - \frac{\sqrt{3}}{3}\right)\left(z - 1 + \frac{\sqrt{3}}{3}\right).$$

2. We have that

$$\Delta = (-8)^2 - 4 \cdot 4 \cdot 4 = 0.$$

Therefore there exists one solution with multiplicity 2. This is given by

$$z = \frac{-(-8)}{2 \cdot 4} = 1$$
.

In particular we have the factorization

$$4z^2 - 8x + 4 = 4(z - 1)^2.$$

3. We have

$$\Delta = 2^2 - 4 \cdot 1 \cdot 3 = -8 < 0 \,.$$

Therefore there are two complex solutions given by

$$z = \frac{-2 \pm i\sqrt{8}}{2 \cdot 1} = -1 \pm i\sqrt{2}.$$

In particular we have the factorization

$$z^{2} + 2z + 3 = (z + 1 - i\sqrt{2})(z + 1 + i\sqrt{2}).$$

So far we have considered the polynomial equation

$$az^2 + bz + c = 0, (5.17)$$

for $a, b, c \in \mathbb{R}$ and $a \neq 0$.

Question 5.58

What if $a, b, c \in \mathbb{C}$?

If $a, b, c \in \mathbb{C}$ then we might have

$$\Delta := b^2 - 4ac \in \mathbb{C}.$$

Therefore it is not clear how to compute

 $\sqrt{\Delta}$.

However, we can still use the quadratic equation to solve (5.17), as outlined in the next Proposition.

Proposition 5.59: Quadratic formula with complex coefficients

Let $a, b, c \in \mathbb{C}$, $a \neq 0$. The two solutions to

$$az^2 + bz + c = 0$$

are given by

$$z_1 = \frac{-b + S_1}{2a}, \quad z_2 = \frac{-b + S_2}{2a},$$

where S_1 and S_2 are the two solutions to

$$z^2 = \Delta, \quad \Delta := b^2 - 4ac$$

Remark 5.60

We want to check that the new formulas

$$z_1 = \frac{-b + S_1}{2a}$$
, $z_2 = \frac{-b + S_2}{2a}$, $S_1^2 = S_2^2 = \Delta$,

agree with the old formulas given in Proposition 2.116, when $a, b, c \in \mathbb{R}$. To this end, note that

 $\Delta \in \mathbb{R}$,

when $a, b, c \in \mathbb{R}$. In this case the equation

$$z^2 = \Delta$$

has the following solutions:

- If $\Delta > 0$ there are two real solutions

$$S_1 = -\sqrt{\Delta}, \quad S_2 = \sqrt{\Delta}.$$

• If $\Delta = 0$ then 0 is the only solution with multiplicity 2. Hence

$$S_1 = S_2 = 0$$
.

• If $\Delta < 0$, there are two complex solutions

$$S_1 = -i\sqrt{-\Delta}, \quad S_2 = i\sqrt{-\Delta}.$$

Therefore the solutions

$$z_1 = \frac{-b + S_1}{2a}, \quad z_2 = \frac{-b + S_2}{2a}$$

given in Proposition 2.116 coincide with the ones given in Proposition 2.118.

Example 5.61

Question Find all the solutions to

$$\frac{1}{2}z^2 - (3+i)z + (4-i) = 0.$$
(5.18)

Solution. We have

$$\Delta = (-(3+i))^2 - 4 \cdot \frac{1}{2} \cdot (4-i)$$

= 8 + 6i - 8 + 2i
= 8i.

Therefore $\Delta \in \mathbb{C}$. We have to find solutions S_1 and S_2 to the equation

$$z^2 = \Delta = 8i. \tag{5.19}$$

We look for solutions of the form z = a + ib. Then we must have that

$$z^{2} = (a + ib)^{2} = a^{2} - b^{2} + 2abi = 8i$$
.

Thus

$$a^2 - b^2 = 0$$
, $2ab = 8$.

From the first equation we conclude that |a| = |b|. From the second equation we have that ab = 4, and therefore *a* and *b* must have the same sign. Hence a = b, and

 $2ab = 8 \implies a = b = \pm 2$.

From this we conclude that the solutions to (5.19) are

 $S_1 = 2 + 2i$, $S_2 = -2 - 2i$.

Hence the solutions to (5.18) are

$$z_1 = \frac{3+i+S_1}{2\cdot\frac{1}{2}} = 3+i+S_1$$
$$= 3+i+2+2i = 5+3i,$$

and

$$z_2 = \frac{3+i+S_2}{2\cdot\frac{1}{2}} = 3+i+S_2$$
$$= 3+i-2-2i = 1-i.$$

In particular, the given polynomial factorizes as

$$\frac{1}{2}z^2 - (3+i)z + (4-i) = \frac{1}{2}(z-z_1)(z-z_2)$$
$$= \frac{1}{2}(z-5-3i)(z-1+i).$$

In the above example it was a bit laborious to compute S_1 and S_2 . In Section 5.8 and Section 5.9, we will explore a more general method to solve problems of the form

$$z^n = \Delta$$
.

5.7.2 Higher order polynomials

Consider now polynomial equations

 $p_n(z)=0\,,$

with p_n of degree n = 3, 4. Although general solution formulas exist for these cases, they are exceedingly lengthy, making them impractical for manual calculations. For example, Figure 5.8 shows the solution formula to the quartic equation

$$z^4 + az^3 + bz^2 + cz + d = 0$$

when $a, b, c, d \in \mathbb{R}$.

Figure 5.8: Quartic formula to solve $z^4 + az^3 + bz^2 + cz + d = 0$. Image from Wikipedia.

A more productive use of time is learning how to perform long polynomial division. A quick tutorial is available here: Polynomial_division.pdf.

Example 5.62

Question. Divide $6z^3 + 5z^2 - 7$ by $3z^2 - 2z - 1$. **Solution.** Using polynomial division, see Figure 5.9, we obtain

$$6z^{3} + 5z^{2} - 7 = (3z^{2} - 2z - 1)(2z + 3) + (8z - 4).$$

$$\begin{array}{r} 2z+3\\3z^2-2z-1\overbrace{) \ \ 6z^3+5z^2\ \ -7}\\ -6z^3+4z^2+2z\\ \hline 9z^2+2z-7\\ -9z^2+6z+3\\ \hline 8z-4\end{array}$$

Figure 5.9: Example of polynomial long division between $6z^3 + 5z^2 - 7$ and $3z^2 - 2z - 1$.

Sometimes, it is possible to solve equations of degree higher than 2, in case it is obvious from inspection that a certain number is a solution, e.g., when z = -1, 0, 1 is a solution.

Example 5.63

Question. Consider the equation

$$z^3 - 7z^2 + 6z = 0.$$

- 1. Check whether z = 0, 1, -1 are solutions.
- 2. Using your answer from Point 1, and polynomial division, find all the solutions.

Solution.

- 1. By direct inspection we see that z = 0 and z = 1 are solutions.
- 2. Since z = 0 is a solution, we can factorize

$$z^3 - 7z^2 + 6z = z \left(z^2 - 7z + 6 \right) \, .$$

We could now use the quadratic formula on the term $z^2 - 7z + 6$ to find the remaining two roots. However, we have already observed that z = 1 is a solution. Therefore z - 1 divides $z^2 - 7z + 6$. Using polynomial long division, see Figure 5.10, we find that

$$\frac{z^2 - 7z + 6}{z - 1} = z - 6 \,.$$

Therefore the last solution is z = 6, and

$$z^3 - 7z^2 + 6z = z(z - 1)(z - 6)$$

$$\begin{array}{r} z-6 \\ z-1 \overline{\smash{\big)} \ z^2 - 7z + 6} \\ \underline{-z^2 + z} \\ -6z + 6 \\ \underline{-6z - 6} \\ 0 \end{array}$$

Figure 5.10: Polynomial long division between $z^2 - 7z + 6$ and z - 1.

Example 5.64

Question. Find all the complex solutions to

$$z^3 - 7z + 6 = 0.$$

Solution. It is easy to see z = 1 is a solution. This means that z - 1 divides $z^3 - 7z + 6$. By using polynomial long division, see Figure 5.11, we compute that

$$\frac{z^3 - 7z + 6}{z - 1} = z^2 + z - 6$$

We are now left to solve

 $z^2 + z - 6 = 0.$

Using the quadratic formula, we see that the above is solved by z = 2 and z = -3. Therefore the given polynomial factorizes as

$$z^{3} - 7z + 6 = (z - 1)(z - 2)(z + 3).$$

$$\begin{array}{r}z^2 + z - 6\\z - 1 \overline{\smash{\big)} z^3} & -7z + 6\\ -z^3 + z^2\\ \hline z^2 - 7z\\ -z^2 + z\\ -6z + 6\\ \hline 6z - 6\\ \hline 0\end{array}$$

Figure 5.11: Polynomial long division between $z^3 - 7z + 6$ and z - 1.

Example 5.65

Question. Consider the equation

$$z^{3} + 5z^{2} + (5 - i)z + (4 - 4i) = 0.$$

- 1. Check whether $z = \pm i$ and $z = 0, \pm 1$ are solutions.
- 2. By using polynomial division with complex coefficients, find all the solutions.

Solution.

- 1. By direct inspection, we see that z = i is a solution.
- 2. Since z = i is a solution, we know that z i divides $z^3 + 5z^2 + (5 i)z + (4 4i)$. We now perform polynomial division by using the same method employed to divide polynomials with real coefficients: We just need to be mindful of the fact that coefficients are now complex, and thus, addition and multiplication have to be carried out in \mathbb{C} . Using polynomial long division, see Figure 5.12, we compute that

$$z^{3} + 5z^{2} + (5-i)z + (4-4i) = (z-i)(z^{2} + (5+i)z + (4-4i)).$$

We are now left to solve the second order equation

$$z^2 + (5+i)z + (4-4i) = 0.$$

We compute that

 $\Delta = b^2 - 4ac = (5+1)^2 - 4(4-4i) = 4(5+4i).$

We now need to solve

$$z^2 = \Delta$$
.

We look for a solution in the form of z = a + ib.

$$z^{2} = (a + ib)^{2} = a^{2} - b^{2} + 2abi = 8i$$
.

Thus

$$a^2 - b^2 = 0$$
, $2ab = 8$.

From the first equation we conclude that |a| = |b|. From the second equation we have that ab = 4, and therefore *a* and *b* must have the same sign. Hence a = b, and

$$2ab = 8 \implies a = b = \pm 2$$

From this we conclude that the solutions to (5.19) are

$$S_1 = 2 + 2i$$
, $S_2 = -2 - 2i$.

Hence the solutions to (5.18) are

$$z_1 = \frac{3+i+S_1}{2 \cdot \frac{1}{2}} = 3+i+S_1$$
$$= 3+i+2+2i = 5+3i,$$

and

$$z_2 = \frac{3+i+S_2}{2\cdot\frac{1}{2}} = 3+i+S_2$$
$$= 3+i-2-2i = 1-i.$$

In particular, the given polynomial factorizes as

$$\frac{1}{2}z^2 - (3+i)z + (4-i) = \frac{1}{2}(z-z_1)(z-z_2)$$
$$= \frac{1}{2}(z-5-3i)(z-1+i)$$

$$\frac{2^{2} + (5+i)2 + (4+4i)}{(2-i)}$$

$$z-i \qquad \boxed{2^{3} + 5z^{2} + (5-i)2 + (4-4i)}$$

$$\frac{z^{3} - iz^{2}}{\sqrt{(5+i)z^{2} + (5-i)z + (4-4i)}}$$

$$\frac{(5+i)z^{2} - i(5+i)z}{\sqrt{(4+4i)z + (4-4i)}}$$

$$\frac{(4+4i)z - i(4+4i)}{\sqrt{(4+4i)z - i(4+4i)}}$$

Figure 5.12: Polynomial long division between $z^3 + 5z^2 + (5 - i)z + (4 - 4i)$ and z - i.

5.8 Roots of unity

Problem

Let $n \in \mathbb{N}$. We want to find all complex solutions to

$$z^n = 1$$
. (5.20)

Note that z = 1 is always a solution to (5.20) if *n* is even. In such case also z = -1 is a solution. If we were only looking for solutions in \mathbb{R} , these two would be the only solutions.

However, the Fundamental Theorem of Algebra, see Theorem 2.111, tells us that there are n complex solutions to (5.20).

Question 5.66

Is there a way to find all *n* solutions?

Example 5.67

We have seen in Example 5.51 that the solutions to

$$z^4 = 1$$

are

$$z=-1,1,i,-i.$$

However, we could only compute these solutions thanks to the clever factorization

 $z^4 - 1 = (z^2 + 1)(z^2 - 1) = (z + i)(z - i)(z + 1)(z - 1).$

It is not clear if and how this trick can be generalized to solve

 $z^{n} = 1$

for arbitrary $n \in \mathbb{N}$.

The trick to find all n solutions to (5.20) is to use the exponential form.

Theorem 5.68

Let $n \in \mathbb{N}$ and consider the equation

$$z^n = 1$$
. (5.21)

All the n solutions to (5.21) are given by

$$z_k = \exp\left(i\frac{2\pi k}{n}\right), \quad k = 0, \dots, n-1,$$

where $\exp(x)$ denotes e^x .

Proof

Rewrite 1 in exponential form:

$$1 = |1|e^{i \arg(1)} = e^{i2\pi k}, \quad k \in \mathbb{Z}.$$

Therefore (5.21) is equivalent to

$$z^n = e^{i2\pi k} \, .$$

By the properties of the exponential, we see that the above is solved by

$$z_k = \exp\left(i\frac{2\pi k}{n}\right), \quad k \in \mathbb{Z}.$$

By choosing k = 0, ..., n - 1 we obtain *n* different solutions.

Definition 5.69

The n solutions to

$$z^{n} = 1$$

are called the **roots of unity**.

Example 5.70

Question. Find all the solutions to

$$z^4 = 1$$
.

Solution. The 4 solutions are given by

$$z_k = \exp\left(i\frac{2\pi k}{4}\right) = \exp\left(i\frac{\pi k}{2}\right),$$

for k = 0, 1, 2, 3. We compute:

$$\begin{aligned} z_0 &= e^{i0} = 1 \,, & z_1 &= e^{i\frac{\pi}{2}} = i \,, \\ z_2 &= e^{i\pi} = -1 \,, & z_3 &= e^{i\frac{3\pi}{2}} = -i \,. \end{aligned}$$

Note that for k = 4 we would again get the solution $z = e^{i2\pi} = 1$.

Example 5.71

Question. Find all the solutions to

 $z^3 = 1$.

Solution. The 3 solutions are given by

$$z_k = \exp\left(i\frac{2\pi k}{3}\right)\,,$$

for k = 0, 1, 2. We compute:

$$z_0 = e^{i0} = 1$$
, $z_1 = e^{i\frac{2\pi}{3}}$, $z_2 = e^{i\frac{4\pi}{3}}$.

We can write z_1 and z_2 in cartesian form:

$$z_1 = e^{i\frac{2\pi}{3}} = \cos\left(\frac{2\pi}{3}\right) + i\sin\left(\frac{2\pi}{3}\right) = -\frac{1}{2} + \frac{\sqrt{3}}{2}i$$

and

$$z_2 = e^{i\frac{4\pi}{3}} = \cos\left(\frac{4\pi}{3}\right) + i\sin\left(\frac{4\pi}{3}\right) = -\frac{1}{2} - \frac{\sqrt{3}}{2}i.$$

5.9 Roots in \mathbb{C}

Let $n \in \mathbb{N}$ and $c \in \mathbb{C}$. We want to give a meaning to

 $\sqrt[n]{c}$.

This means, we want to find all complex solutions to

$$z^n = c$$
.

The Fundamental Theorem of Algebra ensures that the above equation has n complex solutions. To find these solutions, we pass to the exponential form.

Theorem 5.72

Let $n \in \mathbb{N}$, $c \in \mathbb{C}$ and consider the equation

$$z^n = c \,. \tag{5.22}$$

All the n solutions to (5.22) are given by

$$z_k = \sqrt[n]{|c|} \exp\left(i\frac{\theta + 2\pi k}{n}\right), \quad k = 0, \dots, n-1,$$

where $\sqrt[n]{|c|}$ is the *n*-th root of the real number |c|, and $\theta = \arg(c)$.

Proof

Write c in exponential form:

$$c = |c|e^{i\theta} = |c|e^{i(\theta + 2\pi k)}, \quad k \in \mathbb{Z},$$

where $\theta = \arg(c)$. Therefore (5.22) is equivalent to

$$z^n = |c|e^{i(\theta + 2\pi k)}$$

By the properties of the exponential, we see that the above is solved by

$$z_k = \sqrt[n]{|c|} \exp\left(i\frac{\theta + 2\pi k}{n}\right), \quad k \in \mathbb{Z}.$$

By choosing k = 0, ..., n - 1 we obtain *n* different solutions.

Warning: The n-th root function is multi-valued

The above result and the Fundamental Theorem of Algebra tell us that

$$z_0, ..., z_{n-1}$$

are *n* solutions to

 $z^n = c$.

This means that the complex number *c* has multiple *n*-th roots, exaclty *n*. In particular, the *n*-th root function in \mathbb{C} is multi-valued:

$$\sqrt[n]{c} = \{z_0, \dots, z_{n-1}\}.$$

Example 5.73

Question. Find all the $z \in \mathbb{C}$ such that

 $z^5 = -32$.

Solution. Let c = -32. We have

$$|c| = |-32| = 32 = 2^5$$
, $\theta = \arg(-32) = \pi$.

The 5 solutions are given by

$$z_k = \left(2^5\right)^{\frac{1}{5}} \exp\left(i\pi \frac{1+2k}{5}\right), \quad k \in \mathbb{Z},$$

for k = 0, 1, 2, 3, 4. We get

$$z_{0} = 2e^{i\frac{\pi}{5}} \qquad z_{1} = 2e^{i\frac{3\pi}{5}}$$
$$z_{2} = 2e^{i\pi} = -2 \qquad z_{3} = 2e^{i\frac{7\pi}{5}}$$
$$z_{4} = 2e^{i\frac{9\pi}{5}}$$

Example 5.74

Question. Find all the $z \in \mathbb{C}$ such that

$$z^4 = 9\left(\cos\left(\frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{3}\right)\right).$$

Solution. Set

$$c := 9\left(\cos\left(\frac{\pi}{3}\right) + i\sin\left(\frac{\pi}{3}\right)\right).$$

The complex number c is already in the trigonometric form, so that we can immediately obtain

$$|c| = 9$$
, $\theta = \arg(c) = \frac{\pi}{3}$.

The 4 solutions are given by

$$z_k = \sqrt[4]{9} \exp\left(i\frac{\pi/3 + 2\pi k}{4}\right)$$
$$= \sqrt{3} \exp\left(i\pi \frac{1+6k}{12}\right)$$

for k = 0, 1, 2, 3. We compute

$$z_0 = \sqrt{3}e^{i\pi\frac{1}{12}} \qquad z_1 = \sqrt{3}e^{i\pi\frac{7}{12}}$$
$$z_2 = \sqrt{3}e^{i\pi\frac{13}{12}} \qquad z_3 = \sqrt{3}e^{i\pi\frac{19}{12}}$$

6 Sequences in \mathbb{R}

A sequence is an infinite list of real numbers. For example, the following are sequences:

- (1, 2, 3, 4, ...)
- (-1, 1, -1, 1, ...)
- $\left(1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \ldots\right)$

Remark 6.1

• The order of elements in a sequence matters.

For example

- $(1,2,3,4,5,6,\ldots) \neq (2,1,4,3,6,5,\ldots)$
- A sequence is not a set.

For example

$$\{-1, 1, -1, 1, -1, 1, ...\} = \{-1, 1\}$$

but we cannot make a similar statement for the sequence

(-1, 1, -1, 1, -1, 1, ...).

• The above notation is ambiguous.

For example the sequence

$$(1, 2, 3, 4, \ldots)$$

can continue as

$$(1, 2, 3, 4, 1, 2, 3, 4, 1, 2, 3, 4, ...)$$
.

• In the sequence

$$\left(1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \ldots\right)$$

the elements get smaller and smaller, and closer and closer to 0. We say that this sequence converges to 0, or has 0 as a limit.

We would like to make the notions of sequence and convergence more precise.

6.1 Definition of sequence

We start with the definition of sequence of Real numbers.

Definition 6.2: Sequence of Real numbers

A sequence a in \mathbb{R} is a function

 $a: \mathbb{N} \to \mathbb{R}$.

For $n \in \mathbb{N}$, we denote the *n*-th element of the sequence *a* by

 $a_n = a(n)$

and write the sequence as

 $(a_n)_{n\in\mathbb{N}}$.

Notation 6.3

We will sometimes omit the subscript $n \in \mathbb{N}$ and simply write

 (a_n) .

In certain situations, we will also write

 $(a_n)_{n=1}^{\infty}$.

Example 6.4

• In general $(a_n)_{n \in \mathbb{N}}$ is the sequence

 (a_1, a_2, a_3, \ldots) .

• Consider the function

$$a: \mathbb{N} \to \mathbb{N}, \quad n \mapsto 2n.$$

This is also a sequence of real numbers. It can be written as

 $(2n)_{n\in\mathbb{N}}$

and it represents the sequence of even numbers

(2, 4, 6, 8, 10, ...).

• Let

 $a_n = (-1)^n$

Then (a_n) is the sequence

(-1, 1, -1, 1, -1, 1, ...).

• $\left(\frac{1}{n}\right)_{n\in\mathbb{N}}$ is the sequence

$$\left(1,\frac{1}{2},\frac{1}{4},\frac{1}{5},...\right)\,.$$

6.2 Convergent sequences

We have notice that the sequence $\left(\frac{1}{n}\right)_{n \in \mathbb{N}}$ gets close to 0 as *n* gets large. We would like to say that a_n converges to 0 as *n* tends to infinity.

To make this precise, we first have to say what it means for two numbers to be *close*. For this we use the notion of absolute value, and say that:

- *x* and *y* are close if |x y| is small.
- |x y| is called the distance between x and y
- For *x* to be close to 0, we need that |x 0| = |x| is small.

Saying that |x| is *small* is not very precise. Let us now give the formal definition of convergent sequence.

Definition 6.5: Convergent sequence

The real sequence (a_n) **converges** to *a*, or equivalently has limit *a*, denoted by

$$\lim_{n\to\infty}a_n=a$$

if for all $\varepsilon \in \mathbb{R}, \varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \in \mathbb{N}, n \ge N$ it holds that

$$|a_n-a|<\varepsilon.$$

Using quantifiers, we can write this as

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |a_n - a| < \varepsilon.$$

The sequence $(a_n)_{n \in \mathbb{N}}$ is **convergent** if it admits limit.

Notation 6.6

We will often write

in place of

 $a_n \rightarrow a$

$$\lim_{n\to\infty}a_n=a$$

Remark 6.7

- Informally, Definition 2.129 says that, no matter how small we choose ε (as long as it is strictly positive), we always have that a_n has a distance to a of less than or equal to ε from a certain point onwards (i.e., from N onward). The sequence (a_n) may fluctuate wildly in the beginning, but from N onward it should stay within a distance of ε of a.
- In general *N* depends on ε . If ε is chosen smaller, we might have to take *N* larger: this means we need to wait longer before the sequence stays within a distance ε from *a*.

We now prove that the sequence

$$a_n = \frac{1}{n}$$

converges to 0, according to Definition 2.129.

Theorem 6.8

The sequence $\left(\frac{1}{n}\right)_{n \in \mathbb{N}}$ converges to 0, i.e.,

$$\lim_{n\to\infty}\frac{1}{n}=0\,.$$

We give two proofs of the above theorem:

- Long proof, with all the details.
- Short proof, with less details, but still acceptable.

Proof: Proof of Theorem 6.8 (Long version)

We have to show that

$$\lim_{n\to\infty}\frac{1}{n}=0$$

which by definition is equivalent to showing that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, \left| \frac{1}{n} - 0 \right| < \varepsilon.$$
(6.1)

Let $\varepsilon \in \mathbb{R}$ with $\varepsilon > 0$. Choose $N \in \mathbb{N}$ such that

$$N > \frac{1}{\varepsilon}$$

Such natural number N exists thanks to the Archimedean property. The above implies

$$\frac{1}{N} < \varepsilon \,, \tag{6.2}$$

Let $n \in \mathbb{N}$ with $n \ge N$. By (6.2) we have

$$\frac{1}{n} \leq \frac{1}{N} < \varepsilon$$

From this we deduce

$$\left|\frac{1}{n} - 0\right| = \frac{1}{n} \le \frac{1}{N} < \varepsilon$$

Since $n \in \mathbb{N}$, $n \ge N$ was arbitrary, we have proven that

$$\left|\frac{1}{n} - 0\right| < \varepsilon, \quad \forall n \ge N.$$
(6.3)

Condition (6.3) holds for all $\varepsilon > 0$, for the choice of $N \in \mathbb{N}$ such that

$$N > \frac{1}{\varepsilon}$$
.

We have hence shown (6.1), and the proof is concluded.

As the above proof is quite long and includes lots of details, it is acceptable to shorten it. For example:

- We skip some intermediate steps.
- We do not mention the Archimedean property.
- We leave out the conclusion when it is obvious that the statement has been proven.

Proof: Proof of Theorem 6.8 (Short version)

We have to show that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, \left| \frac{1}{n} - 0 \right| < \varepsilon.$$

Let $\varepsilon > 0$. Choose $N \in \mathbb{N}$ such that

$$N > \frac{1}{\varepsilon}$$
.

Let $n \ge N$. Then

$$\left|\frac{1}{n}-0\right|=\frac{1}{n}\leq\frac{1}{N}<\varepsilon\,.$$

In Theorem 6.8 we showed that

$$\lim_{n\to\infty} \frac{1}{n} = 0.$$

We can generalise this statement to prove that

$$\lim_{n \to \infty} \frac{1}{n^p} = 0$$

for any p > 0 fixed.

Theorem 6.9

For all p > 0, the sequence $\left(\frac{1}{n^p}\right)_{n \in \mathbb{N}}$ converges to 0, i.e.,

$$\lim_{n \to \infty} \frac{1}{n^p} = 0$$

Proof

Let p > 0. We have to show that

$$\forall \varepsilon > 0 \,, \, \exists \, N \in \mathbb{N} \, \text{ s.t. } \, \forall \, n \geq N \,, \, \left| \frac{1}{n^p} - 0 \right| < \varepsilon \,.$$

Let $\varepsilon > 0$. Choose $N \in \mathbb{N}$ such that

$$N > \frac{1}{\varepsilon^{1/p}} \,. \tag{6.4}$$

Let $n \ge N$. Since p > 0, we have $n^p \ge N^p$, which implies

$$\frac{1}{n^p} \le \frac{1}{N^p}$$

By (6.4) we deduce

$$\frac{1}{N^p} < \varepsilon \,.$$

Then

$$\left|\frac{1}{n^p} - 0\right| = \frac{1}{n^p} \le \frac{1}{N^p} < \varepsilon \,.$$

Question 6.10

Why did we choose $N \in \mathbb{N}$ such that

$$N > \frac{1}{\varepsilon^p}$$

in the above proof?

The answer is: because it works. Finding a number N that makes the proof work requires some *rough work*: Specifically, such rough work consists in finding $N \in \mathbb{N}$ such that the inequality

 $|a_N - a| < \varepsilon$

is satisfied.

Important

Any *rough work* required to prove convergence must be shown before the formal proof (in assignments).

Example 6.11

Question. Using the definition of convergence, prove that

$$\lim_{n \to \infty} \frac{n}{2n+3} = \frac{1}{2}.$$

Solution.

1. *Rough Work:* Let $\varepsilon > 0$. We want to find $N \in \mathbb{N}$ such that

$$\left|\frac{n}{2n+3}-\frac{1}{2}\right|<\varepsilon\,,\quad\forall\,n\geq N\,.$$

To this end, we compute:

$$\left|\frac{n}{2n+3} - \frac{1}{2}\right| = \left|\frac{-3}{4n+6}\right| = \frac{3}{4n+6}.$$

Therefore

$$\left|\frac{n}{2n+3} - \frac{1}{2}\right| < \varepsilon \quad \iff \quad \frac{3}{4n+6} < \varepsilon$$
$$\iff \quad n > \frac{3}{4\varepsilon} - \frac{6}{4}$$

Looking at the above equivalences, it is clear that $N \in \mathbb{N}$ has to be chosen so that

$$N > \frac{3}{4\varepsilon} - \frac{6}{4} \,. \tag{6.5}$$

2. Formal Proof: We have to show that

$$\forall \varepsilon > 0 , \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N , \left| \frac{n}{2n+3} - \frac{1}{2} \right| < \varepsilon$$

Let $\varepsilon > 0$. Choose $N \in \mathbb{N}$ such that (6.5) holds. By the rough work shown above, inequality (6.5) is equivalent to

$$\frac{3}{4N+6} < \varepsilon$$

Let $n \ge N$. Then

$$\left|\frac{n}{2n+3} - \frac{1}{2}\right| = \frac{3}{4n+6} \le \frac{3}{4N+6} < \varepsilon,$$

where in the third line we used that $n \ge N$.

We conclude by showing that constant sequences always converge.

Theorem 6.12

Let $c \in \mathbb{R}$ and define the constant sequence

 $a_n := c$, $\forall n \in \mathbb{N}$.

We have that

$$\lim_{n\to\infty} a_n = c \, .$$

Proof

We have to prove that

$$\forall \varepsilon > 0, \ \exists N \in \mathbb{N} \ \text{s.t.} \ \forall n \ge N, \ |a_n - c| < \varepsilon.$$
(6.6)

Let $\varepsilon > 0$. We have

 $|a_n - c| = |c - c| = 0 < \varepsilon, \quad \forall n \in \mathbb{N}.$

Therefore we can choose N = 1 and (6.6) is satisfied.

6.3 Divergent sequences

The opposite of convergent sequences are divergent sequences.

Definition 6.13: Divergent sequence

We say that a sequence $(a_n)_{n \in \mathbb{N}}$ in \mathbb{R} is **divergent** if it is not convergent.

Remark 6.14

Proving that a sequence (a_n) is divergent is more complicated than showing it is convergent: To show that (a_n) is divergent, we need to show that (a_n) cannot converge to *a* for any $a \in \mathbb{R}$. In other words, we have to show that there does not exist an $a \in \mathbb{R}$ such that

$$\lim_{n\to\infty} a_n = a \, .$$

Using quantifiers, this means

$$\exists a \in \mathbb{R} \text{ s.t. } \forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |a_n - a| < \varepsilon.$$

The above is equivalent to showing that

 $\forall \, a \in \mathbb{R} \,, \, \exists \, \varepsilon > 0 \; \text{ s.t. } \; \forall \, N \in \mathbb{N} \,, \, \exists \, n \geq N \; \text{ s.t. } \; |a_n - a| \geq \varepsilon \,.$
Theorem 6.15

Let (a_n) be the sequence defined by

 $a_n=(-1)^n\,.$

Then (a_n) does not converge.

Proof

To prove that (a_n) does not converge, we have to show that

$$\forall a \in \mathbb{R}, \exists \varepsilon > 0 \text{ s.t. } \forall N \in \mathbb{N}, \exists n \ge N \text{ s.t. } |a_n - a| \ge \varepsilon.$$

Let $a \in \mathbb{R}$. Choose

$$\varepsilon = \frac{1}{2}$$
.

Let $N \in \mathbb{N}$. We distinguish two cases:

• $a \ge 0$: Choose n = 2N + 1. Note that $n \ge N$. Then

$$\begin{aligned} |a_n - a| &= |a_{2N+1} - a| \\ &= |(-1)^{2N+1} - a| \\ &= |-1 - a| \\ &= 1 + a \\ &\ge 1 \\ &> \frac{1}{2} = \varepsilon \,, \end{aligned}$$

where we used that $a \ge 0$, and therefore

$$|-1-a| = 1+a \ge 1$$
.

• a < 0: Choose n = 2N. Note that $n \ge N$. Then

$$\begin{split} |a_n - a| &= |a_{2N} - a| \\ &= |(-1)^{2N} - a| \\ &= |1 - a| \\ &= 1 - a \\ &> 1 \\ &> \frac{1}{2} = \varepsilon \,, \end{split}$$

where we used that a < 0, and therefore

|1-a| = 1-a > 1.

We propose an alternative proof by contradiction.

Proof: Alternative proof to Theorem 6.15

Suppose by contradiction that $a_n \rightarrow a$ for some $a \in \mathbb{R}$. Let

$$\varepsilon := \frac{1}{2}$$
.

Since $a_n \to a$, there exists $N \in \mathbb{N}$ such that

$$|a_n-a|<\varepsilon=\frac{1}{3} \quad \forall\,n\geq N\,.$$

If we take n = 2N, then $n \ge N$ and

$$|a_{2N} - a| = |1 - a| < \frac{1}{2}$$

If we take n = 2N + 1, then $n \ge N$ and

$$|a_{2N+1}-a| = |-1-a| < \frac{1}{2}.$$

Therefore

$$2 = |(1-a) - (-1-a)|$$

$$\leq |1-a| + |-1-a|$$

$$< \frac{1}{2} + \frac{1}{2} = 1,$$

which is a contradiction. Hence (a_n) is divergent.

6.4 Uniqueness of limit

In Definition 2.129 of convergence, we used the notation

$$\lim_{n\to\infty}a_n=a\,.$$

The above notation makes sense only if the limit is unique, that is, if we do not have that

$$\lim_{n\to\infty}a_n=b\,,$$

for some

 $a \neq b$.

In the next theorem we will show that the limit is unique, if it exists.

Theorem 6.16: Uniqueness of limit

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence. Suppose that

$$\lim_{n \to \infty} a_n = a , \quad \lim_{n \to \infty} a_n = b .$$

Then a = b.

Proof

Assume that,

$$\lim_{n\to\infty}a_n=a\,,\quad \lim_{n\to\infty}a_n=b\,.$$

Suppose by contradiction that

 $a \neq b$.

Choose

$$\varepsilon := \frac{1}{2} \left| a - b \right|.$$

Therefore $\varepsilon > 0$, since |a - b| > 0. By the convergence $a_n \to a$,

 $\exists N_1 \in \mathbb{N} \text{ s.t. } \forall n \geq N_1 \text{ , } |a_n - a| < \varepsilon \text{ .}$

By the convergence $a_n \rightarrow b$,

 $\exists N_2 \in \mathbb{N} \text{ s.t. } \forall n \ge N_2 \text{ , } |a_n - b| < \varepsilon \text{ .}$

Define

 $N := \max\{N_1, N_2\}.$

Choose an $n \in \mathbb{N}$ such that $n \ge N$. In particular

 $n \geq N_1$, $n \geq N_2$.

Then

$$2\varepsilon = |a - b|$$

= $|a - a_n + a_n - b|$
 $\leq |a - a_n| + |a_n - b|$
 $< \varepsilon + \varepsilon$
= 2ε .

where we used the triangle inequality in the first inequality. Hence $2\varepsilon < 2\varepsilon$, which gives a contradiction.

Example 6.17

Question. Prove that

$$\lim_{n \to \infty} \frac{n^2 - 1}{2n^2 - 3} = \frac{1}{2}$$

Solution. According to Theorem 2.134, it suffices to show that the sequence

$$\left(\frac{n^2-1}{2n^2-3}\right)_{n\in\mathbb{N}}$$

converges to $\frac{1}{2}$, since then $\frac{1}{2}$ can be the only limit.

1. *Rough Work:* Let $\varepsilon > 0$. We want to find $N \in \mathbb{N}$ such that

$$\left|\frac{n^2-1}{2n^2-3}-\frac{1}{2}\right|<\varepsilon\,,\quad\forall\,n\geq N\,.$$

To this end, we compute:

$$\left|\frac{n^2 - 1}{2n^2 - 3} - \frac{1}{2}\right| = \left|\frac{2(n^2 - 1) - (2n^2 - 3)}{2(2n^2 - 3)}\right|$$
$$= \left|\frac{1}{4n^2 - 6}\right|$$
$$= \frac{1}{4n^2 - 6}$$
$$= \frac{1}{3n^2 + n^2 - 6}$$
$$\leq \frac{1}{3n^2}$$

which holds if $n \ge 3$, since in this case $n^2 - 6 \ge 0$. Therefore

$$\begin{vmatrix} \frac{n^2 - 1}{2n^2 - 3} - \frac{1}{2} \end{vmatrix} < \varepsilon \qquad \longleftarrow \qquad \frac{1}{3n^2} < \varepsilon$$
$$\iff \qquad 3n^2 > \frac{1}{\varepsilon}$$
$$\iff \qquad n^2 > \frac{1}{3\varepsilon}$$
$$\iff \qquad n > \frac{1}{\sqrt{3\varepsilon}}$$

Looking at the above implications, it is clear that $N \in \mathbb{N}$ has to be chosen so that

$$N > \frac{1}{\sqrt{3\varepsilon}} \,.$$

Moreover we need to recall that N has to satisfy

 $N \ge 3$

for the estimates to hold.

2. Formal Proof: We have to show that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, \left| \frac{n^2 - 1}{2n^2 - 3} - \frac{1}{2} \right| < \varepsilon.$$

Let
$$\varepsilon > 0$$
. Choose $N \in \mathbb{N}$ such that

$$N > \max\left\{\frac{1}{\sqrt{3\varepsilon}}, 3\right\}.$$

Let $n \ge N$. Then

$$\left| \frac{n^2 - 1}{2n^2 - 3} - \frac{1}{2} \right| = \frac{1}{4n^2 - 6}$$
$$= \frac{1}{3n^2 + n^2 - 6}$$
$$\leq \frac{1}{3n^2}$$
$$\leq \frac{1}{3N^2}$$
$$< \varepsilon,$$

where we used that

 $n \ge N \ge 3$

 $n^2-6\geq 0\,,$

which implies

in the third line. The last inequality holds, since it is equivalent to

$$N > \frac{1}{\sqrt{3\varepsilon}}$$

6.5 Bounded sequences

An important property of sequences is boundedness.

Definition 6.18: Bounded sequence

A sequence $(a_n)_{n \in \mathbb{N}}$ is called **bounded** if there exists a constant $M \in \mathbb{R}$, with M > 0, such that

 $|a_n| \leq M$, $\forall n \in \mathbb{N}$.

Definition 2.135 says that a sequence is bounded, if we can find some constant M > 0 (possibly very large), such that for all elements of the sequence it holds that

$$|a_n| \le M\,,$$

or equivalently, that

$$-M \le a_n \le M \, .$$

We now show that any sequence that converges is also bounded

Theorem 6.19

Every convergent sequence is bounded.

Proof

Suppose the sequence $(a_n)_{n \in \mathbb{N}}$ converges and let

$$a := \lim_{n \to \infty} a_n$$

By definition of convergence we have that

$$\forall \, \varepsilon > 0 \,, \, \exists N \in \mathbb{N} \, \text{ s.t. } \, \forall \, n \geq N \,, \, |a_n - a| < \varepsilon \,.$$

In particular, we can choose

 $\varepsilon = 1$

and let $N \in \mathbb{N}$ be that value such that

$$|a_n-a|<1\,,\quad\forall\,n\geq N\,.$$

If $n \ge N$ we have, by the triangle inequality,

$$|a_n| = |a_n - a + a|$$

 $\leq |a_n - a| + |a|$
 $< 1 + |a|$.

Set

$$M := \max\{|a_1|, |a_2|, \dots, |a_{N-1}|, 1+|a|\}.$$

Note that such maximum exists, being the set finite. Then

$$|a_n| \le M, \quad \forall \, n \in \mathbb{N}\,,$$

showing that (a_n) is bounded.

The choice of M in the above proof says that the sequence can behave wildly for a finite number of terms. After that, it will stay close to the value of the limit, if the latter exists.

Example 6.20

Question. Prove that the following sequence is bounded, and find *M*:

$$a_n=\frac{1}{n}$$
.

Solution. In Theorem 6.8 we have shown that

$$\lim_{n \to \infty} \frac{1}{n} = 0$$

Hence, it follows from Theorem 6.19 that the sequence (1/n) is bounded. An explicit constant *M* is found as follows:

$$\left|\frac{1}{n}\right| = \frac{1}{n} \le 1, \quad \forall n \in \mathbb{N},$$

since $n \ge 1$ for all $n \in \mathbb{N}$. Therefore M = 1.

Warning

The converse of Theorem 6.19 does not hold: There exist sequences (a_n) which are bounded, but not convergent.

Example 6.21

The sequence

 $a_n = (-1)^n$

is bounded (M = 1) but not convergent.

Proof. We have proven in Theorem 6.15 that (a_n) is not convergent. However (a_n) is bounded, with M = 1, since

$$|a_n| = |(-1)^n| = 1 = M, \quad \forall n \in \mathbb{N}.$$

Taking the contrapositive of the statement in Theorem 6.19 we get the following corollary:

Corollary 6.22

If a sequence is not bounded, then the sequence does not converge.

Remark 6.23

For a sequence (a_n) to be unbounded, it means that

 $\forall M > 0, \exists n \in \mathbb{N} \text{ s.t. } |a_n| > M.$

Comment. The above is saying that no real number M > 0 can be a bound for $|a_n|$, since there is always an index $n \in \mathbb{N}$ such that

 $|a_n| > M.$

We can use Corollary 6.22 to show that certain sequences do not converge.

Theorem 6.24

Let p > 0. The sequence $a_n = n^p$ is unbounded, and hence divergent.

Proof

Let p > 0. We prove that the sequence $(n^p)_{n \in \mathbb{N}}$ is unbounded, that is,

$$\forall M > 0, \exists n \in \mathbb{N} \text{ s.t. } |a_n| > M.$$

To this end, let M > 0. Choose $n \in \mathbb{N}$ such that

$$n > M^{1/p} \,.$$

Then

$$a_n = n^p > \left(M^{1/p}\right)^p = M.$$

This proves that the sequence (n^p) is unbounded. Hence (n^p) cannot converge, by Corollary 6.22.

Theorem 6.25

The sequence $a_n = \log n$ is unbounded, and hence divergent.

Proof

Let us show that $(\log n)_{n \in \mathbb{N}}$ is unbounded, that is,

 $\forall M > 0, \exists n \in \mathbb{N} \text{ s.t. } |a_n| > M.$

To this end let M > 0. Choose $n \in \mathbb{N}$ such that

 $n \ge e^{M+1}$.

Then

$$|a_n| = |\log n| \ge |\log e^{M+1}| = M + 1 > M$$

This proves that the sequence $(\log n)$ is unbounded. Hence $(\log n)$ cannot converge, by Corollary 6.22.

6.6 Algebra of limits

Proving convergence using Definition 2.129 can be a tedious task. In this section we discuss how to prove convergence, starting from known convergence results.

Theorem 6.26: Algebra of limits

Let $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be sequences in \mathbb{R} . Suppose that

$$\lim_{n\to\infty}a_n=a\,,\quad \lim_{n\to\infty}b_n=b\,,$$

for some $a, b \in \mathbb{R}$. Then,

1. Limit of sum is the sum of limits:

$$\lim_{n \to \infty} \left(a_n \pm b_n \right) = a \pm b$$

2. Limit of product is the product of limits:

$$\lim_{n \to \infty} \left(a_n b_n \right) = a k$$

3. If $b_n \neq 0$ for all $n \in \mathbb{N}$ and $b \neq 0$, then

$$\lim_{n \to \infty} \left(\frac{a_n}{b_n} \right) = \frac{a}{b}$$

Proof

Let $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be sequences in \mathbb{R} and let $c \in \mathbb{R}$. Suppose that, for some $a, b \in \mathbb{R}$

$$\lim_{n\to\infty}a_n=a\,,\quad \lim_{n\to\infty}b_n=b\,.$$

Proof of Point 1. We need to show that

$$\lim_{n\to\infty}(a_n\pm b_n)=a\pm b\,.$$

We only give a proof of the formula with +, since the case with - follows with a very similar proof. Hence, we need to show that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |(a_n + b_n) - (a + b)| < \varepsilon.$$

Let $\varepsilon > 0$ and set

$$\tilde{\varepsilon} := \frac{\varepsilon}{2}.$$

Since $a_n \to a, b_n \to b$, and $\tilde{\varepsilon} > 0$, there exist $N_1, N_2 \in \mathbb{N}$ such that

$$\begin{split} |a_n-a| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_1\,, \\ |b_n-b| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_2\,. \end{split}$$

Define

 $N := \max\{N_1, N_2\}.$

For all $n \ge N$ we have, by the triangle inequality,

$$|(a_n + b_n) - (a + b)| = |(a_n - a) + (b_n - b)|$$

$$\leq |a_n - a| + |b_n - b|$$

$$< \tilde{\varepsilon} + \tilde{\varepsilon}$$

$$= \varepsilon.$$

Proof of Point 2. We need to show that

$$\lim_{n\to\infty}(a_nb_n)=ab\,,$$

which is equivalent to

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |a_n b_n - ab| < \varepsilon.$$

Let $\varepsilon > 0$. The sequence (a_n) converges, and hence is bounded, by Theorem 6.19. This means there exists some M > 0 such that

$$|a_n| \leq M, \quad \forall n \in \mathbb{N}.$$

Define

$$\tilde{\varepsilon} = \frac{\varepsilon}{M + |b|}$$

Since $a_n \to a, b_n \to b$, and $\tilde{\varepsilon} > 0$, there exist $N_1, N_2 \in \mathbb{N}$ such that

$$\begin{split} |a_n-a| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_1\,. \\ |b_n-b| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_2\,. \end{split}$$

Let

$$N := \max\{N_1, N_2\}.$$

For all $n \ge N$ we have

$$\begin{aligned} |a_n b_n - ab| &= |a_n b_n - a_n b + a_n b - ab| \\ &\leq |a_n b_n - a_n b| + |a_n b - ab| \\ &= |a_n| |b_n - b| + |b| |a_n - a| \\ &< M \,\tilde{\varepsilon} + |b| \,\tilde{\varepsilon} \\ &= (M + |b|) \,\tilde{\varepsilon} \\ &= \varepsilon \,. \end{aligned}$$

Proof of Point 3. Suppose in addition that $b_n \neq 0$ and $b \neq 0$. We need to show that

$$\lim_{n \to \infty} \frac{a_n}{b_n} = \frac{a}{b}$$

which is equivalent to

$$\forall \, \varepsilon > 0 \,, \, \exists \, N \in \mathbb{N} \; \text{ s.t. } \; \forall \, n \geq N \,, \, \left| \frac{a_n}{b_n} - \frac{a}{b} \right| < \varepsilon \,.$$

We suppose in addition that b > 0. The proof is very similar for the case b < 0, and is hence omitted. Let $\varepsilon > 0$. Set

$$\delta := \frac{b}{2}.$$

Since $b_n \to b$ and $\delta > 0,$ there exists $N_1 \in \mathbb{N}$ such that

$$|b_n - b| < \delta \quad \forall \, n \ge N_1 \, .$$

In particular we have

$$b_n > b - \delta = b - \frac{b}{2} = \frac{b}{2} \quad \forall n \ge N_1 \,.$$

Define

$$\tilde{\varepsilon} := \frac{b^2}{2(b+|a|)} \, \varepsilon \, .$$

Since $\tilde{\varepsilon} > 0$ and $a_n \to a, b_n \to b$, there exist $N_2, N_3 \in \mathbb{N}$ such that

$$\begin{split} |a_n-a| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_2\,, \\ |b_n-b| &< \tilde{\varepsilon}\,, \quad \forall\, n \geq N_3\,. \end{split}$$

Define

$$N := \max\{N_1, N_2, N_3\}.$$

For all $n \ge N$ we have

$$\begin{aligned} \left|\frac{a_n}{b_n} - \frac{a}{b}\right| &= \left|\frac{a_n b - ab_n}{b_n b}\right| \\ &= \frac{1}{|b_n b|} |a_n b - ab + ab - ab_n| \\ &= \frac{1}{|b_n b|} |(a_n - a)b + a(b - b_n)| \\ &\leq \frac{1}{|b_n b|} (|a_n - a||b| + |a||b - b_n|) \\ &< \frac{1}{\frac{b}{2} b} (\tilde{\epsilon} b + \tilde{\epsilon}|a|) \\ &= \frac{2(b + |a|)}{b^2} \tilde{\epsilon} \\ &= \epsilon . \end{aligned}$$

In the future we will refer to Theorem 2.142 as the *Algebra of Limits*. We now show how to use Theorem 2.142 for computing certain limits.

Example 6.27

Question. Prove that

$$\lim_{n \to \infty} \frac{3n}{7n+4} = \frac{3}{7}$$

Solution. We can rewrite

$$\frac{3n}{7n+4} = \frac{3}{7+\frac{4}{n}}$$

By Theorem 6.12 we know that

$$3 \rightarrow 3$$
, $4 \rightarrow 4$, $7 \rightarrow 7$.

From Theorem 6.8 we know that

$$\frac{1}{n} \to 0$$
.

Hence, it follows from Theorem 2.142 Point 2 that

$$\frac{4}{n} = 4 \cdot \frac{1}{n} \to 4 \cdot 0 = 0.$$

By Theorem 2.142 Point 1 we have

$$7 + \frac{4}{n} \rightarrow 7 + 0 = 7 \,.$$

Finally we can use Theorem 2.142 Point 3 to infer

$$\frac{3n}{7n+4} = \frac{3}{7+\frac{4}{n}} \to \frac{3}{7} \,.$$

Important

The technique shown in Example 6.27 is useful to compute limits of fractions of polynomials. To identify the possible limit, if it exists, it is often best to divide by the largest power of n in the denominator.

Example 6.28

Question. Prove that

$$\lim_{n \to \infty} \frac{n^2 - 1}{2n^2 - 3} = \frac{1}{2} \,.$$

1

Solution. Factor n^2 to obtain

$$\frac{n^2 - 1}{2n^2 - 3} = \frac{1 - \frac{1}{n^2}}{2 - \frac{3}{n^2}}$$

By Theorem 6.9 we have

$$\frac{1}{n^2} \rightarrow 0$$

We can then use the Algebra of Limits Theorem 2.142 Point 2 to infer

$$\frac{3}{n^2} \to 3 \cdot 0 = 0$$

and Theorem 2.142 Point 1 to get

$$1 - \frac{1}{n^2} \to 1 - 0 = 1$$
, $2 - \frac{3}{n^2} \to 2 - 0 = 2$.

Finally we use Theorem 2.142 Point 3 and conclude

$$\frac{1-\frac{1}{n^2}}{2-\frac{3}{n^2}} \to \frac{1}{2}$$

Therefore

$$\lim_{n \to \infty} \frac{n^2 - 1}{2n^2 - 3} = \lim_{n \to \infty} \frac{1 - \frac{1}{n^2}}{2 - \frac{3}{n^2}} = \frac{1}{2}.$$

We can also use the Algebra of Limits to prove that certain limits do not exist.

Example 6.29

Question. Prove that the sequence

$$a_n = \frac{4n^3 + 8n + 1}{7n^2 + 2n + 1}$$

does not converge.

Solution. To show that the sequence (a_n) does not converge, we divide by the largest power in the denominator, which in this case is n^2

$$a_n = \frac{4n^3 + 8n + 1}{7n^2 + 2n + 1}$$
$$= \frac{4n + \frac{8}{n} + \frac{1}{n^2}}{7 + \frac{2}{n} + \frac{1}{n^2}}$$
$$= \frac{b_n}{c_n}$$

where we set

$$b_n := 4n + \frac{8}{n} + \frac{1}{n^2}$$
, $c_n := 7 + \frac{2}{n} + \frac{1}{n^2}$.

Using the Algebra of Limits Theorem 2.142 we see that

$$c_n = 7 + \frac{2}{n} + \frac{1}{n^2} \to 7$$
.

Suppose by contradiction that

$$a_n \rightarrow a$$

for some $a \in \mathbb{R}$. Then, by the Algebra of Limits Theorem 2.142 we would infer

$$b_n = c_n \cdot a_n \to 7a$$
,

concluding that b_n is convergent to 7a. We have that

$$b_n = 4n + d_n$$
, $d_n := \frac{8}{n} + \frac{1}{n^2}$.

Again by the Algebra of Limits Theorem 2.142 we get that

$$d_n=\frac{8}{n}+\frac{1}{n^2}\to 0\,,$$

and hence

$$4n = b_n - d_n \to 7a - 0 = 7a$$

This is a contradiction, since the sequence (4n) is unbounded, and hence cannot be convergent. Hence (a_n) is not convergent.

Warning

Consider the sequence

$$a_n = \frac{4n^3 + 8n + 1}{7n^2 + 2n + 1}$$

from the previous example. We have proven that (a_n) is not convergent, by making use of the Algebra of Limits.

Let us review a **faulty** argument to conclude that (a_n) is not convergent. Write

$$a_n = \frac{b_n}{c_n}$$
, $b_n := 4n^3 + 8n + 1$, $c_n := 7n^2 + 2n + 1$.

The numerator

 $b_n = 4n^3 + 8n + 1$

and denominator

$$c_n = 7n^2 + 2n + 1$$

are both unbounded, and hence (b_n) and (c_n) do not converge. One might be tempted to conclude that (a_n) does not converge. However this is **false** in general: as seen in Example 6.28, we have

$$\lim_{n \to \infty} \frac{n^2 - 1}{2n^2 - 3} = \frac{1}{2} \,,$$

while numerator and denominator are unbounded.

Sometimes it is useful to rearrange the terms of a sequence, before applying the Algebra of Limits.

Example 6.30

Question. Define the sequence

$$a_n := \frac{2n^3 + 7n + 1}{5n + 9} \cdot \frac{8n + 9}{6n^3 + 8n^2 + 3}.$$

Prove that

$$\lim_{n \to \infty} a_n = \frac{8}{15}$$

Solution. The first fraction in (a_n) does not converge, as it is unbounded. Therefore we cannot use Point 2 in Theorem 2.142 directly. However, we note that

$$a_n = \frac{2n^3 + 7n + 1}{5n + 9} \cdot \frac{8n + 9}{6n^3 + 8n^2 + 3}$$
$$= \frac{8n + 9}{5n + 9} \cdot \frac{2n^3 + 7n + 1}{6n^3 + 8n^2 + 3}.$$

Factoring out n and n^3 , respectively, and using the Algebra of Limits, we see that

$$\frac{8n+9}{5n+9} = \frac{8+9/n}{5+9/n} \to \frac{8+0}{5+0} = \frac{8}{5}$$

and

$$\frac{2+7/n^2+1/n^3}{6+8/n+3/n^3} \to \frac{2+0+0}{6+0+0} = \frac{1}{3}$$

Therefore Theorem 2.142 Point 2 ensures that

$$a_n \to \frac{8}{5} \cdot \frac{1}{3} = \frac{8}{15}$$

6.7 Fractional powers

The Algebra of Limits Theorem 2.142 can also be used when fractional powers of *n* are involved.

Example 6.31

Question. Prove that

$$a_n = \frac{n^{7/3} + 2\sqrt{n} + 7}{4n^{3/2} + 5n}$$

does not converge.

Solution. The largest power of *n* in the denominator is $n^{3/2}$. Hence we factor out $n^{3/2}$

$$a_n = \frac{n^{7/3} + 2\sqrt{n} + 7}{4n^{3/2} + 5n}$$

= $\frac{n^{7/3 - 3/2} + 2n^{1/2 - 3/2} + 7n^{-3/2}}{4 + 5n^{-3/2}}$
= $\frac{n^{5/6} + 2n^{-1} + 7n^{-3/2}}{4 + 5n^{-3/2}}$
= $\frac{b_n}{c_n}$

where we set

$$b_n := n^{5/6} + 2n^{-1} + 7n^{-3/2}, \quad c_n := 4 + 5n^{-3/2}.$$

We see that b_n is unbounded while $c_n \rightarrow 4$. By the Algebra of Limits (and usual contradiction argument) we conclude that (a_n) is divergent.

We now present a general result about the square root of a sequence.

Theorem 6.32

Let $(a_n)_{n \in \mathbb{N}}$ be a sequence in \mathbb{R} such that

 $\lim_{n\to\infty} a_n = a\,,$

for some $a \in \mathbb{R}$. If $a_n \ge 0$ for all $n \in \mathbb{N}$ and $a \ge 0$, then

$$\lim_{n\to\infty}\sqrt{a_n}=\sqrt{a}\,.$$

Proof

Let $\varepsilon > 0$. We the two cases a > 0 and a = 0:

• *a* > 0: Define

$$\delta := \frac{a}{2}$$
.

Since $\delta > 0$ and $a_n \to a$, there exists $N_1 \in \mathbb{N}$ such that

$$|a_n-a|<\delta\,,\quad\forall\,n\geq N_1\,.$$

In particular

$$a_n > a - \delta = a - \frac{a}{2} = \frac{a}{2}, \quad \forall n \ge N_1,$$

from which we infer

$$\sqrt{a_n} > \sqrt{a/2} \,, \quad \forall \, n \ge N_1 \,,$$

Now set

$$\tilde{\varepsilon} := \left(\sqrt{a/2} + \sqrt{a}\right) \varepsilon.$$

Since $\tilde{\varepsilon} > 0$ and $a_n \to a$, there exists $N_2 \in \mathbb{N}$ such that

$$|a_n-a|<\tilde{\varepsilon}\,,\quad\forall\,n\geq N_2\,.$$

Let

$$N := \max\{N_1, N_2\}.$$

For $n \ge N$ we have

$$\begin{aligned} \left|\sqrt{a_n} - \sqrt{a}\right| &= \left|\frac{\left(\sqrt{a_n} - \sqrt{a}\right)\left(\sqrt{a_n} + \sqrt{a}\right)}{\sqrt{a_n} + \sqrt{a}}\right| \\ &= \frac{|a_n - a|}{\sqrt{a_n} + \sqrt{a}} \\ &< \frac{\tilde{\varepsilon}}{\sqrt{a/2} + \sqrt{a}} \\ &= \varepsilon \,. \end{aligned}$$

• a = 0: In this case

$$a_n \rightarrow a = 0$$
.

Since $\varepsilon^2 > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - 0| = |a_n| < \varepsilon^2 \,, \quad \forall \, n \ge N \,.$$

Therefore

$$|\sqrt{a_n} - \sqrt{0}| = |\sqrt{a_n}| < \sqrt{\varepsilon^2} = \varepsilon, \quad \forall n \ge N.$$

Let us show an application of Theorem 6.32.

Example 6.33

Question. Define the sequence

$$a_n = \sqrt{9n^2 + 3n + 1} - 3n \, .$$

Prove that

$$\lim_{n\to\infty} a_n = \frac{1}{2} \, .$$

Solution. We first rewrite

$$\begin{split} a_n &= \sqrt{9n^2 + 3n + 1} - 3n \\ &= \frac{\left(\sqrt{9n^2 + 3n + 1} - 3n\right)\left(\sqrt{9n^2 + 3n + 1} + 3n\right)}{\sqrt{9n^2 + 3n + 1} + 3n} \\ &= \frac{9n^2 + 3n + 1 - (3n)^2}{\sqrt{9n^2 + 3n + 1} + 3n} \\ &= \frac{3n + 1}{\sqrt{9n^2 + 3n + 1} + 3n} \,. \end{split}$$

The biggest power of *n* in the denominator is *n*. Therefore we factor out *n*:

$$a_n = \sqrt{9n^2 + 3n + 1} - 3n$$

= $\frac{3n + 1}{\sqrt{9n^2 + 3n + 1} + 3n}$
= $\frac{3 + \frac{1}{n}}{\sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} + 3}$.

By the Algebra of Limits we have

$$9 + \frac{3}{n} + \frac{1}{n^2} \to 9 + 0 + 0 = 9.$$

Therefore we can use Theorem 6.32 to infer

$$\sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} \to \sqrt{9}$$

By the Algebra of Limits we conclude:

$$a_n = \frac{3 + \frac{1}{n}}{\sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} + 3} \to \frac{3 + 0}{\sqrt{9} + 3} = \frac{1}{2}.$$

Example 6.34

Question. Prove that the sequence

$$a_n = \sqrt{9n^2 + 3n + 1} - 2n$$

does not converge. **Solution.** We rewrite a_n as

$$\begin{split} a_n &= \sqrt{9n^2 + 3n + 1} - 2n \\ &= \frac{(\sqrt{9n^2 + 3n + 1} - 2n)(\sqrt{9n^2 + 3n + 1} + 2n)}{\sqrt{9n^2 + 3n + 1} + 2n} \\ &= \frac{9n^2 + 3n + 1 - (2n)^2}{\sqrt{9n^2 + 3n + 1} + 2n} \\ &= \frac{5n^2 + 3n + 1}{\sqrt{9n^2 + 3n + 1} + 2n} \\ &= \frac{5n + 3 + \frac{1}{n}}{\sqrt{9n^2 + 3n + 1} + 2n} \\ &= \frac{5n + 3 + \frac{1}{n}}{\sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} + 2} \\ &= \frac{b_n}{c_n} \,, \end{split}$$

where we factored n, being it the largest power of n in the denominator, and defined

$$b_n := 5n + 3 + \frac{1}{n}$$
, $c_n := \sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} + 2$.

Note that

$$9 + \frac{3}{n} + \frac{1}{n^2} \to 9$$

by the Algebra of Limits. Therefore

$$\sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} \to \sqrt{9} = 3$$

by Theorem 6.32. Hence

$$c_n = \sqrt{9 + \frac{3}{n} + \frac{1}{n^2}} + 2 \rightarrow 3 + 2 = 5.$$

The numerator

$$b_n = 5n + 3 + \frac{1}{n}$$

is instead unbounded. Therefore (a_n) is not convergent, by the Algebra of Limits and the usual contradiction argument.

6.8 Limit Tests

In this section we discuss a number of *Tests* to determine whether a sequence converges or not. These are known as **Limit Tests**.

6.8.1 Squeeze Theorem

When a sequence (a_n) oscillates, it is difficult to compute the limit. Examples of terms which produce oscillations are

```
(-1)^n, \sin(n), \cos(n).
```

In such instance it might be useful to compare (a_n) with other sequences whose limit is known. If we can prove that (a_n) is *squeezed* between two other sequences with the same limiting value, then we can show that also (a_n) converges to this value.

Theorem 6.35: Squeeze theorem

Let (a_n) , (b_n) and (c_n) be sequences in \mathbb{R} . Suppose that

$$b_n \leq a_n \leq c_n$$
, $\forall n \in \mathbb{N}$,

and that

$$\lim_{n \to \infty} b_n = \lim_{n \to \infty} c_n = L$$

Then

 $\lim_{n\to\infty}a_n=L\,.$

Proof

Let $\varepsilon > 0$. Since $b_n \to L$ and $c_n \to L$, there exist $N_1, N_2 \in \mathbb{N}$ such that

$$\begin{split} &-\varepsilon < b_n - L < \varepsilon \;, \; \forall \; n \geq N_1 \;, \\ &-\varepsilon < c_n - L < \varepsilon \;, \; \forall \; n \geq N_2 \;. \end{split}$$

Set

 $N := \max\{N_1, N_2\}.$

Let $n \ge N$. Using the assumption that $b_n \le a_n \le c_n$, we get

 $b_n - L \le a_n - L \le c_n - L \,.$

In particular

 $-\varepsilon < b_n - L \le a_n - L \le b_n - L < \varepsilon \,.$

The above implies

$$-\varepsilon < a_n - L < \varepsilon \quad \Longrightarrow \quad |a_n - L| < \varepsilon \,.$$

Example 6.36

Question. Prove that

$$\lim_{n\to\infty} \frac{(-1)^n}{n} = 0.$$

Solution. For all $n \in \mathbb{N}$ we can estimate

$$-1 \le (-1)^n \le 1 \, .$$

Therefore

$$\frac{-1}{n} \le \frac{(-1)^n}{n} \le \frac{1}{n}, \quad \forall n \in \mathbb{N}.$$

Moreover

$$\lim_{n \to \infty} \frac{-1}{n} = -1 \cdot 0 = 0, \quad \lim_{n \to \infty} \frac{1}{n} = 0.$$

By the Squeeze Theorem 2.151 we conclude

$$\lim_{n\to\infty}\frac{(-1)^n}{n}=0\,.$$

Example 6.37

Question. Prove that

$$\lim_{n \to \infty} \frac{\cos(3n) + 9n^2}{11n^2 + 15\sin(17n)} = \frac{9}{11}$$

Solution. We know that

 $-1 \le \cos(x) \le 1$, $-1 \le \sin(x) \le 1$, $\forall x \in \mathbb{R}$.

Therefore, for all $n \in \mathbb{N}$

$$-1 \le \cos(3n) \le 1, \quad -1 \le \sin(17n) \le 1.$$

We can use the above to estimate the numerator in the given sequence:

$$-1 + 9n^2 \le \cos(3n) + 9n^2 \le 1 + 9n^2.$$
(6.7)

Concerning the denominator, we have

$$11n^2 - 15 \le 11n^2 + 15\sin(17n) \le 11n^2 + 15$$

and therefore

$$\frac{1}{11n^2 + 15} \le \frac{1}{11n^2 + 15\sin(17n)} \le \frac{1}{11n^2 - 15} \,. \tag{6.8}$$

Putting together (6.7)-(6.8) we obtain

$$\frac{-1+9n^2}{11n^2+15} \le \frac{\cos(3n)+9n^2}{11n^2+15\sin(17n)} \le \frac{1+9n^2}{11n^2-15} \,.$$

By the Algebra of Limits we infer

$$\frac{-1+9n^2}{11n^2+15} = \frac{-\frac{1}{n^2}+9}{11+\frac{15}{n^2}} \to \frac{0+9}{11+0} = \frac{9}{11}$$

and

$$\frac{1+9n^2}{11n^2-15} = \frac{\frac{1}{n^2}+9}{11-\frac{15}{n^2}} \to \frac{0+9}{11+0} = \frac{9}{11}$$

Applying the Squeeze Theorem 2.151 we conclude

$$\lim_{n \to \infty} \frac{\cos(3n) + 9n^2}{11n^2 + 15\sin(17n)} = \frac{9}{11}$$

Warning

Suppose that the sequences $(a_n), (b_n), (c_n)$ satisfy

$$b_n \leq a_n \leq c_n$$
, $\forall n \in \mathbb{N}$,

and

$$b_n \to L_1$$
, $c_n \to L_2$, $L_1 \neq L_2$.

In general, we cannot conclude that a_n converges.

Example 6.38

Consider the sequence

$$a_n = \left(1 + \frac{1}{n}\right)(-1)^n$$

For all $n \in \mathbb{N}$ we can bound

$$-1 - \frac{1}{n} \le \left(1 + \frac{1}{n}\right)(-1)^n \le 1 + \frac{1}{n}.$$

However

$$-1 - \frac{1}{n} \longrightarrow -1 - 0 = -1$$

and

$$1 + \frac{1}{n} \longrightarrow 1 + 0 = 1.$$

Since

we cannot apply the Squeeze Theorem 2.151 to conclude convergence of (a_n) . Indeed, (a_n) is a divergent sequence.

 $-1 \neq 1$,

Proof. Suppose by contradiction that $a_n \rightarrow a$. We have

$$a_n = (-1)^n + \frac{(-1)^n}{n} = b_n + c_n$$

where

$$b_n := (-1)^n$$
, $c_n := \frac{(-1)^n}{n}$.

We have seen in Example 6.37 that $c_n \rightarrow 0$. Therefore, by the Algebra of Limits, we have

$$b_n = a_n - c_n \longrightarrow a - 0 = a$$
.

However, Theorem 6.15 says that the sequence $b_n = (-1)^n$ diverges. Contradiction. Hence (a_n) diverges.

6.8.2 Geometric sequences

Definition 6.39: Geometric sequence

A sequence (a_n) is called a **geometric sequence** if

$$a_n = x^n$$

for some $x \in \mathbb{R}$.

The value of |x| determines whether or not a geometric sequence converges, as shown in the following theorem.

Theorem 6.40: Geometric Sequence Test

Let $x \in \mathbb{R}$ and let (a_n) be the sequence defined by $a_n := x^n$. We have:

1. If |x| < 1, then

$$\lim_{n\to\infty}a_n=0\,.$$

2. If |x| > 1, then sequence (a_n) is unbounded, and hence divergent.

Warning

The Geometric Sequence Test in Theorem 2.154 does not address the case

|x| = 1.

This is because, in this case, the sequence

 $a_n = x^n$

might converge or diverge, depending on the value of x. Indeed,

$$|x| = 1 \quad \Longrightarrow \quad x = \pm 1.$$

We therefore have two cases:

• *x* = 1: Then

$$a_n = 1^n = 1$$

so that $a_n \to 1$ and (a_n) is convergent.

• x = -1: Then

$$a_n = x^n = (-1)^n$$

which is divergent by Theorem 6.15.

To prove Theorem 2.154 we need the following inequality, known as Bernoulli's inequality.

Lemma 6.41: Bernoulli's inequality

Let $x \in \mathbb{R}$ with x > -1. Then

$$(1+x)^n \ge 1 + nx, \quad \forall n \in \mathbb{N}.$$
(6.9)

Proof

Let $x \in \mathbb{R}$, x > -1. We prove the statement by induction:

- Base case: (6.9) holds with equality when n = 1.
- Induction hypothesis: Let $k \in \mathbb{N}$ and suppose that (6.9) holds for n = k, i.e.,

$$(1+x)^k \ge 1+kx.$$

Then

$$(1+x)^{k+1} = (1+x)^k (1+x)$$

$$\ge (1+kx)(1+x)$$

$$= 1+kx+x+kx^2$$

$$\ge 1+(k+1)x,$$

where we used that $kx^2 \ge 0$. Then (6.9) holds for n = k + 1.

By induction we conclude (6.9).

We are ready to prove Theorem 2.154.

Proof: Proof of Theorem 2.154

Part 1. The case |x| < 1. If x = 0, then

$$a_n = x^n = 0$$

so that $a_n \to 0$. Hence assume $x \neq 0$. We need to prove that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |x^n - 0| < \varepsilon$$

Let $\varepsilon > 0$. We have

$$|x| < 1 \qquad \Longrightarrow \qquad \frac{1}{|x|} > 1 \,.$$

Therefore

$$|x| = \frac{1}{1+u}$$
, $u := \frac{1}{|x|} - 1 > 0$.

Let $N \in \mathbb{N}$ be such that

 $N > \frac{1}{\varepsilon u}$,

so that

$$\frac{1}{Nu} < \varepsilon$$

Let $n \ge N$. Then

$$\begin{aligned} x^{n} - 0 &| = |x|^{n} \\ &= \left(\frac{1}{1+u}\right)^{n} \\ &= \frac{1}{(1+u)^{n}} \\ &\leq \frac{1}{1+nu} \\ &\leq \frac{1}{nu} \\ &\leq \frac{1}{Nu} \\ &\leq \varepsilon \, . \end{aligned}$$

where we used Bernoulli's inequality (6.9) in the first inequality.

Part 2. The case |x| > 1*.*

To prove that (a_n) does not converge, we prove that it is unbounded. This means showing that

$$\forall M > 0, \exists n \in \mathbb{N} \text{ s.t. } |a_n| > M.$$

Let M > 0. We have two cases:

• $0 < M \le 1$: Choose n = 1. Then

$$|a_1| = |x| > 1 \ge M$$
.

• M > 1: Choose $n \in \mathbb{N}$ such that

$$n > \frac{\log M}{\log |x|}$$

Note that $\log |x| > 0$ since |x| > 1. Therefore

$$n > \frac{\log M}{\log |x|} \iff n \log |x| > \log M$$
$$\iff \log |x|^n > \log M$$
$$\iff |x|^n > M.$$

Then

$$|a_n| = |x^n| = |x|^n > M$$
.

Hence (a_n) is unbounded. By Corollary 6.22 we conclude that (a_n) is divergent.

Example 6.42

We can apply Theorem 2.154 to prove convergence or divergence for the following sequences.

 $\left(\frac{1}{2}\right)^n \longrightarrow 0$

 $\left|\frac{1}{2}\right| = \frac{1}{2} < 1$.

 $\left(\frac{-1}{2}\right)^n \longrightarrow 0$

 $\left|\frac{-1}{2}\right| = \frac{1}{2} < 1$.

1. We have

since

2. We have

since

3. The sequence

$$a_n = \left(\frac{-3}{2}\right)^n$$

does not converge, since

$$\left|\frac{-3}{2}\right| = \frac{3}{2} > 1$$

4.	As $n \to \infty$,	$\frac{3^n}{(-5)^n} = \left(-\frac{3}{5}\right)^n \longrightarrow 0$	
	since	$\left -\frac{3}{5}\right = \frac{3}{5} < 1 .$	
5.	The sequence	$a_n = \frac{(-7)^n}{2^{2n}}$	
	does not converge, since	$\frac{(-7)^n}{2^{2n}} = \frac{(-7)^n}{\left(2^2\right)^n} = \left(-\frac{7}{4}\right)^n$	
	and	$\left -\frac{7}{4}\right = \frac{7}{4} > 1$.	

6.8.3 Ratio Test

Theorem 6.43: Ratio Test

Let (a_n) be a sequence in \mathbb{R} such that

$$a_n \neq 0$$
, $\forall n \in \mathbb{N}$.

1. Suppose that the following limit exists:

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \, .$$

Then,

• If *L* < 1 we have

$$\lim_{n\to\infty}a_n=0$$

- If L > 1, the sequence (a_n) is unbounded, and hence does not converge.
- 2. Suppose that there exists $N \in \mathbb{N}$ and L > 1 such that

$$\left|\frac{a_{n+1}}{a_n}\right| \ge L\,, \quad \forall \, n \ge N\,.$$

Then the sequence (a_n) is unbounded, and hence does not converge.

Proof

Define the sequence $b_n = |a_n|$. Then,

$$\frac{a_{n+1}}{a_n} \bigg| = \frac{|a_{n+1}|}{|a_n|} = \frac{b_{n+1}}{b_n}$$

Part 1. Suppose that there exists the limit

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \, .$$

Therefore

$$\lim_{n \to \infty} \frac{b_{n+1}}{b_n} = L.$$
(6.10)

- L < 1: Choose r > 0 such that
- L < r < 1.

Set

$$\varepsilon := r - L$$

By the convergence at (6.10) there exists $N \in \mathbb{N}$ such that

$$\left|\frac{b_{n+1}}{b_n} - L\right| < \varepsilon = r - L, \quad \forall \, n \ge N \,.$$

In particular

$$\frac{b_{n+1}}{b_n} - L < r - L, \quad \forall n \ge N,$$

which implies

$$b_{n+1} < r \, b_n \,, \quad \forall \, n \ge N \,. \tag{6.11}$$

Let $n \ge N$, we can use (6.11) recursively and obtain

$$0 \le b_n < rb_{n-1} < \dots < r^{n-N} b_N = r^n \frac{b_N}{r^N}.$$

In particular, we have proven that

$$0 \le b_n < r^n \frac{b_N}{r^N}, \quad \forall n \in \mathbb{N}.$$
(6.12)

Since |r| < 1, by the Geometric Sequence Test Theorem 2.154 we infer

 $r^n \rightarrow 0$.

The Algebra of Limits the yields

$$r^n \, \frac{b_N}{r^N} \to 0$$

By the Squeeze Theorem 2.151 applied to (6.12), it follows that

$$b_n = |a_n| \to 0$$
.

Since

$$-|a_n| \le a_n \le |a_n| ,$$

and

$$-|a_n| \to 0$$
, $|a_n| \to 0$,

we can again apply the Squeeze Theorem ${\tt 2.151}$ to infer

$$a_n \to 0$$
.

• L > 1: Choose r > 0 such that

 $1 < r < L \, .$

Define

$$\varepsilon := L - r > 0 \, .$$

By the convergence (6.10), there exists $N \in \mathbb{N}$ such that

$$\left|\frac{b_{n+1}}{b_n} - L\right| < \varepsilon = L - r \,, \quad \forall \, n \geq N \,.$$

In particular,

$$-(L-r) < \frac{b_{n+1}}{b_n} - L, \quad \forall n \ge N,$$

which implies

$$b_{n+1} > r \, b_n \,, \quad \forall \, n \ge N \,. \tag{6.13}$$

Let $n \ge N$. Applying (6.13) recursively we get

$$b_n > r^{n-N} b_N = r^n \frac{b_N}{r^N}, \quad \forall n \ge N.$$
(6.14)

Since |r| > 1, by the Geometric Sequence Test we have that the sequence

 (r^n)

is unbounded. Therefore also the right hand side of (6.14) is unbounded, proving that (b_n) is unbounded. Since

$$b_n = |a_n|,$$

we conclude that (a_n) is unbounded. By Corollary 6.22 we conclude that (a_n) does not converge.

Part 2. Suppose that there exists $N \in \mathbb{N}$ and L > 1 such that

$$\left|\frac{a_{n+1}}{a_n}\right| \ge L, \quad \forall \, n \ge N \, .$$

Since $b_n = |a_n|$, we infer

$$b_{n+1} \ge L b_n$$
, $\forall n \ge N$.

Arguing as above, we obtain

$$b_n \ge L^n \frac{b_N}{L^N}, \quad \forall n \ge N.$$

Since L > 1, we have that the sequence

$$L^n \frac{b_N}{L^N}$$

is unbounded, by the Geometric Sequence Test. Hence also (b_n) is unbounded, from which we conclude that (a_n) is unbounded. By Corollary 6.22 we conclude that (a_n) does not converge.

Let us apply the Ratio Test to some concrete examples.

Example 6.44

Question. Prove that

$$a_n=\frac{3^n}{n!}\to 0\,,$$

where we recall that n! (pronounced n factorial) is defined by

$$n! := n \cdot (n-1) \cdot (n-2) \cdot \ldots \cdot 3 \cdot 2 \cdot 1.$$

Solution. We have

$$\left|\frac{a_{n+1}}{a_n}\right| = \frac{\left(\frac{3^{n+1}}{(n+1)!}\right)}{\left(\frac{3^n}{n!}\right)} = \frac{3}{n+1} \longrightarrow L = 0.$$

Hence, L = 0 < 1 so $a_n \rightarrow 0$ by the Ratio Test in Theorem 2.156.

Example 6.45

Question. Prove that the sequence is divergent

$$a_n = \frac{n! \cdot 3^n}{\sqrt{(2n)!}} \, .$$

Solution. We have

$$\begin{vmatrix} \frac{a_{n+1}}{a_n} \end{vmatrix} = \frac{(n+1)! \cdot 3^{n+1}}{\sqrt{(2(n+1))!}} \frac{\sqrt{(2n)!}}{n! \cdot 3^n} \\ = \frac{(n+1)!}{n!} \cdot \frac{3^{n+1}}{3^n} \cdot \frac{\sqrt{(2n)!}}{\sqrt{(2(n+1))!}} \end{vmatrix}$$

For the first two fractions we have

$$\frac{(n+1)!}{n!} \cdot \frac{3^{n+1}}{3^n} = 3(n+1),$$

while for the third fraction

$$\frac{\sqrt{(2n)!}}{\sqrt{(2(n+1))!}} = \sqrt{\frac{(2n)!}{(2n+2)!}}$$
$$= \sqrt{\frac{(2n)!}{(2n+2)\cdot(2n+1)\cdot(2n)!}}$$
$$= \frac{1}{\sqrt{(2n+1)(2n+2)}}.$$

Therefore, using the Algebra of Limits,

$$\begin{aligned} \frac{a_{n+1}}{a_n} &| = \frac{3(n+1)}{\sqrt{(2n+1)(2n+2)}} \\ &= \frac{3n\left(1+\frac{1}{n}\right)}{\sqrt{n^2\left(2+\frac{1}{n}\right)\left(2+\frac{2}{n}\right)}} \\ &= \frac{3\left(1+\frac{1}{n}\right)}{\sqrt{\left(2+\frac{1}{n}\right)\left(2+\frac{2}{n}\right)}} \longrightarrow \frac{3}{\sqrt{4}} = \frac{3}{2} > 1. \end{aligned}$$

By the Ratio Test we conclude that (a_n) is divergent.

Example 6.46

Question. Prove that the following sequence is divergent

$$a_n = \frac{n!}{100^n}$$

Solution. We have

$$\left|\frac{a_{n+1}}{a_n}\right| = \frac{100^n}{100^{n+1}} \frac{(n+1)!}{n!} = \frac{n+1}{100}.$$

Choose N = 101. Then for all $n \ge N$,

$$\left|\frac{a_{n+1}}{a_n}\right| = \frac{n+1}{100} \ge \frac{N+1}{100} = \frac{101}{100} > 1.$$

Hence a_n is divergent by the Ratio Test.

Warning

The Ratio Test in Theorem 2.156 does not address the case

L = 1 .

This is because, in this case, the sequence (a_n) might converge or diverge. For example:

• Define the sequence

$$a_n = \frac{1}{n}$$
.

We have

$$\left|\frac{a_{n+1}}{a_n}\right| = \frac{n}{n+1} \to L = 1.$$

Hence we cannot apply the Ratio Test. However we know that

$$\lim_{n\to\infty} \frac{1}{n} = 0.$$

• Consider the sequence

We have

$$\frac{|a_{n+1}|}{|a_n|} = \frac{n+1}{n} \to L = 1 \,.$$

 $a_n = n$.

Hence we cannot apply the Ratio Test. However we know that (a_n) is unbounded, and thus divergent.

If the sequence (a_n) is geometric, the Ratio Test of Theorem 2.156 will give the same answer as the Geometric Sequence Test of Theorem 2.154. This is the content of the following remark.

Remark 6.47

Let $x \in \mathbb{R}$ and define the geometric sequence

$$a_n = x^n$$

Then

$$\left|\frac{a_{n+1}}{a_n}\right| = \frac{|x^{n+1}|}{|x^n|} = \frac{|x|^{n+1}}{|x|^n} = |x| \to |x|.$$

Hence:

- If |x| < 1, the sequence (a_n) converges by the Ratio Test
- If |x| > 1, the sequence (a_n) diverges by the Ratio Test.
- If |x| = 1, the sequence (a_n) might be convergent or divergent.

These results are in agreement with the Geometric Sequence Test.

6.9 Monotone sequences

We have seen in Theorem 6.19 that convergent sequences are bounded. We noted that the converse statement is not true. For example the sequence

$$a_n = (-1)^n$$

is bounded but not convergent, as shown in Theorem 6.15. On the other hand, if a bounded sequence is **monotone**, then it is convergent.

Definition 6.48: Monotone sequence

Let (a_n) be a real sequence. We say that:

1. (a_n) is increasing if

 $a_n \leq a_{n+1}$, $\forall n \geq N$.

2. (a_n) is **decreasing** if

$$a_n \ge a_{n+1}$$
, $\forall n \ge N$.

3. (a_n) is **monotone** if it is either increasing or decreasing.

Example 6.49

Question. Prove that the following sequence is increasing

$$a_n=\frac{n-1}{n}\,.$$

Solution. We have

$$a_{n+1} = \frac{n}{n+1} > \frac{n-1}{n} = a_n$$
,

where the inequality holds because

$$\frac{n}{n+1} > \frac{n-1}{n} \quad \iff \quad n^2 > (n-1)(n+1)$$
$$\iff \quad n^2 > n^2 - 1$$
$$\iff \quad 0 > -1$$

Example 6.50

Question. Prove that the following sequence is decreasing

$$a_n=\frac{1}{n}$$
.

Solution. We immediately see that

$$a_n = \frac{1}{n} > \frac{1}{n+1} = a_{n+1}.$$

The main result about monotone sequences is the Monotone Convergence Theorem.

Theorem 6.51: Monotone Convergence Theorem

Let (a_n) be a sequence in \mathbb{R} . Suppose that (a_n) is bounded and monotone. Then (a_n) converges. In particular,

1. If a_n is increasing, then

$$\lim_{n\to\infty}a_n=\sup A\,,$$

2. If a_n is decreasing, then

$$\lim_{n\to\infty}a_n=\inf A\,,$$

where we define $A = \{a_n : n \in \mathbb{N}\}$.

Proof

Assume (a_n) is bounded and monotone. Since (a_n) is bounded, the set

$$A := \{a_n : n \in \mathbb{N}\} \subseteq \mathbb{R}$$

is bounded below and above. By the Axiom of Completeness of \mathbb{R} there exist *i*, *s* \in \mathbb{R} such that

$$i = \inf A$$
, $s = \sup A$.

We have two cases:

1. (a_n) is increasing: We are going to prove that

$$\lim_{n\to\infty}a_n=s\,.$$

Equivalently, we need to prove that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |a_n - s| < \varepsilon.$$
(6.15)

Let $\varepsilon > 0$. Since *s* is the smallest upper bound for *A*, this means

 $s - \varepsilon$
is not an upper bound. Therefore there exists $N \in \mathbb{N}$ such that

$$s - \varepsilon < a_N . \tag{6.16}$$

Let $n \ge N$. Since a_n is increasing, we have

$$a_N \le a_n, \quad \forall n \ge N.$$
 (6.17)

Moreover s is the supremum of A, so that

$$a_n \le s < s + \varepsilon, \quad \forall n \in \mathbb{N}.$$
 (6.18)

Putting together estimates (6.16)-(6.17)-(6.18) we get

$$s - \varepsilon < a_N \le a_n \le s < s + \varepsilon$$
, $\forall n \ge N$.

The above implies

 $s-\varepsilon < a_n < s+\varepsilon$, $\forall n \ge N$,

which is equivalent to (6.15).

2. (a_n) is decreasing: With a similar proof, one can show that

$$\lim_{n\to\infty}a_n=i\,.$$

This is left as an exercise.

6.9.1 Example: Euler's Number

As an application of the Monotone Convergence Theorem we can give a formal definition for the Euler's Number

e = 2.71828182845904523536...

Theorem 6.52

Consider the sequence

$$a_n = \left(1 + \frac{1}{n}\right)^n \, .$$

We have that:

1. (a_n) is monotone increasing,

2. (a_n) is bounded.

In particular (a_n) is convergent.

Proof

Part 1. We prove that (a_n) is increasing

$$a_n \ge a_{n-1}$$
, $\forall n \in \mathbb{N}$,

which by definition is equivalent to

$$\left(1+\frac{1}{n}\right)^n \ge \left(1+\frac{1}{n-1}\right)^{n-1}, \quad \forall n \in \mathbb{N}.$$

Summing the fractions we get

$$\left(\frac{n+1}{n}\right)^n \ge \left(\frac{n}{n-1}\right)^{n-1}$$

Multiplying by $((n-1)/n)^n$ we obtain

$$\left(\frac{n-1}{n}\right)^n \left(\frac{n+1}{n}\right)^n \ge \frac{n-1}{n}$$

which simplifies to

$$\left(1-\frac{1}{n^2}\right)^n \ge 1-\frac{1}{n}, \quad \forall n \in \mathbb{N}.$$
(6.19)

Therefore (a_n) is increasing if and only if (6.19) holds. Recall Bernoulli's inequality from Lemma 9.157: For $x \in \mathbb{R}$, x > -1, it holds

$$(1+x)^n \ge 1+nx$$
, $\forall n \in \mathbb{N}$.

Appliying Bernoulli's inequality with

$$x = -\frac{1}{n^2}$$

yields

$$\left(1-\frac{1}{n^2}\right)^n \ge 1+n\left(-\frac{1}{n^2}\right)=1-\frac{1}{n},$$

which is exactly (6.19). Then (a_n) is increasing. *Part 2.* We have to prove that (a_n) is bounded, that is, that there exists M > 0 such that

$$|a_n| \leq M, \quad \forall n \in \mathbb{N}.$$

To this end, introduce the sequence (b_n) by setting

$$b_n := \left(1 + \frac{1}{n}\right)^{n+1}$$

The sequence (b_n) is decreasing.

To prove (b_n) is decreasing, we need to show that

$$b_{n-1} \ge b_n$$
, $\forall n \in \mathbb{N}$.

By definition of b_n , the above reads

$$\left(1+\frac{1}{n-1}\right)^n \ge \left(1+\frac{1}{n}\right)^{n+1}, \quad \forall n \in \mathbb{N}.$$

Summing the terms inside the brackets, the above is equivalent to

$$\left(\frac{n}{n-1}\right)^n \ge \left(\frac{n+1}{n}\right)^n \left(\frac{n+1}{n}\right)$$
.

Multiplying by $(n/(n+1))^n$ we get

$$\left(\frac{n^2}{n^2-1}\right)^n \ge \left(\frac{n+1}{n}\right) \,.$$

The above is equivalent to

$$\left(1+\frac{1}{n^2-1}\right)^n \ge \left(1+\frac{1}{n}\right).$$
 (6.20)

Therefore (b_n) is decreasing if and only if (6.20) holds for all $n \in \mathbb{N}$. By choosing

$$x = \frac{1}{n^2 - 1}$$

in Bernoulli's inequality, we obtain

$$\left(1 + \frac{1}{n^2 - 1}\right)^n \ge 1 + n\left(\frac{1}{n^2 - 1}\right)$$
$$= 1 + \frac{n}{n^2 - 1}$$
$$\ge 1 + \frac{1}{n},$$

where in the last inequality we used that

$$\frac{n}{n^2-1} > \frac{1}{n} \,,$$

which holds, being equivalent to $n^2 > n^2 - 1$. We have therefore proven (6.20), and hence (b_n) is decreasing.

We now observe that For all $n\in\mathbb{N}$

$$b_n = \left(1 + \frac{1}{n}\right)^{n+1}$$
$$= \left(1 + \frac{1}{n}\right)^n \left(1 + \frac{1}{n}\right)$$
$$= a_n \left(1 + \frac{1}{n}\right)$$
$$> a_n .$$

Since (a_n) is increasing and (b_n) is decreasing, in particular

$$a_n \ge a_1$$
, $b_n \le b_1$.

Therefore

 $a_1 \leq a_n < b_n \leq b_1 \,, \quad \forall \, n \in \mathbb{N} \,.$

We compute

 $a_1 = 2, \quad b_1 = 4,$

from which we get

 $2 \le a_n \le 4 \,, \quad \forall \, n \in \mathbb{N} \,.$

Therefore

 $|a_n| \leq 4$, $\forall n \in \mathbb{N}$,

showing that (a_n) is bounded.

Part 3. The sequence (a_n) is increasing and bounded above. Therefore (a_n) is convergent by the Monotone Convergence Theorem 2.163.

Thanks to Theorem 6.52 we can define the Euler's Number *e*.

Definition 6.53: Euler's Number

The Euler's number is defined as

$$e := \lim_{n \to \infty} \left(1 + \frac{1}{n} \right)^n \, .$$

Setting n = 1000 in the formula for (a_n) , we get an approximation of e:

$$e \approx a_{1000} = 2.7169$$
.

6.10 Special limits

In this section we investigate limits of some sequences to which the Limit Tests do not apply.

Theorem 6.54

Let $x \in \mathbb{R}$, with x > 0. Then

$$\lim_{n\to\infty}\sqrt[n]{x}=1\,.$$

Proof

Step 1. Assume
$$x \ge 1$$
. In this case

 $\sqrt[n]{x} \ge 1$.

Define

 $b_n := \sqrt[n]{x} - 1,$

so that $b_n \ge 0$. By Bernoulli's Inequality we have

$$x = (1+b_n)^n \ge 1+nb_n.$$

Therefore

$$0 \le b_n \le \frac{x-1}{n} \, .$$

Since

$$\frac{x-1}{n}\longrightarrow 0\,,$$

by the Squeeze Theorem we infer $b_n \rightarrow 0$, and hence

$$\sqrt[n]{x} = 1 + b_n \longrightarrow 1 + 0 = 1,$$

by the Algebra of Limits. *Step 2.* Assume 0 < x < 1. In this case

$$\frac{1}{x} > 1$$
.

Therefore

$$\lim_{n\to\infty}\sqrt[n]{1/x}=1.$$

by Step 1. Therefore

$$\sqrt[n]{x} = \frac{1}{\sqrt[n]{1/x}} \longrightarrow \frac{1}{1} = 1,$$

by the Algebra of Limits.

Theorem 6.55

Let (a_n) be a sequence such that $a_n \to 0$. Then

$$\sin(a_n) \to 0$$
, $\cos(a_n) \to 1$.

Proof

Assume that $a_n \rightarrow 0$ and set

$$\varepsilon := \frac{\pi}{2}$$
.

By the convergence $a_m \to 0$ there exists $N \in \mathbb{N}$ such that

$$|a_n| < \varepsilon = \frac{\pi}{2} \quad \forall n \ge N.$$
 (6.21)

Step 1. We prove that

 $\sin(a_n) \to 0$.

By elementary trigonometry we have

$$0 \le |\sin(x)| = \sin |x| \le |x|, \quad \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right].$$

Therefore, since (6.21) holds, we can substitute $x = a_n$ in the above inequality to get

$$0 \le |\sin(a_n)| \le |a_n|, \quad \forall n \ge \mathbb{N}.$$

Since $a_n \to 0$, we also have $|a_n| \to 0$. Therefore $|\sin(a_n)| \to 0$ by the Squeeze Theorem. This immediately implies $\sin(a_n) \to 0$. Step 2. We prove that

 $\cos(a_n) \to 1$.

Inverting the relation

$$\cos^2(x) + \sin^2(x) = 1,$$

we obtain

$$\cos(x) = \pm \sqrt{1 - \sin^2(x)} \,.$$

We have that $\cos(x) \ge 0$ for $-\pi/2 \le x \le \pi/2$. Thus

$$\cos(x) = \sqrt{1 - \sin^2(x)}, \quad \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right].$$

Since (6.21) holds, we can set $x = a_n$ in the above inequality and obtain

$$\cos(a_n) = \sqrt{1 - \sin^2(a_n)}, \quad \forall n \ge N.$$

By Step 1 we know that $sin(a_n) \rightarrow 0$. Therefore, by the Algebra of Limits,

 $1-\sin^2(a_n)\longrightarrow 1-0\cdot 0=1.$

Using Theorem 6.32 we have

$$\cos(a_n) = \sqrt{1 - \sin^2(a_n)} \longrightarrow \sqrt{1} = 1,$$

concluding the proof.

Theorem 6.56

Suppose (a_n) is such that $a_n \to 0$ and $a_n \neq 0$. Then,

$$\lim_{n\to\infty}\frac{\sin(a_n)}{a_n}=1\,.$$

Proof

The following elementary trigonometric inequality holds:

$$\sin(x) < x < \tan(x), \quad \forall x \in \left[0, \frac{\pi}{2}\right].$$

Note that $\sin x > 0$ for $0 < x < \pi/2$. Therefore we can divide the above inequality by $\sin(x)$ and take the reciprocals to get

$$\cos(x) < \frac{\sin(x)}{x} < 1$$
, $\forall x \in \left(0, \frac{\pi}{2}\right]$.

If $-\pi/2 < x < 0$, we can apply the above inequality to -x to obtain

$$\cos(-x) < \frac{\sin(-x)}{-x} < 1.$$

Recalling that $\cos(-x) = \cos(x)$ and $\sin(-x) = -\sin(x)$, we get

$$\cos(x) < \frac{\sin(x)}{x} < 1$$
, $\forall x \in \left(-\frac{\pi}{2}, 0\right]$.

Thus

$$\cos(x) < \frac{\sin(x)}{x} < 1, \quad \forall x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \setminus \{0\}.$$
(6.22)

Let

$$\varepsilon := \frac{\pi}{2}$$
.

Since $a_n \to 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n| < \varepsilon = \frac{\pi}{2}, \quad \forall \, n \ge N \, .$$

Since $a_n \neq 0$ by assumption, the above shows that

$$a_n \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \setminus \{0\}, \quad \forall n \ge \mathbb{N}.$$

Therefore we can substitute $x = a_n$ into (6.22) to get

$$\cos(a_n) < \frac{\sin(a_n)}{a_n} < 1, \quad \forall n \ge N$$

We have

 $\cos(a_n) \to 1$

by Theorem 6.55. By the Squeeze Theorem we conclude that

$$\lim_{n\to\infty}\frac{\sin(a_n)}{a_n}=1\,.$$

Warning

You might be tempted to apply L'Hôpital's rule (which we did not cover in these Lecture Notes) to compute

$$\lim_{x \to 0} \frac{\sin(x)}{x}$$

This would yield the correct limit

$$\lim_{x \to 0} \frac{\sin(x)}{x} = \lim_{x \to 0} \frac{(\sin(x))'}{(x)'} = \lim_{x \to 0} \cos(x) = 1.$$

However this is a circular argument, since the derivative of sin(x) at x = 0 is defined as the limit

$$\lim_{x \to 0} \frac{\sin(x)}{x}$$

Theorem 6.57

Suppose (a_n) is such that $a_n \to 0$ and $a_n \neq 0$. Then,

$$\lim_{n \to \infty} \frac{1 - \cos(a_n)}{(a_n)^2} = \frac{1}{2}, \quad \lim_{n \to \infty} \frac{1 - \cos(a_n)}{a_n} = 0.$$

Proof

Step 1. By Theorem 6.55 and Theorem 6.56, we have

$$\cos(a_n) \to 1$$
, $\frac{\sin(a_n)}{a_n} \to 1$.

Therefore

$$\frac{1 - \cos(a_n)}{(a_n)^2} = \frac{1 - \cos(a_n)}{(a_n)^2} \frac{1 + \cos(a_n)}{1 + \cos(a_n)}$$
$$= \frac{1 - \cos^2(a_n)}{(a_n)^2} \frac{1}{1 + \cos(a_n)}$$
$$= \left(\frac{\sin(a_n)}{a_n}\right)^2 \frac{1}{1 + \cos(a_n)} \longrightarrow 1 \cdot \frac{1}{1 + 1} = \frac{1}{2},$$

where in the last line we use the Algebra of Limits. *Step 2.* We have

$$\frac{1-\cos(a_n)}{a_n} = a_n \cdot \frac{1-\cos(a_n)}{(a_n)^2} \longrightarrow 0 \cdot \frac{1}{2} = 0,$$

using Step 1 and the Algebra of Limits.

Example 6.58

Question. Prove that

$$\lim_{n \to \infty} n \sin\left(\frac{1}{n}\right) = 1.$$
(6.23)

Solution. By Theorem 6.56 with $a_n = 1/n$, we get

$$n\sin\left(\frac{1}{n}\right) = \frac{\sin\left(\frac{1}{n}\right)}{\frac{1}{n}} \longrightarrow 1.$$

Example 6.59

Question. Prove that

$$\lim_{n \to \infty} n^2 \left(1 - \cos\left(\frac{1}{n}\right) \right) = \frac{1}{2}.$$
 (6.24)

Solution. By Theorem 6.57 with $a_n = 1/n$, we have

$$n^{2}\left(1-\cos\left(\frac{1}{n}\right)\right) = \frac{1-\cos\left(\frac{1}{n}\right)}{\frac{1}{n^{2}}} \longrightarrow \frac{1}{2}.$$

Example 6.60

Question. Prove that

$$\lim_{n \to \infty} \frac{n\left(1 - \cos\left(\frac{1}{n}\right)\right)}{\sin\left(\frac{1}{n}\right)} = \frac{1}{2}.$$

Solution. Using (6.24)-(6.23) and the Algebra of Limits

$$\frac{n\left(1-\cos\left(\frac{1}{n}\right)\right)}{\sin\left(\frac{1}{n}\right)} = \frac{n^2\left(1-\cos\left(\frac{1}{n}\right)\right)}{n\sin\left(\frac{1}{n}\right)}$$
$$\longrightarrow \frac{1/2}{1} = \frac{1}{2}.$$

Example 6.61

Question. Prove that

$$\lim_{n\to\infty} n\cos\left(\frac{2}{n}\right)\sin\left(\frac{2}{n}\right) = 2.$$

Solution. We have

$$\cos\left(\frac{2}{n}\right) \longrightarrow 1$$
,

by Theorem 6.55 applied with $a_n = 2/n$. Moreover

$$\frac{\sin\left(\frac{2}{n}\right)}{\frac{2}{n}} \longrightarrow 1,$$

by Theorem 6.56 applied with $a_n = 2/n$. Therefore

$$n\cos\left(\frac{2}{n}\right)\sin\left(\frac{2}{n}\right) = 2\cdot\cos\left(\frac{2}{n}\right)\cdot\frac{\sin\left(\frac{2}{n}\right)}{\frac{2}{n}}$$
$$\longrightarrow 2\cdot1\cdot1 = 2,$$

where we used the Algebra of Limits.

Example 6.62

Question. Prove that

$$\lim_{n \to \infty} \frac{n^2 + 1}{n+1} \sin\left(\frac{1}{n}\right) = 1.$$

Solution. Using (6.23) and the Algebra of Limits,

$$\frac{n^2+1}{n+1}\sin\left(\frac{1}{n}\right) = \left(\frac{1+\frac{1}{n^2}}{1+\frac{1}{n}}\right) \cdot \left(n\sin\left(\frac{1}{n}\right)\right)$$
$$\longrightarrow \frac{1+0}{1+0} \cdot 1 = 1.$$

7 Sequences in C

The theory for sequences in \mathbb{C} is very similar to that of sequences in \mathbb{R} . In \mathbb{R} , we said that a sequence (a_n) converges to some number $a \in \mathbb{R}$ if for all $\varepsilon > 0$, it holds

 $|a_n - a| < \varepsilon$

for all *n* sufficiently large. The definition of convergence in \mathbb{C} is essentially the same, with the absolute value replaced by the complex modulus.

Definition 7.1: Sequence of Complex numbers

A sequence a in \mathbb{C} is a function

 $a: \mathbb{N} \to \mathbb{C}$.

For $n \in \mathbb{N}$, we denote the *n*-th element of the sequence *a* by

 $a_n = a(n)$

and write the sequence as

 $(a_n)_{n\in\mathbb{N}}$ or (a_n) .

In the following we define convergent sequences in \mathbb{C} .

Definition 7.2: Convergent sequence in ℂ

We say that a sequence (a_n) in \mathbb{C} **converges** to $a \in \mathbb{C}$, or equivalently has limit a, denoted by

$$\lim_{n\to\infty}a_n=a\quad\text{or}\quad a_n\to a\,,$$

if it holds:

 $\forall \, \varepsilon > 0, \, \exists \, N \in \mathbb{N} \, \text{ s.t. } \, \forall \, n \geq N \, , \, |a_n - a| < \varepsilon \, .$

If there exists $a \in \mathbb{C}$ such that $\lim_{n\to\infty} a_n = a$, we say that the sequence (a_n) is **convergent**.

Important

In Definition 7.2 we still take ε to be real. This makes sense, since

$$|z| = \sqrt{x^2 + y^2} \in \mathbb{R}$$

for all $z = x + iy \in \mathbb{C}$.

Example 7.3

Question. Using Definition 7.2, prove that

$$\lim_{n\to\infty}\frac{(3+i)n-7i}{n}=3+i\,.$$

Solution.

Part 1. Rough Work. Let $\varepsilon > 0$. We need to clarify for which values of *n* the following holds:

$$\left|\frac{(3+i)n-7i}{n}-(3+i)\right|<\varepsilon$$

We have

$$\left|\frac{(3+i)n-7i}{n}-(3+i)\right|=\frac{|-7i|}{n}=\frac{7}{n}.$$

Therefore

$$\frac{7}{n} < \varepsilon \quad \Longleftrightarrow \quad n > \frac{7}{\varepsilon} \,.$$

Part 2. Formal Proof. We want to prove that for all $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

$$\left|\frac{(3+i)n-7i}{n}-(3+i)\right|<\varepsilon\,,\qquad\forall\,n\geq N\,.$$

Let $\varepsilon > 0$. Choose $N \in \mathbb{N}$ such that

$$N > \frac{7}{\varepsilon}$$
.

The above is equivalent to

$$\frac{7}{N} < \varepsilon$$
.

For $n \ge N$ we have

$$\left|\frac{(3+i)n-7i}{n}-(3+i)\right|=\frac{7}{n}\leq\frac{7}{N}<\varepsilon.$$

Boundedness plays an important role for complex sequences.

Definition 7.4: Bounded sequence in \mathbb{C}

A sequence (a_n) in \mathbb{C} is called **bounded** if there exists a constant $M \in \mathbb{R}$, with M > 0, such that

$$|a_n| \le M, \quad \forall n \in \mathbb{N}.$$

As it happens in \mathbb{R} , we have that complex sequences which converge are also bounded.

Theorem 7.5

If a sequence (a_n) in \mathbb{C} converges, then the sequence is bounded.

The proof is identical to the one in \mathbb{R} , and is hence omitted. Similarly to real sequences, we can define divergent complex sequences.

Definition 7.6: Divergent sequences in C

We say that a sequence (a_n) in \mathbb{C} is **divergent** if it is not convergent.

As a corollary of Theorem 7.5 we have the following.

Corollary 7.7

Let (a_n) be a complex sequence. If (a_n) is not bounded, then it is divergent.

7.1 Algebra of limits in ${\mathbb C}$

Most of the results about limits that we have shown in \mathbb{R} also hold in \mathbb{C} . The first result is the Algebra of Limits.

Theorem 7.8: Algebra of limits in \mathbb{C}

Let (a_n) and (b_n) be sequences in \mathbb{C} . Suppose that

$$\lim_{n\to\infty}a_n=a\,,\quad \lim_{n\to\infty}b_n=b\,,$$

for some $a, b \in \mathbb{C}$. Then,

1. Limit of sum is the sum of limits:

$$\lim_{n \to \infty} \left(a_n \pm b_n \right) = a \pm b$$

2. Limit of product is the product of limits:

$$\lim_{n \to \infty} \left(a_n b_n \right) = a b$$

3. If $b_n \neq 0$ for all $n \in \mathbb{N}$ and $b \neq 0$, then

$$\lim_{n \to \infty} \left(\frac{a_n}{b_n} \right) = \frac{a}{b}$$

The proof of Theorem 7.8 follows word by word the proof of the Algebra of Limits for sequences in \mathbb{R} : one just needs to replace the absolute value by the complex modulus.

We can use the Algebra of Limits to compute limits of complex sequences.

Example 7.9

Question. Compute the limit of

$$a_n = \frac{(2-i)n^2 + 6in - 5 - 3i}{(6+3i)n^2 + 11i}$$

Solution. Factor n^2 , the largest power of *n* in the denominator,

$$a_n = \frac{(2-i) + \frac{6i}{n} - \frac{5}{n^2} - \frac{3i}{n^2}}{(6+3i) + \frac{11i}{n^2}} \longrightarrow \frac{2-i}{6+3i}$$

where we used the Algebra of Limits. Finally,

$$\frac{2-i}{6+3i} = \frac{(2-i)(6-3i)}{(6+3i)(6-3i)} = \frac{1}{5} - \frac{4}{15}i$$

7.2 Convergence to zero

One of the results that cannot hold for complex sequences is the Squeeze Theorem. Indeed the chain of inequalities

$$b_n \leq a_n \leq c_n$$

would not make sense in $\mathbb C,$ since there is no order relation.

We can however prove the following (weaker) result.

Theorem 7.10

Let (a_n) be a sequence in \mathbb{C} and suppose that

$$\lim_{n\to\infty}|a_n|=0.$$

Then

$$\lim_{n\to\infty}a_n=0\,.$$

Proof

Assume that $|a_n| \to 0$. We need to show that

 $\forall \, \varepsilon > 0 \,, \, \exists \, N \in \mathbb{N} \, \text{ s.t. } \, \forall \, n \geq N \,, \, \, |a_n - 0| < \varepsilon \,.$

Let $\varepsilon > 0$. Since $|a_n| \to 0$, there exists $N \in \mathbb{N}$ such that

$$||a_n|-0|<\varepsilon,\quad\forall n\geq N.$$

Let $n \ge N$. Then,

$$\begin{aligned} a_n - 0 &| = |a_n| \\ &= |a_n| - 0 \\ &= ||a_n| - 0| \\ &< \varepsilon \,. \end{aligned}$$

Note that the sequence $|a_n|$ is real. Therefore the convergence of $|a_n|$ can be studied using convergence results in \mathbb{R} .

Example 7.11

Question. Prove that $a_n \rightarrow 0$, where

$$a_n = \left(\frac{1}{2} + \frac{1}{3}i\right)^n \,.$$

Solution. We have

$$|a_n| = \left| \left(\frac{1}{2} + \frac{1}{3}i\right)^n \right|$$
$$= \left| \frac{1}{2} + \frac{1}{3}i \right|^n$$
$$= \left(\sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{1}{3}\right)^2} \right)^n$$
$$= \left(\sqrt{\frac{13}{36}} \right)^n.$$

Since

by the Geometric Sequence Test for real sequences, we conclude that

$$|a_n| \to 0$$
.

 $\left| \sqrt{\frac{13}{36}} \right| < 1$,

Hence $a_n \rightarrow 0$ by Theorem 7.10.

Although the Squeeze Theorem cannot be used for complex sequences, sometimes it can be used to deal with real terms in a complex sequence.

Example 7.12

Question. Consider the sequence

$$a_n := \frac{2i\cos(3n)n + (7-i)n^2}{3n^2 + 2in + \sin(2n)}$$

Prove that

$$\lim_{n\to\infty}a_n=\frac{7}{3}-\frac{1}{3}i\,.$$

Solution. We divide by the largest power in the denominator, to get

$$a_n = \frac{\frac{2i\cos(3n)}{n} + (7-i)}{3 + \frac{2i}{n} + \frac{\sin(2n)}{n^2}}.$$

Notice that

$$-1 \le \cos(3n) \le 1$$
, $\forall n \in \mathbb{N}$,

and thus

$$-\frac{2}{n} \leq \frac{2\cos(3n)}{n} \leq \frac{2}{n}, \quad \forall n \in \mathbb{N}.$$

Since

$$-\frac{2}{n} \longrightarrow 0, \quad \frac{2}{n} \longrightarrow 0,$$

by the Squeeze Theorem we conclude that also

$$\frac{2\cos(3n)}{n} \to 0\,.$$

In particular we have shown that

$$\left|\frac{2i\cos(3n)}{n}\right| = \left|\frac{2\cos(3n)}{n}\right| \to 0.$$

Using Theorem 7.10 we infer

$$\frac{2i\cos(3n)}{n} \to 0\,.$$

Similarly,

$$-\frac{1}{n^2} \le \frac{\sin(2n)}{n^2} \le -\frac{1}{n^2}, \quad \forall n \in \mathbb{N}$$

Since

$$-\frac{1}{n^2} \longrightarrow 0, \quad \frac{1}{n^2} \longrightarrow 0,$$

by the Squeeze Theorem we conclude

$$\frac{\sin(2n)}{n^2} \longrightarrow 0$$

Finally, we have

$$\left|\frac{2i}{n}\right| = \frac{2}{n} \longrightarrow 0,$$

 $\frac{2i}{n} \longrightarrow 0$

and therefore

by Theorem 7.10. Using the Algebra of Limits in C we conclude

$$a_n = \frac{\frac{2i\cos(3n)}{n} + (7-i)}{3 + \frac{2i}{n} + \frac{\sin(2n)}{n^2}} \longrightarrow \frac{0 + (7-i)}{3 + 0 + 0} = \frac{7}{3} - \frac{1}{3}i.$$

7.3 Geometric sequence Test and Ratio Test in ${\mathbb C}$

The Geometric Sequence Test and Ratio Test can be generalized to complex sequences.

Theorem 7.13: Geometric sequence Test in \mathbb{C}

Let $x \in \mathbb{C}$ and let $(a_n)_{n \in \mathbb{N}}$ be the geometric sequence in \mathbb{C} defined by

$$a_n := x^n$$
.

We have:

1. If |x| < 1, then

$$\lim_{n\to\infty}a_n=0$$

2. If |x| > 1, then sequence (a_n) is unbounded, and hence divergent.

The proof can be obtained as in the real case, replacing the absolute value by the modulus.

Example 7.14

Question. Prove that $a_n \rightarrow 0$, where

$$a_n = \frac{(-1+4i)^n}{(7+3i)^n}$$

Solution. We first rewrite

$$a_n = \frac{(-1+4i)^n}{(7+3i)^n} = \left(\frac{-1+4i}{7+3i}\right)^n$$

Then, we compute

$$\frac{-1+4i}{7+3i} = \frac{|-1+4i|}{|7+3i|}$$
$$= \frac{\sqrt{(-1)^2+4^2}}{\sqrt{7^2+3^2}}$$
$$= \frac{\sqrt{17}}{\sqrt{58}}$$
$$= \sqrt{\frac{17}{58}}$$
$$< 1$$

By the Geometric Sequence Test $a_n \rightarrow 0$.

Example 7.15

Question. Prove that a_n diverges, where

$$a_n = \frac{(-5+12i)^n}{(3-4i)^n} \,.$$

Solution. We first rewrite

$$a_n = \frac{(-5+12i)^n}{(3-4i)^n} = \left(\frac{-5+12i}{3-4i}\right)^n \,.$$

We compute

$$\left|\frac{-5+12i}{3-4i}\right| = \frac{\left|-5+12i\right|}{\left|3-4i\right|}$$
$$= \frac{\sqrt{5^{2}+(-12)^{2}}}{\sqrt{3^{2}+(-4)^{2}}}$$
$$= \frac{13}{5}$$
$$\ge 1.$$

By the Geometric Sequence Test, the sequence a_n diverges.

Example 7.16

Question. Prove that a_n diverges, where

$$a_n = \exp\left(\frac{i\pi}{2}n\right)$$
.

Solution. We have

$$|a_n| = \left| e^{\frac{i\pi}{2}n} \right| = 1 ,$$

and hence the Geometric Sequence Test cannot be applied. However, we can see that

$$a_n = (i, -1, -i, 1, i, -1, -i, 1, ...),$$

that is, a_n assumes only the values $\{i, -1, -i, 1\}$, and each of them is assumed infinitely many times. Therefore a_n is oscillating, and thus divergent.

We now provide the statement of the Ratio Test in \mathbb{C} .

Theorem 7.17: Ratio Test in ℂ

Let (a_n) be a sequence in \mathbb{C} such that

$$a_n \neq 0$$
, $\forall n \in \mathbb{N}$.

1. Suppose that the following limit exists:

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \, .$$

Then,

• If *L* < 1 we have

$$\lim_{n\to\infty}a_n=0\,.$$

- If L > 1, the sequence (a_n) is unbounded, and hence does not converge.
- 2. Suppose that there exists $N \in N$ and L > 1 such that

$$\left|\frac{a_{n+1}}{a_n}\right| \ge L, \quad \forall \, n \ge N \, .$$

Then the sequence a_n is unbounded, and hence does not converge.

The proof of Theorem 7.17 follows word by word the proof of the Ratio Test Theorem in \mathbb{R} , and only two minor modifications are needed:

- Replace the absolute value with the complex modulus,
- Instead of the Squeeze Theorem, use Theorem 7.10.

Example 7.18

Question. Study the convergence / divergence of the sequence

$$a_n = \frac{(4-3i)^n}{(2n)!}$$

Solution. We compute

$$\begin{aligned} \left| \frac{a_{n+1}}{a_n} \right| &= \left| \frac{(4-3i)^{n+1}}{(2(n+1))!} \frac{(2n)!}{(4-3i)^n} \right| \\ &= \frac{|4-3i|^{n+1}}{|4-3i|^n} \cdot \frac{(2n)!}{(2n+2)!} \\ &= \frac{|4-3i|}{(2n+2)(2n+1)} \\ &= \frac{\sqrt{4^2 + (-3)^2}}{(2n+2)(2n+1)} \\ &= \frac{5}{(2n+2)(2n+1)} \\ &= \frac{5}{(2n+2)(2n+1)} \\ &= \frac{5}{n^2} \\ &= \frac{5}{n^2} \\ &\xrightarrow{\left(2+\frac{2}{n}\right)\left(2+\frac{1}{n}\right)} \longrightarrow L = 0 \,. \end{aligned}$$

Since L = 0 < 1, by the Ratio Test in \mathbb{C} we infer $a_n \to 0$.

7.4 Convergence of real and imaginary part

A complex number $z \in \mathbb{C}$ can be written as

$$z = a + bi$$

for some $a, b \in \mathbb{R}$, where

- a is the real part of z,
- *b* the imaginary part of *z*.

We can prove that a complex sequence converges if and only if both the real parts and the imaginary parts converge.

Theorem 7.19

Let $(z_n)_{n \in \mathbb{N}}$ be a sequence in \mathbb{C} . For $n \in \mathbb{N}$, let $a_n, b_n \in \mathbb{R}$ such that

$$z_n = a_n + b_n i.$$

Let z = a + bi, with $a, b \in \mathbb{R}$. Then

$$\lim_{n\to\infty} z_n = z \quad \iff \quad \lim_{n\to\infty} a_n = a \,, \quad \lim_{n\to\infty} b_n = b.$$

Proof

Part 1. Suppose that

$$\lim_{n\to\infty} z_n = z \, .$$

To prove that $a_n \rightarrow a$ we need to show that

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ s.t. } \forall n \ge N, |a_n - a| < \varepsilon.$$

Let $\varepsilon > 0$. Since $z_n \to z$, there exists $N \in \mathbb{N}$ such that

$$|z_n-z|<\varepsilon\,,\quad\forall\,n\geq N\,.$$

Let $n \ge N$. Then

$$|a_n - a| = \sqrt{(a_n - a)^2}$$

$$\leq \sqrt{(a_n - a)^2 + (b_n - b)^2}$$

$$= |(a_n - a) + (b_n - b)i|$$

$$= |(a_n + b_n i) - (a + bi)|$$

$$= |z_n - z|$$

$$\leq \varepsilon.$$

The proof for $b_n \rightarrow b$ is similar. *Part 2.* Suppose that

$$\lim_{n\to\infty}a_n=a\,,\quad \lim_{n\to\infty}b_n=b\,.$$

To prove that $z_n \rightarrow z$ we need to show that

 $\forall\,\varepsilon>0\,,\,\exists\,N\in\mathbb{N}\;\;\mathrm{s.t.}\;\;\forall\,n\geq N\,,\;\;|z_n-z|<\varepsilon\,.$

Let $\varepsilon > 0$. Since $a_n \to a$, there exists $N_1 \in \mathbb{N}$ such that

$$|a_n-a|<rac{arepsilon}{2}\,,\quad \forall\,n\geq N_1\,.$$

Since $b_n \to b$, there exists $N_2 \in \mathbb{N}$ such that

$$|b_n - b| < \frac{\varepsilon}{2} \,, \quad \forall \, n \ge N_2 \,.$$

Let

$$N := \max\{N_1, N_2\}$$
.

Let $n \ge N$. By the triangle inequality,

$$\begin{aligned} |z_n - z| &= |(a_n + b_n i) - (a + bi)| \\ &= |(a_n - a) + (b_n - b)i| \\ &\leq |a_n - a| + |b_n - b| \cdot |i| \\ &= |a_n - a| + |b_n - b| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \,. \end{aligned}$$

Example 7.20

Question. Consider the complex sequence

$$z_n := \frac{\left(4n + 3n^2i\right)\left(2n^2 + i\right)}{5n^4}.$$

Show that

$$\lim_{n\to\infty} z_n = \frac{6}{5}i.$$

Solution. We find the real and imaginary parts of z_n

$$z_n = \frac{(4n + 3n^2i)(2n^2 + i)}{5n^4}$$

= $\frac{8n^3 + 4ni + 6n^4i + 3n^2i^2}{5n^4}$
= $\frac{8n^3 - 3n^2}{5n^4} + \frac{6n^4 + 4n}{5n^4}i$
= $a_n + b_ni$.

Using the Algebra of Limits for real sequences we have that

$$a_n = \frac{8n^3 - 3n^2}{5n^4} = \frac{\frac{8}{n} - \frac{3}{n^2}}{5} \longrightarrow \frac{0 - 0}{5} = 0,$$

$$b_n = \frac{6n^4 + 4n}{5n^4} = \frac{6 + \frac{4}{n^3}}{5} \longrightarrow \frac{6 + 0}{5} = \frac{6}{5}.$$

By Theorem 7.19 we conclude

$$\lim_{n \to \infty} z_n = \lim_{n \to \infty} a_n + i \lim_{n \to \infty} b_n = 0 + \frac{6}{5}i = \frac{6}{5}i.$$

8 Series

A series is the sum of all terms in a sequence (a_n) :

$$a_1 + a_2 + a_3 + \dots + a_n + \dots$$

Since we are dealing with infinitely many terms, we need to be careful. For example, consider the series

$$\sum_{n=1}^{\infty} (-1)^n = -1 + 1 - 1 + 1 - 1 + 1 - \dots$$
(8.1)

If we sum the terms in pairs, we obtain

$$\sum_{n=1}^{\infty} (-1)^n = (-1+1) + (-1+1) + (-1+1) + \dots = 0.$$

If we reorder the terms, we obtain a different result

$$\sum_{n=1}^{\infty} (-1)^n = (1+1) - 1 + (1+1) - 1 + \dots$$
$$= (2-1) + (2-1) + \dots$$
$$= 1+1+\dots$$
$$= \infty.$$

We can also swap terms pairwise, and then sum each pair, starting from the second term. This way we obtain

$$\sum_{k=1}^{\infty} (-1)^n = (-1+1) + (-1+1) + (-1+1) + \dots$$
$$= (1-1) + (1-1) + (1-1) + \dots$$
$$= 1 + (-1+1) + (-1+1) + (-1+1) + \dots$$
$$= 1.$$

If we do one more swap, and start summing each pair starting from the third term, we get

$$\sum_{k=1}^{\infty} (-1)^n = 1 + (-1+1) + (-1+1) + (-1+1) + \dots$$
$$= 1 + (1-1) + (1-1) + (1-1) + \dots$$
$$= 1 + 1 + (-1+1) + (-1+1) + \dots$$
$$= 2.$$

With similar arguments, we see that we can rearrange terms so that

$$\sum_{n=1}^{\infty} (-1)^n = m \,,$$

for each $m \in \mathbb{Z}$. This example shows that commutativity of the sum does not hold when summing infinitely many terms: The result of the sum depends on the order in which we sum. This means we need a good definition of convergence for series.

8.1 Convergent series

We will develop the theory of series for sequences in \mathbb{C} . All the results for complex series will also hold for series in \mathbb{R} . Some results will only hold for series in \mathbb{R} : this will be the case for comparison tests in which the order relation of \mathbb{R} is needed.

We start by defining partial sums.

Definition 8.1: Partial sums

Let (a_n) be a sequence in \mathbb{C} . The *k*-th partial sum of (a_n) is

$$s_k := a_1 + a_2 + \dots + a_k = \sum_{n=1}^k a_n$$

This sequence $(s_k)_{k \in \mathbb{N}}$ is called the sequence of **partial sums**.

We can use the sequence of partial sums to define convergence of a series.

Definition 8.2: Convergent series

Let (a_n) be a sequence in \mathbb{C} . We denote the series of $(a_n)_{n \in \mathbb{N}}$ by

$$\sum_{n=1}^{\infty} a_n$$

We say that this series **converges** to $s \in \mathbb{C}$ if

$$\lim_{k\to\infty}\sum_{n=1}^k a_n = \lim_{k\to\infty} s_k = s\,.$$

In this case we write

$$\sum_{n=1}^{\infty} a_n = s$$

Definition 8.3: Divergent series

Let (a_n) be a sequence in \mathbb{C} . The series

$$\sum_{n=1}^{\infty} a_n$$

is **divergent** if the sequence of partial sums (s_k) is divergent.

Example 8.4

Question. Prove that

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1.$$

Solution. The idea to prove convergence is to split the general term into the sum of two fraction:

$$\frac{1}{n(n+1)} = \frac{A}{n} + \frac{B}{n(n+1)}$$
$$= \frac{A(n+1) + Bn}{n(n+1)}$$
$$= \frac{(A+B)n + A}{n(n+1)}$$

In order for the LHS and RHS to be the same, we need to impose

$$(A+B)n+A=1,$$

which holds if and only if

$$A + B = 1$$
, $A = 1 \implies A = 1$, $B = -1$.

Therefore, we conclude that

$$\frac{1}{n(n+1)} = \frac{1}{n} - \frac{1}{n+1} \, .$$

We can now compute the partial sums s_k as follows:

$$s_k = \sum_{n=1}^k \frac{1}{n(n+1)}$$

= $\sum_{n=1}^k \left(\frac{1}{n} - \frac{1}{n+1}\right)$
= $\frac{1}{1} - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \frac{1}{3} - \frac{1}{4} + \dots + \frac{1}{k} - \frac{1}{k+1}$
= $1 - \frac{1}{k+1}$.

Therefore,

$$\lim_{k\to\infty}s_k=\lim_{k\to\infty}\left(1-\frac{1}{k+1}\right)=1\,,$$

which means that the series converges to 1, that is,

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1.$$

A series of this kind is called a **telescopic series**, since we can *fold* the entire partial sum together, in such a way that only two terms remain.

Let us go back to the series considered in (8.1).

Example 8.5

Question. Prove that the following series diverges

$$\sum_{n=1}^{\infty} (-1)^n \, .$$

Solution. The partial sums s_k are given by

 $s_k = \sum_{n=1}^k (-1)^n = \begin{cases} -1 & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even.} \end{cases}$

Therefore s_k diverges, so also the series $\sum (-1)^n$ diverges.

In general, it is a difficult taks to compute the exact sum of a series. Therefore, we will mainly focus our effort on determining whether a series converges or not.

The following Theorem shows that if the terms in the sequence do not converge to 0, then the series cannot converge.

Theorem 8.6: Necessary Condition for Convergence

Let (a_n) be a sequence in \mathbb{C} . If the series

$$\sum_{n=1}^{\infty} a_n$$

converges, then

 $\lim_{n\to\infty}a_n=0\,.$

Proof

Suppose that

converges. By definition of convergent series there exists some $s \in \mathbb{C}$ such that

 $\lim_{k\to\infty}s_k=\lim_{k\to\infty}\sum_{n=1}^k a_n=s\,.$

 $\sum_{n=1}^{\infty} a_n$

Then also

 $\lim_{k\to\infty}s_{k-1}=s\,.$

Hence, by the Algebra of Limits in C, we have that

 $\lim_{k \to \infty} (s_k - s_{k-1}) = \lim_{k \to \infty} s_k - \lim_{k \to \infty} s_{k-1} = s - s = 0.$

Noting that

 $s_k - s_{k-1} = a_k$, $\forall k \in \mathbb{N}$,

we obtain

$$\lim_{k\to\infty}a_k=\lim_{k\to\infty}(s_k-s_{k-1})=0\,.$$

Important

Theorem 8.6 is saying that, if

$$\lim_{n\to\infty}a_n\neq 0\,,$$

then the series

$$\sum_{n=1}^{\infty} a_n$$

does not converge.

Example 8.7

Consider the series

 $\sum_{n=1}^{\infty} (-1)^n \,. \tag{8.2}$

We have that

$$\lim_{n \to \infty} a_n = \lim_{n \to \infty} (-1)^n \neq 0$$

being (a_n) divergent. Therefore the series at (8.2) diverges by Theorem 8.6.

Example 8.8

Question. Discuss converge/divergence for the following series

$$\sum_{n=1}^{\infty} \frac{n}{5n+11}$$

Solution. We have

$$a_n := \frac{n}{5n+11} = \frac{1}{5+\frac{11}{n}} \longrightarrow \frac{1}{5} \neq 0$$

Hence, the series $\sum a_n$ diverges.

Important

Theorem 8.6 says that if $\sum_{n=1}^{\infty} a_n$ converges, then

 $a_n \rightarrow 0$.

The converse is false: In general the condition $a_n \rightarrow 0$ does not guarantee convergence of the associated series, as shown in the example below.

Example 8.9

Question. Discuss convergence/divergence for the following series

$$\sum_{n=1}^{\infty} a_n \,, \quad a_n \,:= \frac{1}{\sqrt{n+1} + \sqrt{n}}$$

Solution. By the Algebra of Limits we have

$$\lim_{n\to\infty}a_n=0\,.$$

Therefore, we cannot conclude anything yet: The series might converge or diverge. Let us

compute the partial sums:

$$s_k = \sum_{n=1}^k \frac{1}{\sqrt{k+1} + \sqrt{k}}$$

= $\sum_{n=1}^k \frac{1}{\sqrt{k+1} + \sqrt{k}} \cdot \frac{\sqrt{k+1} - \sqrt{k}}{\sqrt{k+1} - \sqrt{k}}$
= $\sum_{n=1}^k \sqrt{k+1} - \sqrt{k}$
= $\sqrt{2} - \sqrt{1} + \sqrt{3} - \sqrt{2} + \dots + \sqrt{k+1} - \sqrt{k}$
= $\sqrt{k+1} - 1$.

We have shown that the partial sums are

$$s_k = \sum_{n=1}^k a_n = \sqrt{k+1} - 1.$$

Therefore (s_k) is divergent, and so the series $\sum a_n$ is divergent.

Remark 8.10

It is customary to sum a series starting at n = 1. However one could start the sum at any n = N with $N \in \mathbb{N}$. This does not affect the convergence of the series, in the sense that

$$\sum_{n=1}^{\infty} a_n \text{ converges } \iff \sum_{n=N}^{\infty} a_n \text{ converges.}$$

In case of convergence, we would of course have

$$\sum_{n=N}^{\infty} a_n = \sum_{n=1}^{\infty} a_n - (a_1 + \dots + a_{N-1}) \; .$$

Example 8.11

Question. Prove that

$$\sum_{n=7}^{\infty} \frac{1}{n(n+1)} = \frac{1}{7} \, .$$

Solution. We have seen that

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1.$$

Hence also the series

$$\sum_{n=7}^{\infty} \frac{1}{n(n+1)}$$

converges. In this case, the partial sums are given by

$$\begin{split} s_k &= \sum_{n=7}^k \frac{1}{n(n+1)} \\ &= \sum_{n=7}^k \left(\frac{1}{n} - \frac{1}{n+1} \right) \\ &= \frac{1}{7} - \frac{1}{8} + \frac{1}{8} - \frac{1}{9} + \ldots + \frac{1}{k} - \frac{1}{k+1} \\ &= \frac{1}{7} - \frac{1}{k+1} \,. \end{split}$$

Therefore

$$\sum_{n=7}^{\infty} \frac{1}{n(n+1)} = \lim_{k \to \infty} s_k = \frac{1}{7} \, .$$

8.2 Geometric series

Definition 8.12: Geometric Series in ℂ

Let $x \in \mathbb{C}$. The **geometric series** of ratio x is the series

$$\sum_{n=0}^{\infty} x^n \, .$$

Geometric series are one of the few types of series that can be explicitly computed, as stated in the following theorem.

Theorem 8.13: Geometric Series Test

Let $x \in \mathbb{C}$. We have:

1. If |x| < 1, then the geometric series of ratio *x* converges, with

$$\sum_{n=0}^{\infty} x^n = \frac{1}{1-x} \,. \tag{8.3}$$

2. If $|x| \ge 1$, then the geometric series of ratio *x* diverges.

Important

Recall that the Geometric Sequence Test does not cover the case |x| = 1, since in general the sequence

$$a_n = x^n$$

could be convergent or divergent. However, the Geometric Series Test does cover the case |x| = 1, in which case

$$\sum_{n=0}^{\infty} x^n$$

diverges.

Proof

Part 1. Suppose that |x| < 1. Let us compute the partial sums:

$$s_{k} = \sum_{n=0}^{k} x^{n}$$

$$= \sum_{n=0}^{k} x^{n} \cdot \frac{1-x}{1-x}$$

$$= \sum_{n=0}^{k} \frac{x^{n} - x^{n+1}}{1-x}$$

$$= \frac{1-x}{1-x} + \frac{x-x^{2}}{1-x} + \dots + \frac{x^{n} - x^{n+1}}{1-x}$$

$$= \frac{1-x^{k+1}}{1-x}.$$

We have therefore shown that

$$s_k = \sum_{n=0}^k x^n = \frac{1 - x^{k+1}}{1 - x}, \quad \forall k \in \mathbb{N}.$$
 (8.4)

Notice that (8.4) could have also been proven by induction:

1. Base case: For k = 0, we get that

$$s_0 = x^0 = 1 = \frac{1 - x^1}{1 - x}$$
,

showing that (8.4) holds.

2. Induction step: Let $k \in \mathbb{N} \cup \{0\}$ and suppose that

$$s_k = \frac{1 - x^{k+1}}{1 - x}$$

Then,

$$s_{k+1} = s_k + x^{k+1}$$

= $\frac{1 - x^{k+1}}{1 - x} + x^{k+1}$
= $\frac{1 - x^{k+1} + (1 - x)x^{k+1}}{1 - x}$
= $\frac{1 - x^{k+1} + x^{k+1} - x^{k+2}}{1 - x}$
= $\frac{1 - x^{k+2}}{1 - x}$,

concluding the proof of the inductive step.

By the Principle of Induction, formula (8.4) holds for all $k \in \mathbb{N}$. Since |x| < 1, by the Geometric Sequence Test we infer

$$\lim_{k\to\infty} x^k = 0.$$

Hence

$$\sum_{n=0}^{\infty} x^n = \lim_{k \to \infty} s_k$$
$$= \lim_{k \to \infty} \frac{1 - x^{k+1}}{1 - x}$$
$$= \lim_{k \to \infty} \frac{1 - x \cdot x^k}{1 - x}$$
$$= \frac{1}{1 - x},$$

where the last equality follows from the Algebra of Limits. *Part 2.* Suppose $|x| \ge 1$. Then

$$\lim_{n \to \infty} x^n \neq 0.$$
 (8.5)

Indeed, suppose by contradiction that $x_n \to 0$. Hence, for $\varepsilon = 1/2$, there exists $N \in \mathbb{N}$ such that

$$|x^n - 0| < \varepsilon = \frac{1}{2}, \quad \forall n \ge \mathbb{N}$$

However

$$|x^n - 0| = |x^n| = |x|^n \ge 1$$
,

which yields

$$1<\varepsilon=\frac{1}{2}\,,$$

contradiction. Then (8.5) holds and the series

$$\sum_{n=0}^{\infty} x^n$$

diverges by the Necessary Condition in Theorem 8.6.

Let us apply the Geometric Series Test of Theorem 8.13 to some examples.

Example 8.14

Question. Discuss convergence/divergence of the following series. If the series converges, compute the limit.

$$\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n, \qquad \qquad \sum_{n=0}^{\infty} \left(-\frac{3}{2}\right)^n,$$
$$\sum_{n=0}^{\infty} \left(\frac{-3}{4}\right)^n, \qquad \qquad \sum_{n=0}^{\infty} (-1)^n.$$

Solution.

1. Since $\left|\frac{1}{2}\right| < 1$, by the GST we have

$$\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n = \frac{1}{1 - \frac{1}{2}} = 2$$

2. Since $\left|\frac{-3}{2}\right| = \frac{3}{2} > 1$, by the GST the series

$$\sum_{n=0}^{\infty} \left(-\frac{3}{2}\right)^n$$

diverges.

3. Since $\left|\frac{-3}{4}\right| = \frac{3}{4} < 1$, we have

$$\sum_{n=0}^{\infty} \left(\frac{-3}{4}\right)^n = \frac{1}{1 - \frac{-3}{4}} = \frac{1}{\frac{7}{4}} = \frac{4}{7}$$

4. Since |-1| = 1, the series

$$\sum_{n=0}^{\infty} (-1)^n$$

diverges.

Remark 8.15

If the sum of a Geometric Sries does not start at n = 0, we need to tweak the summation formula at (8.3). For example, if |x| < 1, and we start the series at n = 1, we get

$$\sum_{n=1}^{\infty} x^k = \frac{1}{1-x} - 1 = \frac{x}{1-x}$$

Example 8.16

Question. Prove that

$$\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n = 1$$

Solution. We have that

$$\sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^n = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n - 1$$
$$= \frac{1}{1 - \frac{1}{2}} - 1 = 1.$$
The Geometric Series Test of Theorem 8.13 can be applied to complex geometric series as well.

Example 8.17

Question. Discuss convergence/divergence of the following series. If the series converges, compute the limit.

$$\sum_{n=0}^{\infty} \frac{1}{(1+i)^n}, \quad \sum_{n=0}^{\infty} \left(\frac{1-5i}{3+3i}\right)^n, \quad \sum_{n=0}^{\infty} \left(\frac{2+i}{3-2i}\right)^n.$$

Solution.

1. We have

$$\frac{1}{(1+i)^n} = \left(\frac{1}{1+i}\right)^n$$

and

$$\left|\frac{1}{1+i}\right| = \frac{1}{\sqrt{1^2 + 1^2}} = \frac{1}{\sqrt{2}} < 1 \,.$$

Therefore, the series converges by the Geometric Series Test, and

$$\sum_{n=0}^{\infty} \frac{1}{(1+i)^n} = \frac{1}{1-\frac{1}{1+i}} = 1-i.$$

2. Since

$$\begin{split} \left| \frac{1-5i}{3+3i} \right| &= \frac{|1-5i|}{|3+3i|} \\ &= \frac{\sqrt{(1)^2 + (-5)^2}}{3\sqrt{1^2 + 1^2}} \\ &= \frac{\sqrt{26}}{3\sqrt{2}} \\ &= \frac{\sqrt{13}}{3} > 1 \,, \end{split}$$

the series diverges by the Geometric Series Test.

3. We have

$$\begin{aligned} \left|\frac{2+i}{3-2i}\right| &= \frac{|2+i|}{|3-2i|} \\ &= \frac{\sqrt{2^2+1^2}}{\sqrt{3^2+(-2)^2}} \\ &= \sqrt{\frac{5}{13}} < 1 \,. \end{aligned}$$

Therefore the series converges by the Geometric Series Test, and

$$\sum_{n=0}^{\infty} \left(\frac{2+i}{3-2i}\right)^n = \frac{1}{1-\frac{2+i}{3-2i}}$$
$$= \frac{1}{\frac{3-2i-(2+i)}{3-2i}}$$
$$= \frac{3-2i}{1-3i}$$
$$= \frac{3-2i}{1-3i} \frac{1+3i}{1+3i}$$
$$= \frac{3-2i+9i-6i^2}{1-9i^2}$$
$$= \frac{9}{10} + \frac{7}{10}i$$

8.3 Algebra of Limits for Series

We have proven the Algebra of Limit Theorem for sequences in \mathbb{C} . A similar results holds for series as well.

Theorem 8.18: Algebra of Limits for Series

Let $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be sequences in \mathbb{C} and let $c \in \mathbb{C}$. Suppose that

1

$$\sum_{n=1}^{\infty} a_n = a, \qquad \sum_{n=1}^{\infty} b_n = b.$$

Then:

1. The sum of series is the series of the sums:

$$\sum_{n=1}^{\infty} \left(a_n \pm b_n \right) = a \pm b \, .$$

2. The product of a series with a number obeys

$$\sum_{n=1}^{\infty} c \cdot a_n = c \cdot a \, .$$

Proof

Part 1. We prove the formula with the + sign, since in the other case the proof is the same. To this end, define the partial sums

$$s_k := \sum_{n=1}^k a_n$$
, $t_k := \sum_{n=1}^k b_n$, $v_k := \sum_{n=1}^k (a_n + b_n)$.

We can write

$$v_k = \sum_{n=1}^k (a_n + b_n)$$

= $(a_1 + b_1) + \dots + (a_k + b_k)$
= $(a_1 + \dots + a_k) + (b_1 + \dots + b_k)$
= $s_k + t_k$.

By assumption $s_k \to a$ and $t_k \to b$. Hence, by the Algebra of Limits in \mathbb{C} , we infer

$$\sum_{n=1}^{\infty} (a_n + b_n) = \lim_{k \to \infty} v_k$$
$$= \lim_{k \to \infty} (s_k + t_k)$$
$$= \lim_{k \to \infty} s_k + \lim_{k \to \infty} t_k$$
$$= a + b.$$

Part 2. Denote the partial sums by

$$s_k := \sum_{n=1}^k a_n$$
, $t_k := \sum_{n=1}^k c \cdot a_n$.

We can write

$$t_k = \sum_{n=1}^k c \cdot a_k$$

= $c \cdot a_1 + c \cdot a_2 + \dots + c \cdot a_k$
= $c \cdot (a_1 + a_2 + \dots + a_k) =$
= $c \cdot s_k$

By assumption $s_k \rightarrow a$, so that the Algebra of Limits in \mathbb{C} allows to conclude

$$\sum_{n=1}^{\infty} c \cdot a_n = \lim_{k \to \infty} t_k$$
$$= \lim_{k \to \infty} c \cdot s_k$$
$$= c \cdot a.$$

Let us apply the Algebra of Limits for Series to a concrete example.

Example 8.19

Question. Prove that

$$\sum_{n=0}^{\infty} \left(2\left(\frac{1}{3}\right)^n + \left(\frac{2}{3}\right)^n \right) = 6.$$

Solution. Note that

$$\sum_{n=0}^{\infty} \left(\frac{1}{3}\right)^n = \frac{1}{1-\frac{1}{3}} = \frac{3}{2},$$
$$\sum_{n=0}^{\infty} \left(\frac{2}{3}\right)^n = \frac{1}{1-\frac{2}{3}} = 3,$$

by the Geometric Series Test. Therefore, we can apply the Algebra of Limit for Series to conclude that

$$\sum_{n=0}^{\infty} \left(2\left(\frac{1}{3}\right)^n + \left(\frac{2}{3}\right)^n \right) = 2 \cdot \sum_{n=0}^{\infty} \left(\frac{1}{3}\right)^n + \sum_{n=0}^{\infty} \left(\frac{2}{3}\right)^n$$
$$= 2 \cdot \frac{3}{2} + 3 = 6$$

Important

The Algebra of Limits Theorem 8.18 does not discuss product and quotient of series. The situation becomes more complicated in this case: Indeed, we have

$$(a_1 + a_2) \cdot (a_2 + b_2) = a_1 b_1 + a_2 b_2 + a_1 b_2 + a_2 b_1.$$

Therefore, in general, we can expect

$$\left(\sum_{n=0}^{\infty} a_n\right) \cdot \left(\sum_{n=0}^{\infty} b_n\right) \neq \sum_{n=0}^{\infty} a_n \cdot b_n \,. \tag{8.6}$$

A way to compute

$$\left(\sum_{n=0}^{\infty} a_n\right) \cdot \left(\sum_{n=0}^{\infty} b_n\right)$$

is through the so-called **Cauchy Product** of two series. We do not cover the latter, and the interested reader can refer to Page 82 in [1].

Similarly, we expect that

$$\frac{\sum_{n=0}^{n}a_n}{\sum_{n=0}^{\infty}b_n} \neq \sum_{n=0}^{\infty}\frac{a_n}{b_n}.$$
(8.7)

Let us give two examples to show that formulas (8.6) and (8.7) hold.

 ∞

Example 8.20

1. We know that

$$\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n = \frac{1}{1-\frac{1}{2}} = 2.$$
(8.8)

Therefore

$$\left(\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n\right) \cdot \left(\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n\right) = 2 \cdot 2 = 4.$$

However

$$\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \cdot \left(\frac{1}{2}\right)^n = \sum_{n=0}^{\infty} \left(\frac{1}{4}\right)^n = \frac{1}{1-\frac{1}{4}} = \frac{4}{3}.$$

Hence

$$\left(\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n\right) \cdot \left(\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n\right) \neq \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \cdot \left(\frac{1}{2}\right)^n.$$

2. Using (8.8) we have

$$\frac{\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n}{\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n} = \frac{2}{2} = 1.$$

On the other hand

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)^n}{\left(\frac{1}{2}\right)^n} = \sum_{n=0}^{\infty} 1,$$

which does not converge by the Necessary Condition, since 1 does not converge to 0. Therefore

$$\frac{\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n}{\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n} \neq \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)^n}{\left(\frac{1}{2}\right)^n}.$$

8.4 Non-negative series

We now investigate series of which all terms are non-negative. To be precise:

 Definition 8.21: Non-negative series

 Let (a_n) be a sequence in \mathbb{R} . We call the series

 $\sum_{n=1}^{\infty} a_n$

 a non-negative series if

 $a_n \ge 0$, $\forall n \in \mathbb{N}$.

The key remark for non-negative series is that the partial sums are increasing.

Lemma 8.22

Let (a_n) be a sequence in \mathbb{R} with

$$a_n \ge 0$$
, $\forall n \in \mathbb{N}$.

Define the partial sums as

$$s_k := \sum_{n=1}^k a_n \, .$$

The sequence (s_k) is increasing.

Proof

For all $k \in \mathbb{N}$ we have

$$s_{k+1} = \sum_{n=1}^{k+1} a_n = s_k + a_{k+1} \ge s_k$$

where we used that $a_{k+1} \ge 0$. Therefore (s_k) is increasing.

Therefore, if we have a series with non-negative terms

$$\sum_{n=1}^{\infty} a_n$$

there are only 2 options:

1.
$$\sum_{n=1}^{\infty} a_n$$
 converges,

2. $\sum_{n=1}^{\infty} a_n$ diverges to $+\infty$.

This is because the partial sums (s_k) are increasing. Therefore we have either:

- 1. (s_k) is bounded above: Then s_k converges by the Monotone Convergence Theorem
- 2. (s_k) is not bounded above: Therefore s_k diverges to $+\infty$.

We present 4 test for the convergence of non-negative series:

- 1. Cauchy Condensation Test
- 2. Comparison Test
- 3. Limit Comparison Test
- 4. Ratio Test (positive series only)

8.4.1 Cauchy Condensation Test

Let us start with the study of the two non-negative series:

$$\sum_{n=1}^{\infty} \frac{1}{n^2} , \quad \sum_{n=1}^{\infty} \frac{1}{n} .$$

Question 8.23

Do the above series converge or diverge?

Answer: the first series converges, while the second diverges. We prove it in the next two theorems.

Theorem 8.24

The following series converges

$$\sum_{n=1}^{\infty} \frac{1}{n^2}.$$

Proof

For $k \in \mathbb{N}$ define the sequence of partial sums

$$s_k := \sum_{n=1}^k \frac{1}{n^2}.$$

Note that

$$\begin{split} s_k &= 1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots + \frac{1}{k^2} \\ &= 1 + \frac{1}{2 \cdot 2} + \frac{1}{3 \cdot 3} + \frac{1}{4 \cdot 4} + \dots + \frac{1}{k \cdot k} \\ &< 1 + \frac{1}{2 \cdot 1} + \frac{1}{3 \cdot 2} + \frac{1}{4 \cdot 3} + \dots + \frac{1}{k \cdot (k-1)} \\ &= 1 + \left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \dots + \left(\frac{1}{k-1} - \frac{1}{k}\right) \\ &= 1 + 1 - \frac{1}{k} \\ &= 2 - \frac{1}{k} \\ &\leq 2 . \end{split}$$

showing that s_k is bounded above. Recall that s_k is increasing, by Lemma 8.22. Therefore, by the Monotone Convergence Theorem, we conclude that s_k converges. Hence the series

$$\sum_{n=1}^{\infty} \frac{1}{n^2}$$

is convergent.

Theorem 8.25: Harmonic series

The harmonic series

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

is divergent.

Proof

For $k \in \mathbb{N}$ define the sequence of partial sums

Note that

while

$$s_k := \sum_{n=1}^k \frac{1}{n} \, .$$

$$s_2 = 1 + \frac{1}{2}$$

$$s_{4} = 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right)$$

> $1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right)$
= $1 + \frac{1}{2} + 2\left(\frac{1}{4}\right)$
= $1 + \frac{1}{2} + \frac{1}{2}$
= $1 + 2\left(\frac{1}{2}\right)$

Similarly

$$\begin{split} s_4 &= 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right) \\ &> 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\right) \\ &= 1 + \frac{1}{2} + 2\left(\frac{1}{4}\right) + 4\left(\frac{1}{8}\right) \\ &= 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} \\ &= 1 + 3\left(\frac{1}{2}\right) \end{split}$$

Proceeding in a similar way, for $k \in \mathbb{N}$ we have

$$\begin{split} s_{2^k} &= 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4}\right) + \left(\frac{1}{5} + \frac{1}{6} + \frac{1}{7} + \frac{1}{8}\right) \\ &+ \dots + \left(\frac{1}{2^{k-1} + 1} + \dots + \frac{1}{2^k}\right) \\ &> 1 + \frac{1}{2} + \left(\frac{1}{4} + \frac{1}{4}\right) + \left(\frac{1}{8} + \frac{1}{8} + \frac{1}{8} + \frac{1}{8}\right) \\ &+ \dots + \left(\frac{1}{2^k} + \dots + \frac{1}{2^k}\right) \\ &= 1 + \frac{1}{2} + 2\left(\frac{1}{4}\right) + 4\left(\frac{1}{8}\right) + \dots + 2^{k-1}\left(\frac{1}{2^k}\right) \\ &= 1 + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \dots + \frac{1}{2} \\ &= 1 + k\left(\frac{1}{2}\right) \end{split}$$

Hence

$$s_{2^k} > 1 + k\left(\frac{1}{2}\right), \quad \forall k \in \mathbb{N},$$

showing that s_{2^k} is unbounded. Therefore s_k is unbounded, and s_k does not converge. We conclude that the series

$$\sum_{n=1}^{n} \frac{1}{n}$$

is divergent.

The proofs of the above theorems inspire the Cauchy Condensation Test.

Theorem 8.26: Cauchy Condensation Test

Let (a_n) be a sequence in \mathbb{R} . Suppose that (a_n) is non-negative and decreasing, that is,

$$a_n \ge a_{n+1}$$
, $\forall n \in \mathbb{N}$.

They are equivalent:

1. The following series converges

$$\sum_{n=1}^{\infty} a_n$$

2. The following series converges

$$\sum_{n=0}^{\infty} 2^n a_{2^n} = a_1 + 2a_2 + 8a_8 + 16a_{16} + \dots$$

Proof

For $k \in \mathbb{N}$ denote the partial sums by

$$s_k := \sum_{n=1}^k a_n, \quad t_k := \sum_{n=0}^k 2^n a_{2^n}.$$

Since $a_n \ge 0$, it is immediate to check that the sequences (s_k) and (t_k) are increasing. *Part 1.* Assume that the series

$$\sum_{n=0}^{\infty} 2^n a_{2^n}$$

diverges. Hence the sequence (t_k) diverges and therefore (t_k) is not bounded above.

Indeed, suppose (t_k) was bounded above. Since (t_k) is increasing, we would conclude that (t_k) is convergent by the Monotone Convergence Theorem. Contradiction.

We want to estimate s_k from below. To this end, we notice that

$$s_{2} = a_{1} + a_{2}$$

$$\geq \frac{1}{2}a_{1} + a_{2}$$

$$= \frac{1}{2}(a_{1} + 2a_{2})$$

$$= \frac{1}{2}t_{1},$$

where we used that $a_n \ge 0$, and so $a_1 > a_1/2$. Moreover

$$s_{4} = a_{1} + a_{2} + (a_{3} + a_{4})$$

$$\geq a_{1} + a_{2} + (a_{4} + a_{4})$$

$$= a_{1} + a_{2} + 2a_{4}$$

$$\geq \frac{1}{2} a_{1} + a_{2} + 2a_{4}$$

$$= \frac{1}{2} (a_{1} + 2a_{2} + 4a_{4})$$

$$= \frac{1}{2} t_{2},$$

where we used also that (a_n) is decreasing. Similar reasoning yields

$$s_8 = a_1 + a_2 + (a_3 + a_4) + (a_5 + a_6 + a_7 + a_8)$$

$$\geq a_1 + a_2 + (a_4 + a_4) + (a_8 + a_8 + a_8 + a_8)$$

$$= a_1 + a_2 + 2a_4 + 4a_8$$

$$\geq \frac{1}{2} a_1 + a_2 + 2a_4 + 4a_8$$

$$= \frac{1}{2} (a_1 + 2a_2 + 4a_4 + 8a_8)$$

$$= \frac{1}{2} t_3.$$

where again we used that (a_n) is decreasing, and $a_n \ge 0$. Iterating, we obtain that for all $k \in \mathbb{N}$ it holds:

$$\begin{split} s_{2^{k}} &= a_{1} + a_{2} + (a_{3} + a_{4}) + (a_{5} + a_{6} + a_{7} + a_{8}) \\ &+ \dots + (a_{2^{k-1}} + \dots + a_{2^{k}}) \\ &\geq a_{1} + a_{2} + (a_{4} + a_{4}) + (a_{8} + a_{8} + a_{8} + a_{8}) \\ &+ \dots + (a_{2^{k}} + \dots + a_{2^{k}}) \\ &= a_{1} + a_{2} + 2a_{4} + 4a_{8} + \dots + 2^{k-1}a_{2^{k}} \\ &\geq \frac{1}{2}a_{1} + a_{2} + 2a_{4} + 4a_{8} + \dots + 2^{k-1}a_{2^{k}} \\ &= \frac{1}{2}\left(a_{1} + 2a_{2} + 4a_{4} + 8a_{8} + \dots + 2^{k}a_{2^{k}}\right) \\ &= \frac{1}{2}t_{k} \,. \end{split}$$

We have shown that

$$s_{2^k} \ge \frac{1}{2} t_k$$
, $\forall k \in \mathbb{N}$.

Since (t_k) is not bounded above, we infer that (s_{2^k}) is not bounded above. In particular (s_k) is not bounded, and hence divergent. Thus the series

$$\sum_{n=1}^{\infty} a_n$$

diverges. *Part2.* Suppose that the series

 $\sum_{n=0}^{\infty} 2^n a_{2^n}$

converges. Hence the sequence (t_k) converges, and therefore (t_k) is bounded. This means that there exists M > 0 such that

$$|t_k| \le M$$
, $\forall n \in \mathbb{N}$.

Since $t_k \ge 0$, the above reads

 $t_k \leq M$, $\forall n \in \mathbb{N}$.

Fix $k \in \mathbb{N}$ and let $m \in \mathbb{N}$ be such that

$$k \le 2^{m+1} - 1$$
.

In this way

$$s_k \leq s_{2^{m+1}-1}.$$

We have

$$\begin{split} s_{2^{m+1}-1} &= a_1 + (a_2 + a_3) + (a_4 + a_5 + a_6 + a_7) \\ &+ \dots + (a_{2^m} + \dots + a_{2^{m+1}-1}) \\ &\leq a_1 + (a_2 + a_2) + (a_4 + a_4 + a_4 + a_4) \\ &+ \dots + (a_{2^m} + \dots + a_{2^m}) \\ &= a_1 + 2a_2 + 4a_4 + \dots + 2^m a_{2^m} \\ &= t_m \,, \end{split}$$

where we used that (a_n) is decreasing. We have then shown

$$s_k \leq s_{2^{m+1}-1} \leq t_m \leq M.$$

Since M does not depend on k, we conclude that

$$s_k \leq M$$
, $\forall k \in \mathbb{N}$.

As $s_k \ge 0$, we conclude that s_k is bounded. Recalling that (s_k) is increasing, by the Monotone Convergence Theorem we infer that (s_k) converges. This proves that the series

$$\sum_{n=1}^{\infty} a_n$$

is convergent, ending the proof.

Thanks to the Cauchy Condensation Test of Theorem 8.26 we can prove the following result.

Theorem 8.27: Convergence of *p*-series

Let $p \in \mathbb{R}$. Consider the *p*-series

$$\sum_{n=1}^{\infty} \frac{1}{n^p} \, .$$

We have:

- 1. If p > 1 the *p*-series converges.
- 2. If $p \leq 1$ the *p*-series diverges.

Proof

The series in question is

$$\sum_{n=1}^{\infty} a_n \,, \quad a_n \,:= \frac{1}{n^p} \,.$$

Note that (a_n) is decreasing and non-negative. Hence, by the Cauchy Condensation Test of Theorem 8.26, the *p*-series converges if and only if

$$\sum_{n=0}^{\infty} 2^n a_{2^n}$$

converges. We have

$$\sum_{n=0}^{\infty} 2^n a_{2^n} = \sum_{n=0}^{\infty} 2^{n-np} = \sum_{n=0}^{\infty} (2^{1-p})^n,$$

and the latter is a Geometric Series of ratio

$$x := 2^{1-p}$$
.

By the Geometric Series Test, we have convergence if and only if

|x| < 1,

which is equivalent to

$$2^{1-p} < 1 = 2^0 \quad \Longleftrightarrow \quad 1-p < 0$$
$$\iff \quad p > 1.$$

Therefore

$$\sum_{n=1}^{\infty} \frac{1}{n^p}$$

converges if and only if p > 1, ending the proof.

The following is another notable application of the Cauchy Condensation Test.

Theorem 8.28

Let $p \in \mathbb{R}$. Consider the series

$$\sum_{n=2}^{\infty} \frac{1}{n \left(\log n\right)^p}.$$

We have:

- 1. If p > 1 the series converges.
- 2. If $p \leq 1$ the series diverges.

Proof

The series in question is

$$\sum_{n=2}^{\infty} a_n, \quad a_n := \frac{1}{n \left(\log n\right)^p}.$$

Note that (a_n) is non-negative and decreasing. Therefore we can apply the Cauchy Condensation Test to conclude that the above series is convergent if and only if the series

$$\sum_{n=1}^{\infty} 2^n a_{2^n}$$

is convergent. We have

$$2^{n}a_{2^{n}} = 2^{n} \frac{1}{2^{n} \left(\log 2^{n}\right)^{p}} = \frac{1}{n^{p} \log 2}$$

so that

$$\sum_{n=1}^{\infty} 2^n a_{2^n} = \frac{1}{\log 2} \sum_{n=1}^{\infty} \frac{1}{n^p}$$

The latter is a *p*-series, which by Theorem 8.27 converges if and only if p > 1. Hence

$$\sum_{n=2}^{\infty} \frac{1}{n \left(\log n\right)^{p}}$$

converges if and only if p > 1, and the proof is concluded.

8.4.2 Comparison Test

Another really useful result to study the convergence of non-negative series is the Comparison Test.

Theorem 8.29: Comparison test

Let $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ be non-negative sequences. Suppose that there exists $N \in \mathbb{N}$ such that

$$a_n \leq b_n$$
, $\forall n \geq N$.

They hold:

$$\sum_{n=1}^{\infty} b_n \text{ converges} \implies \sum_{n=1}^{\infty} a_n \text{ converges},$$
$$\sum_{n=1}^{\infty} a_n \text{ diverges} \implies \sum_{n=1}^{\infty} b_n \text{ diverges}.$$

Proof

Part 1. Define the partial sums starting at n = N

$$s_k := \sum_{n=N}^k a_k, \quad t_k := \sum_{n=N}^k b_k.$$

Suppose that

$$\sum_{n=1}^{\infty} b_n$$

converges. Hence also the series

$$\sum_{n=N}^{\infty} b_n$$

converges. Then (t_k) is a convergent sequence, which implies that (t_k) is bounded, and hence bounded above. Moreover, by assumption

$$s_k = a_N + a_{N+1} + \dots + a_k$$

 $\leq b_N + b_{N+1} + \dots + b_k$
 $= t_k$,

which reads

Therefore (s_k) is bounded above, being (t_k) bounded above. Recall that s_k is increasing, by Lemma 8.22. By the Monotone Convergence Theorem we conclude that s_k is convergent, showing that the series

 $s_k \leq t_k$, $\forall k \geq N$.



converges. Hence also the series

$$\sum_{n=1}^{\infty} a_n$$

converges, concluding the proof of Point 1.

Part 2. Note that Point 2 is the contrapositive of Point 1, and hence it holds.

Let us give two applications of the Comparison Test.

Example 8.30

Question. Discuss convergence/divergence of the following series:

$$\sum_{n=1}^{\infty} \frac{1}{n^2 + 3n - 1},$$
(8.9)

$$\sum_{n=0}^{\infty} \frac{3^n + 6n + \frac{1}{n+1}}{2^n} \,. \tag{8.10}$$

Solution.

1. Since $3n - 1 \ge 0$ for all $n \in \mathbb{N}$, we get

$$\frac{1}{n^2+3n-1} \le \frac{1}{n^2}, \quad \forall n \in \mathbb{N}.$$

By Theorem 8.27 the *p*-series

$$\sum_{n=1}^{\infty} \frac{1}{n^2}$$

converges. Therefore also the series at (8.9) converges by the Comparison Test in Theorem 8.29.

2. Note that

$$\frac{3^n + 6n + \frac{1}{n+1}}{2^n} \ge \frac{3^n}{2^n} = \left(\frac{3}{2}\right)^n, \quad \forall n \in \mathbb{N}.$$

Since $\left|\frac{3}{2}\right| = \frac{3}{2} > 1$, the series

diverges by the Geometric Series Test in Theorem 8.13. Therefore, by the Comparison Test, also the series at (8.10) diverges.

 $\sum_{n=0}^{\infty} \left(\frac{3}{2}\right)^n$

8.4.3 Limit Comparison Test

To apply the Comparison Test to the series

$$\sum_{n=1}^{\infty} a_n \, ,$$

one needs to find another sequence (b_n) such that

$$a_n \leq b_n$$
, $\forall n \geq N$,

or

$$b_n \leq a_n$$
, $\forall n \geq N$.

This is not always possible. However, one might be able to show that

$$\lim_{n \to \infty} \frac{a_n}{b_n} = L$$

for some $L \in \mathbb{R}$. In this case, the series of (a_n) and (b_n) can still be compared, in the sense specified in the Theorem below.

Theorem 8.31: Limit Comparison Test

Let (a_n) and (b_n) be sequences such that

$$a_n \ge 0$$
, $b_n > 0$, $\forall n \in \mathbb{N}$.

Suppose there exists $L \in \mathbb{R}$ such that

$$L = \lim_{n \to \infty} \frac{a_n}{b_n} \, .$$

They hold:

1. If $0 < L < \infty$, then

$$\sum_{n=1}^{\infty} a_n \text{ converges } \iff \sum_{n=1}^{\infty} b_n \text{ converges.}$$

2. If L = 0, then

$$\sum_{n=1}^{\infty} b_n \text{ converges} \implies \sum_{n=1}^{\infty} a_n \text{ converges},$$
$$\sum_{n=1}^{\infty} a_n \text{ diverges} \implies \sum_{n=1}^{\infty} b_n \text{ diverges}.$$

Proof

Part 1. Suppose that 0 < L < 1. Set

$$\varepsilon := \frac{L}{2}$$
.

Since $\varepsilon > 0$ and $a_n/b_n \to L$, there exists $N \in \mathbb{N}$ such that

$$\left|\frac{a_n}{b_n} - L\right| < \varepsilon \,, \quad \forall \, n \ge N \,.$$

The above is equivalent to

$$L - \varepsilon < \frac{a_n}{b_n} < \varepsilon + L$$
, $\forall n \ge N$.

Since $\varepsilon = L/2$, we get

$$\frac{L}{2} < \frac{a_n}{b_n} < \frac{3L}{2} \,, \quad \forall \, n \ge N \,,$$

or equivalently,

$$\frac{L}{2} b_n < a_n < \frac{3L}{2} b_n \,, \quad \forall \, n \ge N \,.$$

We are now ready to prove the main claim:

1. Suppose that

$$\sum_{n=1}^{\infty} a_n$$

converges. Then also

$$\sum_{n=N}^{\infty} a_n$$

converges, since we are only discarding a finite number of terms. As

$$\frac{L}{2}b_n \leq a_n\,, \quad \forall \, n \geq N\,,$$

it follows from the Comparison Test in Theorem 8.29 that the series

$$\sum_{n=N}^{\infty} \frac{L}{2} b_n \, .$$

converges. Since L/2 is a constant, we also conclude that

$$\sum_{n=1}^{\infty} b_n$$

converges.

2. Suppose that

$$\sum_{n=1}^{\infty} b_n$$

converges. Then also

$$\sum_{n=N}^{\infty} \frac{3L}{2} b_n$$

converges. Since

$$a_n < \frac{3L}{2} b_n \,, \quad \forall \, n \ge N \,,$$

by the Comparison Test we infer that

$$\sum_{n=N}^{\infty} a_n$$

converges. Therefore, also

$$\sum_{n=1}^{\infty} a_n$$

converges.

Part 2. Suppose that L = 0. Note that the second condition is the contrapositive of the first. Hence we only need to show that

$$\sum_{n=1}^{\infty} b_n \text{ converges } \Longrightarrow \quad \sum_{n=1}^{\infty} a_n \text{ converges.}$$

Let $\varepsilon = 1$. Since $a_n/b_n \to 0$, there exists $N \in \mathbb{N}$ such that

$$\left|\frac{a_n}{b_n} - 0\right| < \varepsilon = 1, \quad \forall \, n \ge N \, .$$

Therefore

 $a_n < b_n$, $\forall n \ge N$.

The thesis follows immediately by the Comparison Test in Theorem 8.29.

Important

It might happen that

$$\lim_{n \to \infty} \frac{a_n}{b_n} = \infty$$

In this case, we can still apply the Limit Comparison Test to the series

$$\sum_{n=1}^{\infty} \frac{1}{b_n/a_n}$$

since in this case

$$\lim_{n\to\infty}\frac{b_n}{a_n}=0$$

Let us give a few applications of the Limit Comparison Tets.

Example 8.32

Question. Prove that the following series converges

$$\sum_{n=1}^{\infty} \frac{2n^3 + 5n + 1}{7n^6 + 2n + 5}$$

Solution. Set

$$a_n := \frac{2n^3 + 5n + 1}{7n^6 + 2n + 5}, \quad b_n := \frac{1}{n^3}.$$

We have

$$L := \lim_{n \to \infty} \frac{u_n}{b_n}$$

= $\lim_{n \to \infty} \frac{2n^3 + 5n + 1}{7n^6 + 2n + 5} / \frac{1}{n^3}$
= $\lim_{n \to \infty} \frac{2n^6 + 5n^4 + n^3}{7n^6 + 2n + 5}$
= $\lim_{n \to \infty} \frac{2 + \frac{5}{n^2} + \frac{1}{n^3}}{7 + \frac{2}{n^5} + \frac{5}{n^6}} = \frac{2}{7}.$

The series

$$\sum_{n=1}^{\infty} \frac{1}{n^3}$$

converges, being a *p*-series with p = 3 > 1. Since $L = \frac{2}{7} > 0$, also the series

$$\sum_{n=1}^{\infty} \frac{2n^3 + 5n + 1}{7n^6 + 2n + 5}$$

converges, by the Limit Comparison Test.

Example 8.33

Question. Prove that the following series diverges

$$\sum_{n=1}^{\infty} \frac{n + \cos(n)}{n^2}$$

Solution. Since sin(n) is bounded, we expect the terms in the series to behave like 1/n for large n. Hence we set

$$a_n := \frac{n + \cos(n)}{n^2}, \quad b_n = \frac{1}{n}.$$

We compute

$$L := \frac{a_n}{b_n} = \lim_{n \to \infty} \frac{n + \cos(n)}{n^2} / \frac{1}{n}$$
$$= \lim_{n \to \infty} \frac{n^2 + n \cos(n)}{n^2}$$
$$= \lim_{n \to \infty} \left(1 + \frac{\cos(n)}{n}\right)$$

Note that

$$-1 \le \cos(n) \le 1 \implies -\frac{1}{n} \le \frac{\cos(n)}{n} \le \frac{1}{n}$$

As both $-\frac{1}{n} \to 0$ and $\frac{1}{n} \to 0$, by the Squeeze Theorem

$$\frac{\cos(n)}{n} \longrightarrow 0$$

Hence

$$L = \lim_{n \to \infty} \left(1 + \frac{\cos(n)}{n} \right) = 1.$$

The harmonic series $\sum_{n=1}^{\infty} \frac{1}{n}$ diverges. Since L = 1 > 0, the series

$$\sum_{n=1}^{\infty} \frac{n + \cos(n)}{n^2}$$

diverges by the Limit Comparison Test.

Example 8.34

Question. Prove that the following series converges

$$\sum_{n=1}^{\infty} \left(1 - \cos\left(\frac{1}{n}\right) \right) \, .$$

Solution. Since

$$\cos\left(\frac{1}{n}\right) \leq 1$$
,

the above is a non-negative series. Recall the limit

$$\lim_{n\to\infty}\frac{1-\cos(a_n)}{(a_n)^2}=\frac{1}{2}\,,$$

where (a_n) is a sequence in \mathbb{R} such that $a_n \to 0$ and

$$a_n \neq 0 \quad \forall n \in \mathbb{N}$$
.

In particular, for $a_n = 1/n$, we obtain

$$\lim_{n \to \infty} n^2 \left(1 - \cos\left(\frac{1}{n}\right) \right) = \frac{1}{2}.$$

Set

$$b_n := 1 - \cos\left(\frac{1}{n}\right), \quad c_n := \frac{1}{n^2}.$$

We have

$$L := \lim_{n \to \infty} \frac{b_n}{c_n} = \lim_{n \to \infty} n^2 \left(1 - \cos\left(\frac{1}{n}\right) \right) = \frac{1}{2}.$$

Note that the series $\sum_{n=1}^{\infty} \frac{1}{n^2}$ converges, being a *p*-series with p > 2. Therefore, since L = 1/2 > 0, also the series

$$\sum_{n=1}^{\infty} \left(1 - \cos\left(\frac{1}{n}\right) \right)$$

converges, by the Limit Comparison Test.

Sometimes the Limit Comparison Test fails, but the Comparison Test works.

Example 8.35

Question. Prove that the following series converges

$$\sum_{n=1}^{\infty} \frac{1+\sin(n)}{n^2}$$

Solution. Since

$$\sin(n)\geq -1\,,$$

the above is a non-negative series. As sin(n) is bounded, the series behaves similarly to

$$\sum_{n=1}^{\infty} \frac{1}{n^2} \, .$$

However

$$\frac{1+\sin(n)}{n^2} \bigg/ \frac{1}{n^2} = 1 + \sin(n)$$

does not converge. Hence, we cannot use the Limit Comparison Test. In alternative, we note that

$$\frac{1+\sin(n)}{n^2} \le \frac{2}{n^2}, \quad \forall n \in \mathbb{N}.$$

The series

$$\sum_{n=1}^{\infty} \frac{2}{n^2}$$

converges, being a *p*-series with p = 2 > 1. Therefore also

$$\sum_{n=1}^{\infty} \frac{1+\sin(n)}{n^2}$$

converges, by the Comparison Test of Theorem 8.29.

8.4.4 Ratio Test for positive series

The Ratio Test can be generalized to series. Notice that in this case the terms of the series need to be positive.

Theorem 8.36: Ratio Test for positive series

Let (a_n) be a sequence in \mathbb{R} such that

$$a_n > 0$$
, $\forall n \in \mathbb{N}$.

1. Suppose that the following limit exists:

$$L := \lim_{n \to \infty} \frac{a_{n+1}}{a_n} \, .$$

They hold:

- If L < 1 then ∑_{n=1}[∞] a_n converges.
 If L > 1 then ∑_{n=1}[∞] a_n diverges.
- 2. Suppose that there exists $N \in \mathbb{N}$ and L > 1 such that

$$\frac{a_{n+1}}{a_n} \ge L \,, \quad \forall \, n \ge N \,.$$

Then the series $\sum_{n=1}^{\infty} a_n$ diverges.

Proof

Part 1. Let

$$L := \lim_{n \to \infty} \frac{a_{n+1}}{a_n} \, .$$

• Suppose that L < 1. Therefore there exists *r* such that

$$L < r < 1.$$

Define

$$\varepsilon := r - L$$

so that $\varepsilon > 0$. By the convergence $a_{n+1}/a_n \to L$ there exists $N \in \mathbb{N}$ such that

$$\left|\frac{a_{n+1}}{a_n} - L\right| < \varepsilon = r - L, \quad \forall \, n \ge N \,.$$

In particular

$$\frac{a_{n+1}}{a_n} - L < r - L, \quad \forall n \ge N,$$

which implies

 $a_{n+1} < r a_n \,, \quad \forall \, n \ge N \,.$

Applying n - N times the above estimate we get

$$0 < a_n < r \, a_{n-1} < \dots < r^{n-N} \, a_N \,, \quad \forall \, n \ge N \,. \tag{8.11}$$

Note that the series of $r^{n-N} a_N$ converges, since

$$\sum_{n=N}^{\infty} r^{n-N} a_N = a_N \sum_{k=0}^{\infty} r^k = a_N \frac{1}{1-r} \,,$$

where the last equality follows because $\sum_{k=0}^{\infty} r^k$ is a geometric series and 0 < r < 1. Since (8.11) holds, by the Comparison Test in Theorem 8.29 we conclude that the series



converges. Therefore also the series

$$\sum_{n=1}^{\infty} a_n$$

converges, ending the proof in the case L < 1.

- Suppose L > 1: Then, by the Ratio Test for sequences, it follows that a_n diverges. Therefore

$$\lim_{n\to\infty}a_n\neq 0\,,$$

and the series $\sum_{n=1}^{\infty}$ diverges by the Necessary Condition, see Theorem 8.6.

Part 2. Suppose there exists L > 1 and $N \in \mathbb{N}$ such that

$$\frac{a_{n+1}}{a_n} \ge L, \quad \forall \, n \ge N \, .$$

By the Ratio Test for sequences it follows that a_n diverges. Therefore $\sum_{n=1}^{\infty}$ diverges by the Necessary Condition.

Example 8.37

Question. Discuss convergence/divergence of the following series

$$\sum_{n=1}^{\infty} a_n \, , \quad a_n = \frac{(n!)^2}{(2n)!} \, .$$

Solution. We compute

$$\lim_{n \to \infty} \frac{a_{n+1}}{a_n} = \lim_{n \to \infty} \frac{((n+1)!)^2}{(2(n+1))!} / \frac{(n!)^2}{(2n)!}$$
$$= \lim_{n \to \infty} \frac{(n+1)^2}{(2n+2)(2n+1)}$$
$$= \lim_{n \to \infty} \frac{\left(1 + \frac{1}{n}\right)^2}{\left(2 + \frac{2}{n}\right)\left(2 + \frac{1}{n}\right)} = \frac{1}{4}$$

Since L = 1/4 < 1, by the Ratio Test we conclude that $\sum a_n$ converges.

Important

Like with the Ratio Test for sequences, the case L = 1 is not covered by Theorem 8.36. This is because, in this case, the series

 $\sum_{n=1}^{\infty} a_n$

might be convergent or divergent, as shown in the next example.

Example 8.38

1. Consider the series

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

Setting $a_n = 1/n$, we have

$$L = \lim_{n \to \infty} \frac{a_{n+1}}{a_n}$$
$$= \lim_{n \to \infty} \frac{1}{n+1} / \frac{1}{n}$$
$$= \lim_{n \to \infty} \frac{n}{n+1} = 1.$$

Therefore L = 1 and we cannot apply the Ratio Test. However the series in question diverges, being the harmonic series.

2. Consider the series

$$\sum_{n=1}^{\infty} \frac{1}{n^2}$$

Setting $a_n = 1/n^2$, we have

$$L = \lim_{n \to \infty} \frac{a_{n+1}}{a_n}$$
$$= \lim_{n \to \infty} \frac{1}{(n+1)^2} / \frac{1}{n^2}$$
$$= \lim_{n \to \infty} \frac{n^2}{n^2 + 2n + 1} = 1$$

Therefore L = 1 and we cannot apply the Ratio Test. However the series in question diverges, being a *p*-series with p = 2 > 1.

The Ratio Test can often be combined with other convergence tests, as seen in the following example.

Example 8.39

Question. Using the Cauchy Condensation Test and the Ratio Test, prove that the following series converges

$$\sum_{n=1}^{\infty} \frac{\log(n)}{n^2} \, .$$

Solution. Set $a_n = \log n/n^2$. By the Cauchy Condensation Test, we know that $\sum a_n$ converges if and only if $\sum 2^n a_{2^n}$ converges. We have:

$$\sum_{n=0}^{\infty} 2^n a_{2^n} = \sum_{n=0}^{\infty} 2^n \frac{\log(2^n)}{(2^n)^2}$$
$$= \log(2) \sum_{n=0}^{\infty} \frac{n}{2^n}$$
$$= \log(2) \sum_{n=0}^{\infty} b_n, \qquad b_n := \frac{n}{2^n}$$

Apply the Ratio Test to the series $\sum b_n$

$$\frac{b_{n+1}}{b_n} = \frac{n+1}{2^{n+1}} \bigg/ \frac{n}{2^n} = \frac{n+1}{2n} \longrightarrow \frac{1}{2} < 1.$$

Therefore, $\sum b_n$ converges by the Ratio Test, so that also $\sum 2^n a_{2^n}$ converges. We conclude that $\sum a_n$ converges by the Cauchy Condensation Test.

8.5 General series

In the previous section we presented several tests for non-negative series. For non-negative series we showed that the partial sums (s_k) are increasing, see Lemma 8.22. This makes non-negative terms series relatively easy to study.

When a series contains both positive and negative terms, the partial sums (s_k) might oscillate, making the series harder to study. In this section we present some tests for general series in \mathbb{C} .

8.5.1 Absolute Convergence Test

To study general series, we introduce a stronger notions of convergence, known as **absolute con-vergence**.

Definition 8.40: Absolute convergence

Let (a_n) be a sequence in \mathbb{C} . The series $\sum_{n=1}^{\infty} a_n$ is said to **converge absolutely** if the following non-negative series converges

$$\sum_{n=1}^{\infty} |a_n| \; .$$

Let us show that absolute convergence implies convergence.

Theorem 8.41: Absolute Convergence Test

Let (a_n) be a sequence in \mathbb{C} . If the series $\sum_{n=1}^{\infty} a_n$ converge absolutely, then the series converges.

Before proceeding with the proof, we introduce some notation.

Notation 8.42: Positive and negative part

For a number $x \in \mathbb{R}$, we define

$$x^{+} := \begin{cases} x & \text{if } x \ge 0\\ 0 & \text{if } x < 0 \end{cases} \qquad x^{-} := \begin{cases} -x & \text{if } x < 0\\ 0 & \text{if } x \ge 0 \end{cases}$$

These are called the **positive part** and **negative part** of *x*, respectively. Note that

$$x^+ \ge 0, \quad x^- \ge 0.$$

Moreover they hold

$$x = x^+ - x^-$$
, $|x| = x^+ + x^-$.

The above relations are easy to check, and the proof is omitted. In particular, it holds that

$$x^+ \le |x|, \quad x^- \le |x|.$$

Proof: Proof of Theorem 8.41

Part 1. Suppose first that (a_n) is a sequence in \mathbb{R} such that

$$\sum_{n=1}^{\infty} |a_n|$$

converges. Since

$$0 \le a_n^+ \le |a_n| , \quad \forall \, n \in \mathbb{N} \,,$$

we can use the Comparison Test for non-negative series (Theorem 8.29) and conclude that the

series

$$\sum_{n=1}^{\infty} a_n^+ \, .$$

converges. Similarly, we have

$$0 \le a_n^- \le |a_n| , \quad \forall \, n \in \mathbb{N} \,.$$

Therefore also the series

$$\sum_{n=1}^{\infty} a_n^{-1}$$

converges by the Comparison Test. Since

$$a_n = a_n^+ - a_n^-, \quad \forall n \in \mathbb{N},$$

we can use the Algebra of Limits for series (Theorem 8.18) to conclude that

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} \left(a_n^+ - a_n^- \right) = \sum_{n=1}^{\infty} a_n^+ - \sum_{n=1}^{\infty} a_n^-$$

converges.

Part 2. Suppose now that (a_n) is a sequence in \mathbb{C} such that

$$\sum_{n=1}^{\infty} |a_n|$$

converges. Let $x_n, y_n \in \mathbb{R}$ denote the real and imaginary part of a_n . Therefore

$$|x_n| = \sqrt{x_n^2} \le \sqrt{x_n^2 + y_n^2} = |a_n|, \quad \forall n \in \mathbb{N}.$$

Therefore the series

$$\sum_{n=1}^{\infty} |x_n|$$

converges by the Comparison Test for non-negative series (Theorem 8.29). Since (x_n) is a real sequence, from Part 1 of the proof we have that the series

$$\sum_{n=1}^{\infty} x_n$$

converges. Arguing in the same way for the imaginary part y_n we conclude that also

$$\sum_{n=1}^{\infty} y_n$$

converges. Finally, by the Algebra of Limits in C, we get

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (x_n + iy_n) = \sum_{n=1}^{\infty} x_n + i \sum_{n=1}^{\infty} y_n \,,$$

proving that $\sum_{n=1}^{\infty} a_n$ converges.

Example 8.43

Question. Discuss absolute convergence of the series

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n} \, .$$

Solution. The series does not converge absolutely, since

$$\sum_{n=1}^{\infty} \left| (-1)^n \frac{1}{n} \right| = \sum_{n=1}^{\infty} \frac{1}{n}$$

does not converge, being the harmonic series.

Example 8.44

Question. Prove that the following series converges

$$\sum_{n=1}^{\infty} a_n \,, \qquad a_n = (-1)^n \frac{n^2 - 5n + 2}{n^4} \,.$$

Solution. We have

$$|a_n| = rac{|n^2 - 5n + 2|}{n^4} = rac{n^2 + 5n + 2}{n^4}$$

for *n* sufficiently large (e.g. $n \ge 10$). Note that

$$\frac{n^2 + 5n + 2}{n^4} \bigg/ \frac{1}{n^2} = \frac{n^4 + 5n^3 + 2n^2}{n^4}$$
$$= 1 + \frac{5}{n} + \frac{2}{n^2} \longrightarrow 1$$

The series $\sum 1/n^2$ converges, being a *p*-series with p = 2. Hence, also

$$\sum_{n=1}^{\infty} \frac{n^2 + 5n + 2}{n^4}$$

converges, by the Limit Comparison Test for non-negative series (Theorem 8.31). This shows $\sum |a_n|$ converges, which means that $\sum a_n$ converges absolutely. In particular, $\sum a_n$ converges by the Absolute Convergence Test.

8.5.2 Ratio Test for general series

As an application of the Absolute Convergence Test, we obtain the Ratio Test for general series.

Theorem 8.45: Ratio Test for general series

Let (a_n) be a sequence in \mathbb{C} , such that

$$a_n \neq 0 \quad \forall n \in \mathbb{N}$$
.

1. Suppose that the following limit exists:

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| \, .$$

They hold:

• If L < 1 then $\sum_{n=1}^{\infty} a_n$ converges absolutely, and hence converges.

• If L > 1 then $\sum_{n=1}^{\infty} a_n$ diverges.

2. Suppose that there exists $N \in \mathbb{N}$ and L > 1 such that

$$\left|\frac{a_{n+1}}{a_n}\right| \ge L\,, \quad \forall \, n \ge N\,.$$

Then the series $\sum_{n=1}^{\infty} a_n$ diverges.

Proof

Part 1. Let

$$b_n := |a_n| ,$$

so that

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lim_{n \to \infty} \frac{b_{n+1}}{b_n} \, .$$

• Suppose that L < 1. Since (b_n) is a sequence with non-negative terms, we have that

$$\sum_{n=1}^{\infty} b_n$$

converges by the Ratio Test for non-negative series, see Theorem 8.36. Since, by definition

$$\sum_{n=1}^{\infty} b_n = \sum_{n=1}^{\infty} |a_n|$$

also the latter series converges, i.e., $\sum_{n=1}^{\infty} a_n$ converges absolutely. In particular $\sum_{n=1}^{\infty} a_n$ converges, by the Absolute Convergence Test in Theorem 8.41.

• Suppose that L > 1. Then the sequence (a_n) diverges by the Ratio Test for sequences. Hence the series

$$\sum_{n=1}^{\infty} a_n$$

diverges by the Necessary Condition in Theorem 8.6.

Part 2. If there exists $N \in \mathbb{N}$ and L > 1 such that

$$\left|\frac{a_{n+1}}{a_n}\right| \ge L\,, \quad \forall \, n \ge N\,,$$

then the sequence (a_n) diverges by the Ratio Test for sequences, and we conclude as above.

Example 8.46

Question. Prove that the series converges

$$\sum_{n=1}^{\infty} a_n, \quad a_n = \frac{(4-3i)^n}{(n+1)!}$$

Solution. We have

$$L := \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right|$$

= $\lim_{n \to \infty} \left| \frac{(4-3i)^{n+1}}{((n+1)+1)!} \right| / \frac{(4-3i)^n}{(n+1)!} \right|$
= $\lim_{n \to \infty} \frac{5}{n+2} = 0.$

As L = 0 < 1, we conclude that $\sum a_n$ converges absolutely, by the Ratio Test. Hence, $\sum a_n$ converges by the Absolute Convergence Test.

8.5.3 Exponential function and Euler's Number

We have already encountered the exponential function in the Euler's identity

$$e^{i\theta} = \cos(\theta) + i\sin(\theta), \quad \theta \in \mathbb{R}.$$

We have also defined the exponential of $z = a + ib \in \mathbb{C}$ as

$$e^z = e^a e^{ib} \,.$$

We have also anticipated that

$$e^z = \sum_{n=0}^{\infty} \frac{z^n}{n!} \, .$$

Using the Ratio Test for general series, we can give a precise meaning to the above expression. We start by studying convergence of the exponential series.

Theorem 8.47: Exponential series

Let $z \in \mathbb{C}$. The **exponential series**

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges absolutely.

Proof

Set $a_n = z^n/n!$. Then

$$L = \lim_{n \to \infty} \left| \frac{a_{n+1}}{a_n} \right|$$
$$= \lim_{n \to \infty} \left| \frac{z^{n+1}}{(n+1)!} / \frac{z^n}{n!} \right|$$
$$= \lim_{n \to \infty} \frac{|z|}{n+1} = 0.$$

Therefore the series converges absolutely by the Ratio Test in Theorem 8.45.

Definition 8.48: Exponential function

Define the exponential function

$$\exp: \mathbb{C} \to \mathbb{C} \,, \quad \exp(z) \,:= \sum_{n=0}^{\infty} \frac{z^n}{n!}$$

for $z \in \mathbb{C}$. We denote

 $e^z := \exp(z), \quad e := e^1.$

Remark 8.49

1. Using the definition of e^z , one can show that

$$e^{z+w} = e^z e^w, \quad (e^z)^w = e^{zw},$$

for all $z, w \in \mathbb{C}$.

2. This way, we see that the new definition of exponential agrees with the old one:

$$e^z = e^{a+ib} = e^a e^{ib} \,.$$

3. We had also defined

$$e := \lim_{n \to \infty} \left(1 + \frac{1}{n} \right)^n \, .$$

Using the binomial theorem one can prove that

$$\lim_{n \to \infty} \left(1 + \frac{1}{n} \right)^n = \sum_{n=0}^{\infty} \frac{1}{n!}$$

8.5.4 Conditional convergence

Some series do not converge absolutely, but still converge. Such series are said to converge conditionally.

Definition 8.50: Conditional convergence

Let (a_n) be a sequence in \mathbb{C} . We say that the series

$$\sum_{n=1}^{\infty} a_n$$

converges conditionally if it converges, but it does not converge absolutely.

In practice, conditional convergence means that the convergence of the series depends on the order in which we perform the summation. Changing the order of summation of a series is called rearrangement.

Definition 8.51: Rearrangement of a series

Let (a_n) be a sequence in \mathbb{C} . Then:

- 1. A **permutation** is a bijection $\sigma : \mathbb{N} \to \mathbb{N}$.
- 2. A **rearrangement** of the series $\sum_{n=1}^{\infty} a_n$ is a series

$$\sum_{n=1}^{\infty} a_{\sigma(n)}$$

for some permutation σ .

If a series of complex numebers converges absolutely, then all its rearrangements converge to the same limit.

Theorem 8.52

Let (a_n) be a sequence in \mathbb{C} such that

$$\sum_{n=1}^{\infty} |a_n|$$

converges. For any permutation σ we have

$$\sum_{n=1}^{\infty} a_{\sigma(n)} = \sum_{n=1}^{\infty} a_n \, .$$

For a proof, see Theorem 3.55 in [3]. A very surprising result is the following: If a series of real numbers converges conditionally, then the series can be rearranged to converge to any real number.

Theorem 8.53: Riemann rearrangement Theorem

Let (a_n) be a real sequence such that the series

$$\sum_{n=1}^{\infty} a_r$$

converges conditionally. Let

$$L \in \mathbb{R}$$
 or $L = \pm \infty$.

There exists a permutation σ such that the corresponding rearrangement $\sum_{n=1}^{\infty} a_{\sigma(n)}$ converges conditionally to *L*, that is,

$$\sum_{n=1}^{\infty} a_{\sigma(n)} = L$$
For a proof, we refer the reader to Theorem 3.54 in [3].

In simpler terms, the last two Theorems are saying that:

- 1. If a series converges absolutely, then all the rearrangements converge to the same number:
 - Absolute convergence \implies Commutativity holds for infinite sums
- 2. If a series converges conditionally, then we can rearrange the terms to converge to any real number, or even diverge:
 - Conditional convergence \implies Commutativity does not hold for infinite sums

8.5.5 Dirichlet and Alternating Series Tests

There are very few conditional convergence tests available. We present the Dirichlet Test and the Alternating Series Test.

Theorem 8.54: Dirichlet Test

Let (c_n) be a sequence in \mathbb{C} and (q_n) a sequence in \mathbb{R} . Suppose that

- q_n is decreasing,
- $q_n \rightarrow 0$,
- $q_n \ge 0$ for all $n \in \mathbb{N}$.
- Suppose there exists M > 0 such that

$$\left|\sum_{n=1}^k c_n\right| \le M, \quad \forall k \in \mathbb{N}.$$

Then the following series converges

$$\sum_{n=1}^{\infty} c_n q_n \, .$$

Proof

Define the partial sums

$$s_k := \sum_{n=1}^k c_n, \quad \forall k \in \mathbb{N}.$$

By assumption it holds

$$|s_k| \le M, \quad \forall k \in \mathbb{N}.$$
 (8.12)

Note that

$$c_1 = s_1$$

$$c_2 = s_2 - s_1$$

$$\dots$$

$$c_n = s_n - s_{n-1}$$

Therefore

$$\sum_{n=1}^{k} c_k q_k = c_1 q_1 + c_2 q_2 + \dots + c_k q_k$$

= $s_1 q_1 + (s_2 - s_1) q_2 + \dots + (s_k - s_{k-1}) q_k$
= $s_1 (q_1 - q_2) + s_2 (q_2 - q_3) + \dots + s_{k-1} (q_{k-1} - q_k) + s_k q_k$
= $\left(\sum_{n=1}^{k-1} s_n (q_n - q_{n+1})\right) + s_k q_k$

Since (s_k) is bounded and $q_k \rightarrow 0$, we conclude that

 $s_k q_k \to 0$.

Further, notice that

 $q_n-q_{n+1}\geq 0\,,$

since q_n is decreasing. Therefore, using (8.12), we get

$$|s_n||q_n - q_{n+1}| = |s_n|(q_n - q_{n+1})$$

$$\leq M(q_n - q_{n+1})$$

for all $n \in \mathbb{N}$. Note that

$$\sum_{n=1}^{k-1} M(q_n - q_{n+1}) = M(q_1 - q_k).$$

Since $q_k \to 0$ as $k \to \infty$, we conclude that

$$\sum_{n=1}^{\infty} M(q_n - q_{n+1}) = Mq_1$$

Hence, by the Comparison Test for non-negative series, we infer that

$$\sum_{n=1}^{\infty} |s_n| |q_n - q_{n+1}|.$$

In particular the series

$$\sum_{n=1}^{\infty} s_n (q_n - q_{n+1})$$

converges by the Absolute Convergence Test. Since we have shown

$$\sum_{n=1}^{k} c_n q_n = \left(\sum_{n=1}^{k-1} s_n (q_n - q_{n+1})\right) + s_k q_k$$

and $s_k q_k \to 0$, we conclude that $\sum_{n=1}^{\infty} c_n q_n$ converges.

Example 8.55

Question. Let $\theta \in \mathbb{R}$, with

$$\theta \neq 2k\pi$$
, $\forall k \in \mathbb{Z}$.

Prove that the below series are conditionally convergent

$$\sum_{n=1}^{\infty} \frac{e^{i\theta n}}{n}, \quad \sum_{n=1}^{\infty} \frac{\cos(\theta n)}{n}, \quad \sum_{n=1}^{\infty} \frac{\sin(\theta n)}{n}$$

Solution.

1. Recalling the Euler's Identity

$$e^{i\theta} = \cos(\theta) + i\sin(\theta),$$

we obtain that

$$\sum_{n=1}^{\infty} \frac{e^{i\theta n}}{n} = \sum_{n=1}^{\infty} \frac{\cos(n\theta)}{n} + i \sum_{n=1}^{\infty} \frac{\sin(n\theta)}{n} \,.$$

Therefore, the series $\sum e^{i\theta n}/n$ converge conditionally if and only if $\sum \cos(\theta n)/n$ and $\sum \sin(\theta n)/n$ converge conditionally. It is then sufficient to study $\sum e^{i\theta n}/n$.

2. The series $\sum e^{i\theta n}/n$ does not converge absolutely, since

$$\sum_{n=1}^{\infty} \left| \frac{e^{i\theta n}}{n} \right| = \sum_{n=1}^{\infty} \frac{1}{n}$$

diverges, being the Harmonic Series.

3. Set $c_n = e^{i\theta n}$, $q_n = 1/n$, so that

$$\sum_{n=1}^{\infty} \frac{e^{i\theta n}}{n} = \sum_{n=1}^{\infty} c_n q_n \, .$$

We have that q_n is decreasing, $q_n \to 0$ and $q_n \ge 0.$ Let us prove that there exists M > 0 such that

$$\left|\sum_{n=1}^{k} e^{i\theta n}\right| \le M, \quad \forall k \in \mathbb{N}.$$
(8.13)

Note that

 $1-e^{i\theta}\neq 0$,

since $\theta \neq 2k\pi$ for all $k \in \mathbb{Z}$. Therefore we can use the Geometric Series (truncated) summation formula to get

$$\sum_{n=1}^{k} e^{i\theta n} = \sum_{n=1}^{k} (e^{i\theta})^n$$
$$= \frac{1 - e^{i(k+1)\theta}}{1 - e^{i\theta}} - 1$$
$$= e^{i\theta} \frac{1 - e^{ik\theta}}{1 - e^{i\theta}}$$

Taking the modulus

$$\begin{split} \sum_{n=1}^{k} e^{i\theta n} \Bigg| &= \left| e^{i\theta} \frac{1 - e^{ik\theta}}{1 - e^{i\theta}} \right| = \left| e^{i\theta} \right| \left| \frac{1 - e^{ik\theta}}{1 - e^{i\theta}} \right| \\ &= \frac{\left| 1 - e^{ik\theta} \right|}{\left| 1 - e^{i\theta} \right|} \le \frac{\left| 1 \right| + \left| e^{ik\theta} \right|}{\left| 1 - e^{i\theta} \right|} = \frac{2}{\left| 1 - e^{i\theta} \right|} \,, \end{split}$$

where we used the triangle inequality. Since the RHS does not depend on k, we can set

$$M = \frac{2}{|1 - e^{i\theta}|},$$

so that (8.13) holds. Therefore, $\sum e^{i\theta n}/n$ converges by the Dirichlet Test.

4. We have shown that $\sum e^{i\theta n}/n$ converges, but not absolutely. Hence, it converges conditionally.

As a corollary of the Dirichlet Test we obtain the Alternate Convergence Test.

Theorem 8.56: Alternate Convergence Test

Let (q_n) be a sequence in \mathbb{R} such that

• q_n is decreasing,

- $q_n \rightarrow 0$,
- $q_n \ge 0$ for all $n \in \mathbb{N}$.

The following series converges

$$\sum_{n=1}^{\infty} (-1)^n q_n$$

Proof

Define the sequence $c_n := (-1)^n$. Then,

$$\sum_{n=1}^{k} c_n = \begin{cases} 0 & \text{if } k \text{ even} \\ -1 & \text{if } k \text{ odd} \end{cases}$$

Hence

$$\left|\sum_{n=1}^k c_n\right| \le 1, \quad \forall \, k \in \mathbb{N}.$$

By the Dirichlet Test we have convergence of

$$\sum_{n=1}^{\infty} c_n q_n = \sum_{n=1}^{\infty} (-1)^n q_n \, .$$

Example 8.57

Question. Prove that the series converges conditionally

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n} \, .$$

Solution. The series does not converge absolutely, since

$$\sum_{n=1}^{\infty} \left| (-1)^n \frac{1}{n} \right| = \sum_{n=1}^{\infty} \frac{1}{n}$$

diverges, being the Harmonic Series. Set $q_n = 1/n$, so that

$$\sum_{n=1}^{\infty} (-1)^n \frac{1}{n} = \sum_{n=1}^{\infty} (-1)^n q_n \, .$$

Clearly, $q_n \ge 0$, $q_n \to 0$ and q_n is decreasing. Hence, the series converges by the Alternating Series Test. Thus, the series converges conditionally.

8.5.6 Abel's Test

The Abel Test is another test for conditional convergence. It looks similar to the Dirichlet Test, however notice that the Abel Test only deals with real sequences.

Theorem 8.58: Abel's Test

Let (a_n) and (q_n) be sequences in \mathbb{R} . Suppose that

- q_n is monotone and bounded,
- The series $\sum a_n$ converges.

Then the following series converges

$$\sum_{n=1}^{\infty} a_n q_n \, .$$

The proof is similar to the one of the Dirichlet Test. We decided to omit it.

Example 8.59

Question. Prove that the series converges conditionally

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{n} \left(1 + \frac{1}{n}\right)^n$$

Solution. Set

$$a_n := rac{(-1)^n}{n}, \quad q_n := \left(1 + rac{1}{n}\right)^n.$$

We have seen that q_n is monotone increasing and bounded (recall that $q_n \to \varepsilon$). Moreover, the series $\sum_{n=1}^{\infty} a_n$ converges by the Alternating Series Test, as seen in Example 8.57. Hence the series $\sum_{n=1}^{\infty} a_n q_n$ converges by the Abel Test.

However, the series in question does not converge absolutely. Indeed,

$$\left|\frac{(-1)^n}{n}\left(1+\frac{1}{n}\right)^n\right| = \frac{1}{n}q_n \ge \frac{1}{n}q_1 = \frac{2}{n},$$

since (q_n) is increasing. As the series $\sum 2/n$ diverges, by the Comparison Test we conclude that also

$$\sum_{n=1}^{\infty} \left| \frac{(-1)^n}{n} \left(1 + \frac{1}{n} \right)^n \right|$$

diverges. Therefore, the series in the example converges conditionally.

License

Reuse

This work is licensed under CC-BY-NC-ND 4.0



Citation

For attribution, please cite this work as:

```
Fanzon, Silvio. (2024). Lecture Notes of Numbers Sequences and Series. 
https://www.silviofanzon.com/2024-NSS-Notes/
```

BibTex citation:

```
@electronic{Fanzon-NSS-2024,
    author = {Fanzon, Silvio},
    title = {Lecture Notes of Numbers Sequences and Series},
    url = {https://www.silviofanzon.com/2024-NSS-Notes/},
    year = {2024}}
```

References

- [1] S. Abbott. Understanding Analysis. Second Edition. Springer, 2015.
- [2] Bartle, Robert G. and Sherbert, Donald R. *Introduction to Real Analysis*. Fourth Edition. Wiley, 2011.
- [3] W. Rudin. Principles of Mathematical Analysis. Third Edition. McGraw Hill, 1976.